

Contributors

Stephanie Allen, *School of Pharmacy, University of Nottingham, University Park, Nottingham NG7 2RD, UK.*

G. Michael Blackburn, *Department of Chemistry, University of Sheffield, Brook Hill, Sheffield S3 7HF, UK.*

Martin Egli, *Department of Biochemistry, Vanderbilt University, School of Medicine, Nashville, TN 37232, USA.*

Julie Fisher, *School of Chemistry, University of Leeds, Woodhouse Lane, Leeds LS2 9JT, UK.*

Andrew J. Flavell, *Plant Research Unit, University of Dundee at SCRI, Invergowrie, Dundee DD2 5DA, UK.*

Michael J. Gait, *MRC Laboratory of Molecular Biology, Hills Road, Cambridge CB2 2QH, UK.*

Ihtshamul Haq, *Department of Chemistry, University of Sheffield, Brook Hill, Sheffield S3 7HF, UK.*

Charles Laughton, *School of Pharmacy, University of Nottingham, University Park, Nottingham NG7 2RD, UK.*

David Loakes, *MRC Laboratory of Molecular Biology, Hills Road, Cambridge CB2 2QH, UK.*

Ben Luisi, *Department of Biochemistry, University of Cambridge, 80 Tennis Court Road, Cambridge CB2 1GA, UK.*

Anna Marie Pyle, *Department of Molecular Biophysics and Biochemistry, Yale University, 266 Whitney Avenue, P.O. Box 208114, New Haven, CT 06520-8114, USA.*

Elliott Stollar, *The Hospital for Sick Children Research Institute, 555 University Avenue, Toronto, Ont., Canada M5G 1X8.*

David M. Williams, *Department of Chemistry, University of Sheffield, Brook Hill, Sheffield S3 7HF, UK.*

Nicholas H. Williams, *Department of Chemistry, University of Sheffield, Brook Hill, Sheffield S3 7HF, UK.*

