

Electronic Supplementary Information

Facile Synthesis and Photovoltaic Applications of a New Alkylated Bismethano Fullerene as Electron Acceptor for High Open Circuit Voltage Solar Cells

Derya Baran^a, Sule Erten-Ela^{*b,d}, Andreas Kratzer^b, Tayebah Ameri^a, Christoph Brabec^{a,c},
Andreas Hirsch^{*b}

^a *Institute of Materials for Electronics and Energy Technology, Department of Materials Science and Engineering, Friedrich-Alexander-University Erlangen-Nurnberg, Martensstrasse 7, 91058, Erlangen, Germany*

^b *Department of Chemistry and Pharmacy, Interdisciplinary Center of Molecular Materials (ICMM), Friedrich-Alexander- Universität Erlangen-Nürnberg, Henkestrasse 42, 91054 Erlangen, Germany*

^c *Bavarian Center for Applied Energy Research (ZAE Bayern), Haberstrasse 2, 91058, Erlangen, Germany*

^d *Solar Energy Institute, Ege University, Bornova, 35100 Izmir, Turkey Phone: 0090 232 3111231; E-mail: suleerten@yahoo.com*





Fig. S1. HPLC gel-permeation separation of bis-4-nonyl [6, 6] methanofullerene




Fig. S2. Mass data of of bis-4-propylpentyl [6, 6] methanofullerene




Figure S3. Measured space-charge limited J - V characteristics of the NCBA and P3HT: NCBA blend devices under dark conditions for a) hole-only b) electron-only devices.