

Table S1

Individual recoveries (RSD, %) in six real samples spiked at 100 $\mu\text{g g}^{-1}$.

Compound	Li1	Li2	MC1	HG	Su1	NP	Mean recovery
PBSA	82.7	75.0	75.0 (10)	90.8 (8.0)	-	85.0	81.7 (6.8)
BP4	88.2	107 (1.6)	93.7 (8.6)		96.2 (9.6)	68.6 (3.7)	96.2 (7.8)
BP2	99.8	111 (2.3)	72.5 (6.7)	106 (9.8)	96.7 (8.0)	85 (14)	99.8 (9.8)
BP1	91.6	77.3 (6.2)	92 (15)	98.3 (7.8)	86.0 (6.3)	-	89.0 (7.9)
BP8	96.1	104 (2.8)	95.0 (7.0)	66.8 (8.5)	112 (3.1)	105 (1.9)	102 (6.9)
Eto	95.0	112 (4.4)	84.2 (4.8)	112 (8.4)	112 (2.4)	88.9 (3.7)	101 (13)
BP6	98.6	105 (0.1)	105 (6.9)	86.6 (6.6)	95.4 (5.6)	88.3 (0.8)	96.5 (7.9)
BP3	96.6	108 (2.0)	105 (6.9)	107 (3.8)	83.6 (1.5)	83.4 (5.4)	97 (11)
IAMC	103	86.4 (5.2)	104 (5.3)	103 (7.8)	100 (1.5)	91.6 (0.7)	97.9 (7.2)
4MBC	101	97.9 (1.9)	90.4 (8.3)	101 (7.6)	92.4 (1.3)	105 (0.8)	98.0 (5.6)
OCR	103	86.3 (5.4)	104 (7.3)	92.6 (3.5)	-	91.8 (0.2)	95.6 (7.7)
MA	97.4	104 (1.6)	102 (8.9)	105 (9.4)	94.2 (0.9)	105 (0.3)	101 (4.5)
EHPABA	93.2	104 (2.0)	88.2 (6.1)	94.2 (4.8)	73.6 (0.7)	84.1 (1.0)	90 (10)
2EHMC	99.9	71.2 (3.6)	96.5 (8.1)	94.1 (4.3)	77.3 (1.9)	93.0 (2.3)	89 (12)
BMDM	99.2	79.5 (6.7)	95.4 (7.3)	63.5 (4.4)	-	109 (0.8)	96 (12)

Li: lipstick; MC: moisturizing cream; HG: hair gel; Su: sunscreen; NP: nail polish.