

## Supplementary Information

### **Microfluidic Continuum Sorting of Sub-Populations of Tumor Cells via Surface Antibody Expression Levels**

Rhonda M. Jack, Khadijah Hussain, Danika Rodrigues, Ebrahim Azizi, Max Wicha, Diane M. Simeone, and Sunitha Nagrath\*

#### Contents

**Figure S1:** Cancer Cell magnetic bead coverage as a function of incubation times


**Figure S2:** Recoveries of cells with  $> 2/3$  bead attachment on the cell surface

**Figure S3:** EpCAM bead attachment on PANC-1 cells cultured for six days after sorting according to low, moderate and high EpCAM expression


**Figure S4:** Magnetic Flux normal to contour along channel

**Figure S5:** Magnetic Force experienced by  $1\mu\text{m}$  particle along channel


**Figure S6:** Magnetic Force experienced by  $1\mu\text{m}$  particle just before exiting the sorter channel




**Figure S1:** Cancer Cell magnetic bead coverage as a function of incubation times using both PANC-1 and MiAPaca cell lines and EpCAM and CD44 protein-coated magnetic beads. Times of 5, 10, 15 and 30min were evaluated for each cell line using either type of bead.




**Figure S2:** Recoveries of cells with > 2/3 bead attachment on the cell surface, as a function of magnetic sorter gap width




**Figure S3:** EpCAM bead attachment on PANC-1 cells cultured for six days after sorting according to low, moderate and high EpCAM expression then relabeled with EpCAM beads.



**Figure S4:** Magnetic Flux normal to contour along channel for the four different separation widths studied. Plots were generated using FEMM.



**Figure S5:** Magnetic Force experienced by 1 μm particle along channel, moving from inlets to outlets, for the four different separation widths studied. Plots were generated using FEMM. Red line indicates channel exit point, plotted in figure S6.



**Figure S6:** Magnetic Force experienced by 1μm particle just before exiting the sorter channel, for the four different separation widths studied. Plots were generated using FEMM.