

Supplementary

Table 6. Coagulation Function in different groups at the end of treatment and recovery period. ($\bar{X} \pm S$)

Groups	Gender	Treatment Period d35, (n=10)		Recovery Period d49, (n=5)	
		PT(s)	aPTT(s)	PT(s)	aPTT(s)
I	♂	21.1±1.9	13.9±1.8	22.3±3.7	13.7±2.8
	♀	21.7±2.2	12.4±1.2	18.7±3.5	12.3±2.4
II	♂	21.8±2.3	13.7±1.6	21.8±2.0	14.4±2.2
	♀	22.3±2.7	12.3±1.3	18.9±3.5	13.4±2.1
III	♂	20.6±1.6	13.8±2.0	21.5±3.7	14.3±1.4
	♀	22.2±1.7	12.4±1.4	19.9±2.9	12.4±1.1
IV	♂	20.2±2.9	13.9±2.0	19.9±3.9	14.1±3.5
	♀	21.8±2.1	12.7±1.8	20.0±2.5	13.2±2.2
V	♂	22.1±3.1	12.9±2.0	21.0±2.1	13.0±1.8
	♀	21.6±2.3	12.4±1.4	21.3±3.4	12.4±1.5
VI	♂	21.7±2.5	13.0±2.3	21.7±1.4	12.7±2.6
	♀	23.4±2.1	11.2±1.2	20.0±3.4	12.4±1.5

