

Supporting Information

Lysophosphatidylcholine modulates the aggregation of human islet amyloid polypeptide

Yanting Xing,[†] Emily H. Pilkington,[‡] Miaoyi Wang,[‡] Cameron J. Nowell,[¶] Aleksandr Kaminen,[‡] Yunxiang Sun,[†] Bo Wang,[†] Thomas P. Davis,^{‡§} Feng Ding^{†*} and Pu Chun Ke^{‡*}*

[†]Department of Physics and Astronomy, Clemson University, Clemson, SC 29634, USA

[‡]ARC Centre of Excellence in Convergent Bio-Nano Science and Technology, Monash Institute of Pharmaceutical Sciences, Monash University, 381 Royal Parade, Parkville, VIC 3052, Australia

[¶]Monash Institute of Pharmaceutical Sciences, Monash University, 381 Royal Parade, Parkville, VIC 3052, Australia

[§]Department of Chemistry, University of Warwick, Gibbet Hill, Coventry, CV4 7AL, United Kingdom

Supplementary Figures




Figure S1 TEM image of LPC micelles at 2 mM (>CMC). Scale: 10 nm.




Figure S2 Normalized far UV circular dichroism spectra of IAPP (25 μM) in the presence of LPC above and below its critical micelle point (CMC) over 24 h.




Figure S3 The IAPP amyloid fibril model from by two u-shaped protofibrils. Each peptide in the fibril is shown as cartoon with sidechains in stick representation.