

Supplementary information

A synthetic dodecanolide library for the identification of putative semiochemicals emitted by mantellid frogs

Pardha Saradhi Peram,^a Miguel Vences^b and Stefan Schulz^{a,*}

^aInstitut für Organische Chemie, Technische Universität Braunschweig, Hagenring 30, 38106 Braunschweig, Germany.

^bZoologisches Institut, Technische Universität Braunschweig, Mendelssohnstraße 4, 38106 Braunschweig, Germany

Contents

1. Mass spectra of diastereomers of synthesized compounds and comparison with natural compounds	2
2. ¹ H-NMR and ¹³ C-NMR spectra of intermediates leading to 2-methyl-11-dodecanolide (7)	3-8
3. ¹ H-NMR and ¹³ C-NMR spectra of intermediates leading to 4-methyl-11-dodecanolide (8)	9-12
4. ¹ H-NMR and ¹³ C-NMR spectra of intermediates leading to 6-methyl-11-dodecanolide (9)	13-19
5. ¹ H-NMR and ¹³ C-NMR spectra of intermediates leading to 8-methyl-11-dodecanolide (10)	20-25
6. ¹ H-NMR and ¹³ C-NMR spectra of intermediates leading to 10-methyl-11-dodecanolide (11)	26-30

1. Mass spectra of diastereomers of synthesized compounds and comparison with natural compounds










Fig. S1 Mass spectra and gas chromatographic retention indices of natural compounds (left) and synthetic compounds (right).




2. ^1H -NMR and ^{13}C -NMR spectra of intermediates leading to 2-methyl-11-dodecanolide (7)








30











32









7








3. ^1H -NMR and ^{13}C -NMR spectra of intermediates leading to 4-methyl-11-dodecanolide (8)








22











8












4. ^1H -NMR and ^{13}C -NMR spectra of intermediates leading to 6-methyl-11-dodecanolide (9)

















18











9









5. ^1H -NMR and ^{13}C -NMR spectra of intermediates leading to 8-methyl-11-dodecanolide (10)















28











10








6. ^1H -NMR and ^{13}C -NMR spectra of intermediates leading to 10-methyl-11-dodecanolide (11)








36





11

