

SUPPORTING INFORMATION FOR
Diplatinum, Triplatinum, and Tetraplatinum Complexes Featuring
 $\text{Cl}[\text{PtC}\equiv\text{CC}\equiv\text{CC}\equiv\text{CC}\equiv\text{C}]_m\text{PtCl}$ Segments; Iterative Syntheses and
Functionalization for Single Molecule Conductivity Studies

**Qinglin Zheng,[‡] Jakob Schneider,[‡] Hashem Amini,^{✉†} Frank Hampel,[‡] and John A.
Gladysz^{✉†*}**

Department of Chemistry, Texas A&M University, PO Box 30012, College Station, Texas
77842-3012, USA and Institut für Organische Chemie and Interdisciplinary Center for Molecular
Materials, Friedrich-Alexander-Universität Erlangen-Nürnberg, Henkestraße 42, 91054
Erlangen, Germany




Figure s1. Cyclic voltammograms used for the data in Table 7. Conditions: $3\text{-}11 \times 10^{-4}$ M in substrate and 0.10 M in $n\text{-Bu}_4\text{N}^+ \text{BF}_4^-$ in CH_2Cl_2 at 22.5 ± 1 °C; Pt working and counter electrodes, potential vs. Ag wire pseudoreference; scan rate 100 mV/s, calibrated vs. added ferrocene = 0.46 V.