

Supporting Information

Heteroleptic iron(II) complexes with naphthoquinone-type ligands

Takuya Shiga,^{*a} Rina Kumamaru,^a

Graham N. Newton^b and Hiroki Oshio^{a, c}

- a. Graduate School of Pure and Applied Sciences, University of Tsukuba, Tennodai 1-1-1, Tsukuba, Ibaraki 305-8571, Japan.
- b. GSK Carbon Neutral Laboratories for Sustainable Chemistry, The University of Nottingham, Nottingham NG7 2TU, U.K.
- c. State Key Laboratory of Fine Chemicals, Dalian University of Technology
2 Linggong Rd., 116024 Dalian, China

Corresponding author

Dr. Takuya Shiga

Graduate School of Pure and Applied Sciences,

University of Tsukuba

Tennodai 1-1-1, Tsukuba, Ibaraki 305-8571 (Japan)

TEL: (+81)29-852-5923

FAX: (+81)29-852-4426

E-mail:

shiga@chem.tsukuba.ac.jp

Contents

Figure S1 One dimensional network structure of **1**. Blue dotted lines represent intrachain hydrogen bond interactions.

Figure S2 One dimensional network structure of **2**. Blue dotted lines represent intrachain hydrogen bond interactions.

Figure S3. The derivative of the $\chi_m T$ data for complexes **1** (left) and **2** (right).

Figure S4. Simulation of magnetic data with spin crossover transition for complexes **1** (left) and **2** (right).

Figure S5 Mössbauer spectra of **1** at 300 K (top) and 20 K (bottom). Parameters are provided in Table S3.

Figure S6 Mössbauer spectra of **2** at 300 K (top) and 20 K (bottom). Parameters are provided in Table S3.

Figure S7 Mössbauer spectra of **3** (top) and **4** (bottom) at 20 K. Parameters are provided in Table S3.

Figure S8 Mössbauer spectra of **5** (top) and **6** (bottom) at 20 K. Parameters are provided in Table S3.

Table S1 Crystal parameters for **1** and **2**.

Table S2 Crystal parameters of **3**, **4**, **5** and **6** at 100 K.

Table S3 Mössbauer parameters for all complexes.

Explanation of alerts:




Figure S1. One dimensional network structure of **1**. Blue dotted lines represent intrachain hydrogen bond interactions.




Figure S2. One dimensional network structure of **2**. Blue dotted lines represent intrachain hydrogen bond interactions.




Figure S3. The derivative of the $\chi_m T$ data for complexes **1** (left) and **2** (right).




Figure S4. Simulation of magnetic data with spin crossover transition for complexes **1** (left) and **2** (right).




Figure S5. Mössbauer spectra of **1** at 300 K (top) and 20 K (bottom). Parameters are provided in Table S3.




Figure S6 Mössbauer spectra of **2** at 300 K (top) and 20 K (bottom). Parameters are provided in Table S3.




Figure S7 Mössbauer spectra of **3** (top) and **4** (bottom) at 20 K. Parameters are provided in Table S3.




Figure S8 Mössbauer spectra of **5** (top) and **6** (bottom) at 20 K. Parameters are provided in Table S3.

Table S1. Crystal parameters of **1** and **2**.

	Comp. 1		Comp. 2	
	100 K	270 K	100 K	270 K
Formula	C ₄₅ H ₄₇ FeN ₉ O ₉	C ₄₅ H ₄₇ FeN ₉ O ₉	C ₄₁ H ₃₇ FeN ₉ O ₅	C ₄₁ H ₃₇ FeN ₉ O ₅
M / g mol ⁻¹	913.76	913.76	791.64	791.64
Temp. / K	100(2)	270(2)	100(2)	270(2)
Crystal system	Triclinic	Triclinic	Monoclinic	Monoclinic
Space group	<i>P</i> Error!	<i>P</i> Error!	<i>P</i> 2 ₁ /c	<i>P</i> 2 ₁ /c
<i>a</i> / Å	11.088(3)	11.339(3)	14.218(3)	14.317(5)
<i>b</i> / Å	12.398(3)	12.582(3)	17.474(4)	17.962(6)
<i>c</i> / Å	16.239(4)	16.370(4)	15.627(4)	15.991(5)
α / °	101.544(4)	100.776(4)	-	-
β / °	100.791(4)	101.014(4)	109.818(3)	110.230(5)
γ / °	100.566(4)	101.059(4)	-	-
<i>V</i> / Å ³	2091.3(8)	2188.4(8)	3652.6(14)	3859(2)
<i>Z</i>	2	2	4	4
<i>d</i> / g cm ⁻³	1.451	1.387	1.440	1.363
μ / mm ⁻¹	0.431	0.412	0.473	0.448
F(000)	956	956	1648	1648
Reflections				
collected / unique	10032 / 7103	12931 / 9734	20780 / 8347	18274 / 8379
<i>R</i> _{int}	0.0631	0.0651	0.0507	0.1810
GOF	1.067	1.028	1.039	1.219
<i>R</i> ₁ (<i>I</i> > 2σ(<i>I</i>))	0.0770	0.0886	0.0603	0.1810
<i>R</i> _w 2 (<i>I</i> > 2σ(<i>I</i>))	0.1762	0.1564	0.1455	0.2736
$\Delta\rho_{\max}$ / e Å ⁻³	1.106	0.393	1.933	0.640
$\Delta\rho_{\min}$ / e Å ⁻³	-0.827	-0.480	-0.729	-0.825
CCDC No.	1956636	1956637	1956638	1956639

Table S2. Crystal parameters of **3**, **4**, **5** and **6** at 100 K.

	Comp. 3	Comp. 4	Comp. 5	Comp. 6
Formula	C ₄₄ H ₃₆ FeN ₁₀ O ₆	C ₄₂ H ₃₂ FeN ₁₂ O ₆	C ₆₄ H ₃₂ Fe ₂ N ₁₂ O ₁₂	C ₄₈ H ₂₂ BF ₄ FeN ₉ O ₆
M / g mol ⁻¹	856.68	856.64	1272.71	963.40
Temp. / K	100(2)	100(2)	100(2)	100(2)
Crystal system	Orthorhombic	Triclinic,	Triclinic	Monoclinic
Space group	<i>Fdd2</i>	<i>PError!</i>	<i>PError!</i>	<i>C2/c</i>
<i>a</i> / Å	17.374(3)	11.6257(11)	10.55(2)	9.260(7)
<i>b</i> / Å	28.891(6)	13.6920(12)	13.14(3)	32.13(3)
<i>c</i> / Å	16.217(3)	13.7914(12)	28.26(7)	18.517(15)
α / °	-	106.5717(12)	95.12(2)	-
β / °	-	111.1607(11)	91.640(19)	103.237(9)
γ / °	-	94.9087(12)	107.52(5)	-
<i>V</i> / Å ³	8140(3)	1918.1(3)	3715(15)	5362(7)
<i>Z</i>	8	2	2	4
<i>d</i> / g cm ⁻³	1.398	1.483	1.138	1.193
μ / mm ⁻¹	0.433	0.461	0.450	0.346
F(000)	3552	884	1296	1952
Reflections				
collected / unique	10689 / 4009	11328 / 8506	21449 / 16347	14165 / 6036
<i>R</i> _{int}	0.0335	0.0188	0.0410	0.1105
GOF	1.018	1.288	1.413	0.870
<i>R</i> ₁ (<i>I</i> > 2 σ (<i>I</i>))	0.0321	0.0541	0.1197	0.0977
<i>R</i> _w 2 (<i>I</i> > 2 σ (<i>I</i>)]	0.0662	0.1647	0.2882	0.2502
$\Delta\rho_{\max}$ / e Å ⁻³	0.311	1.828	1.910	0.775
$\Delta\rho_{\min}$ / e Å ⁻³	-0.311	-1.194	-1.532	-0.441
CCDC No.	1956640	1956641	1956642	1956643

Table S3. Mössbauer parameters for all complexes.

	δ_{IS} (mm / s)	ΔE_{Q} (mm / s)	spin state	Area fraction (%)
Comp.1 (300 K)	0.94	1.39	Fe(II) HS	83
	0.50	0.28	Fe(II) LS	17
Comp.1 (20 K)	0.50	0.28	Fe(II) LS	-
Comp.2 (300 K)	0.94	1.30	Fe(II) HS	-
Comp.2 (20 K)	0.51	0.33	Fe(II) LS	-
Comp.3 (20 K)	0.26	1.48	Fe(II) LS	-
Comp.4 (20 K)	0.22	1.59	Fe(II) LS	-
Comp.5 (20 K)	1.18	2.94	Fe(II) HS	-
Comp.6 (20 K)	1.20	2.93	Fe(II) HS	-

Explanation of alerts in CIFCHECK:

Datablock: Comp1_270K

Alert Level B

The following B-level alerts arise from the diffraction data being slightly weak.

PLAT026_ALERT_3_B Ratio Observed / Unique Reflections (too) Low .. 34% Check

PLAT260_ALERT_2_B Large Average Ueq of Residue Including O5 0.187 Check

PLAT260_ALERT_2_B Large Average Ueq of Residue Including O9 0.172 Check

The following B-level alerts arise from the movement of solvent molecules due to high-temperature measurement.

PLAT360_ALERT_2_B Short C(sp³)-C(sp³) Bond C35 - C36 . 1.24 Ang.

PLAT360_ALERT_2_B Short C(sp³)-C(sp³) Bond C37 - C38 . 1.25 Ang.

PLAT410_ALERT_2_B Short Intra H...H Contact H35A ..H48 . 1.80 Ang.

x,y,z = 1_555 Check

PLAT410_ALERT_2_B Short Intra H...H Contact H35B ..H47 . 1.81 Ang.

x,y,z = 1_555 Check

PLAT410_ALERT_2_B Short Intra H...H Contact H37A ..H38B . 1.82 Ang.

x,y,z = 1_555 Check

PLAT410_ALERT_2_B Short Intra H...H Contact H37B ..H38A . 1.81 Ang.

x,y,z = 1_555 Check

Datablock: Comp2_270K

Alert Level B

The following B-level alerts arise from the diffraction data being slightly weak.

RINTA01_ALERT_3_B The value of Rint is greater than 0.18

Rint given 0.181

PLAT020_ALERT_3_B The Value of Rint is Greater Than 0.12 0.181 Report

PLAT026_ALERT_3_B Ratio Observed / Unique Reflections (too) Low .. 36% Check

PLAT082_ALERT_2_B High R1 Value 0.18 Report

PLAT341_ALERT_3_B Low Bond Precision on C-C Bonds 0.01608 Ang.