

Supporting Information for:

ChemComm
www.rsc.org/chemcomm

Cell-Permeable GPNA Containing Appropriate Backbone Stereochemistry and Spacing Binds Sequence-Specifically to RNA

Anca Dragulescu-Andrasi, Peng Zhou, Gaofei He, Danith H. Ly*

*Department of Chemistry, Carnegie Mellon University, 4400 Fifth Ave.,
Pittsburgh, Pennsylvania*




Figure S1. UV melting curves for PNA1-RNA and GPNA(6-8)-RNA duplexes. Thermal denaturation studies were performed on a Varian Cary 3 spectrophotometer equipped with thermoelectrically-controlled multicell holder, using 2 μ M complementary strands in 100mM NaCl, 10mM sodium phosphate, and 1mM EDTA, pH 7. All samples were annealed before using for melting experiments by heating up to 90 °C and slowly cooling down to room temperature. Thermal denaturation was monitored at 260 nm at a heating rate of 1 °C/min from 20 to 90 °C. The melting transitions were determined from the first derivatives of the UV-melting curves.




Figure S2. CD spectra of PNA1-RNA and GPNA(6-7)-RNA duplexes. CD experiments were performed on a Jasco J-715 spectropolarimeter using the same samples used for the melting experiments. Scans were run from 320 to 200 nm taking measurements every 1 nm. All spectra represent an average of at least 15 scans and were recorded at a rate of 100 nm/min. A 1-cm path-length cuvette was used, and the temperature was maintained at 22 °C.




Figure S3. UV melting curves of GPNA8-RNA and PNA1-RNA duplexes for perfectly matched and mismatched sequences. Thermal denaturation studies were performed on a Varian Cary 3 spectrophotometer equipped with thermoelectrically-controlled multicell holder, using 2 μ M complementary strands in 100 mM NaCl, 10 mM sodium phosphate, and 1 mM EDTA, pH 7. All samples were annealed before using for melting experiments by heating up to 90 °C and slowly cooling down to room temperature. Thermal denaturation was monitored at 260 nm at a heating rate of 1 °C/min from 20 to 90 °C. The melting transitions were determined from the first derivatives of the UV-melting curves.




Figure S4. CD melting curves for GPNA8-RNA duplexes containing perfectly matched and mismatched sequences. The CD spectra were recorded at 260 nm as a function of temperature from 20 to 90°C. CD melting experiments were performed on a Jasco J-715 spectropolarimeter using 2 μ M complementary strands in 100 mM NaCl, 10mM sodium phosphate, and 1mM EDTA, pH 7. All samples were annealed before using for melting experiments by heating up to 90 °C and slowly cooling down to room temperature. The melting transitions were determined from the first derivatives of the UV-melting curves.




Figure S5. UV melting curves for antiparallel and parallel GPNA8-RNA and PNA1-RNA duplexes. Thermal denaturation studies were performed on a Varian Cary 3 spectrophotometer equipped with thermoelectrically-controlled multicell holder, using 2 μM complementary strands in 100 mM NaCl, 10 mM sodium phosphate, and 1 mM EDTA, pH 7. All samples were annealed before using for melting experiments by heating up to 90 $^\circ\text{C}$ and slowly cooling down to room temperature. Thermal denaturation was monitored at 260 nm at a heating rate of 1 $^\circ\text{C}/\text{min}$ from 20 to 90 $^\circ\text{C}$. The melting transitions were determined from the first derivatives of the UV-melting curves.

Supplementary Material (ESI) for Chemical Communications
 # This journal is © The Royal Society of Chemistry 2005

	PNA1 H-GCATGTTTGA- ^L Lys-NH ₂	GPNA8 H- H-G ^D CA ^D TG ^D TT ^D TG ^D A-NH ₂
RNA 5'-UCA <u>A</u> ACAUGC-3'	$T_m = 53\text{ }^\circ\text{C}$	$T_m = 52\text{ }^\circ\text{C}$
RNA-U 5'-UCA <u>U</u> ACAUGC-3'	$T_m = 38\text{ }^\circ\text{C}$	$T_m = 31\text{ }^\circ\text{C}$
RNA-C 5'-UCA <u>C</u> ACAUGC-3'	$T_m = 35\text{ }^\circ\text{C}$	$T_m = 38\text{ }^\circ\text{C}$
RNA-G 5'-UCA <u>G</u> ACAUGC-3'	$T_m = 42\text{ }^\circ\text{C}$	$T_m = 39\text{ }^\circ\text{C}$
RNAp 3'-UCA <u>G</u> ACAUGC-5'	$T_m = 37\text{ }^\circ\text{C}$	$T_m = 31\text{ }^\circ\text{C}$

S6. Tabulation of melting transitions between perfectly matched and mismatched antiparallel GPN8-RNA and PNA1-RNA, and perfectly matched parallel GPNA8-RNAp and PNA1-RNAp duplexes. The indicated T_m s were determined from the data shown in S3, S4, and S6.