

Electronic Supplementary Information

Electronic Energy Harvesting Multi BODIPY – Zinc Porphyrin Dyads Accommodating Fullerene as Photosynthetic Composite of Antenna-Reaction Center

Eranda Maligaspe,^a Tatu Kumpulainen,^b Navaneetha K. Subbaiyan,^a Melvin E. Zandler,^a
Helge Lemmetyinen,^b Nikolai V. Tkachenko,^{b,*} and Francis D'Souza,^{a,*}

^a*Department of Chemistry, Wichita State University, 1845 Fairmount, Wichita, KS 67260-0051, USA; E-mail: Francis.DSouza@wichita.edu; FAX: +316-978-3431*

^b*Department of Chemistry and Bioengineering, Tampere University of Technology, P. O. Box 541, 33101 Tampere, Finland; E-mail: nikolai.tkachenko@tut.fi*

Received:




Fig. S1. Transient absorption decay component spectra of BDP-ZnP:ImC₆₀ complex in DCB. The dashed line shows the differential absorption spectrum right after excitation (at 0 ps delay time).




Fig. S2. Transient absorption decay component spectra of (BDP)-ZnP:ImC₆₀ complex in *o*-dichlorobenzene calculated for a bi-exponential model. The dashed line shows the differential absorption spectrum right after excitation (at 0 ps delay time).