

ELECTRONIC SUPPLEMENTARY INFORMATION

Photoinduced energy transfer processes in hybrid organic-inorganic multichromophoric arrays arranged on a truxene-based platform

Stéphane Diring,^a Barbara Ventura,^b Andrea Barbieri*^b and Raymond Ziessel*^a

^a *Laboratoire de Chimie Organique et Spectroscopies Avancées, (LCOSA) associé au CNRS (UMR 7515), Ecole de Chimie, Polymères, Matériaux (ECPM), Université de Strasbourg (UdS), 25 rue Becquerel, 67087 Strasbourg Cedex 02, France. E-mail: ziessel@unistra.fr*

^b *Istituto per la Sintesi Organica e la Fotoreattività (ISOF), Consiglio Nazionale delle Ricerche (CNR), Via Gobetti 101, 40129 Bologna BO, Italy. E-mail: andrea.barbieri@isof.cnr.it*

Contents

Figure S1. Spectral sum of the model components **Os₂L** and **R**, **B** or **G**.

Figure S2. Emission spectra of iso-absorbing solutions of **Os₂L**, **Os₂LR**, **Os₂LB** and **Os₂LG** in CH₃CN at rt; $\lambda_{ex} = 465$ nm. Inset: zoom on the spectral region between 650 and 1,050 nm.




Figure S1. Spectral sum of the model components Os_2L and R , B or G .




Figure S2. Emission spectra of iso-absorbing solutions of Os_2L , Os_2LR , Os_2LB and Os_2LG in CH_3CN at rt; $\lambda_{\text{ex}} = 465$ nm. Inset: zoom on the spectral region between 650 and 1,050 nm.