

Supplemental Information

Streamlining Gene Expression Analysis: Integration of Co-Culture and mRNA Purification

Scott M. Berry^{1,2}, Chandresh Singh^{1,2}, Jessica D. Lang^{2,3}, Lindsay N. Strotman^{1,2}, Elaine T. Alarid^{2,3}, and David J. Beebe^{1,2}

Departments of Biomedical Engineering¹, The Carbone Cancer Center², and Oncology³, University of Wisconsin-Madison

Supplemental Information Table of Contents:

Table S-1 – PCR primers and annealing temperatures

Table S-1 – PCR primers and annealing temperatures. PCR reactions were run for 40 cycles of 95° C for 15 seconds and the annealing temperature given here for 60 seconds.

Gene	Forward Primer	Reverse Primer	Annealing Temp.
RPLP0	GACAATGGCAGCATCTACAAC	GCAGACAGACACTGGCAAC	60° C
eGFP	CATCTTCTTCAAGGACGAC	TTGTGGCTGTTGTAGTTG	60° C
Vimentin	ACACCCTGCAATCTTTCAGACA	GATTCCACTTTGCGTTCAAGGT	60° C
ER α	CCTGATGATTGGTCTCGTCTG	GGCACACAAACTCCTCTCC	57° C