

Aliphatic Polyester Elastomers Derived from Erythritol and 5-, 7-, and 9-Carbon Diacids

Devin G. Barrett, Wei Luo, and Muhammad N. Yousaf\*

Chemistry Department and the Carolina Center for Genome Science, University of North Carolina at

Chapel Hill, Chapel Hill, 27599

mnyousaf@email.unc.edu

Supplementary Material (ESI)




Figure S1. Fully labeled <sup>1</sup>H spectrum of OErG in d<sub>6</sub>-DMSO.




Figure S2. Fully labeled  $^1\text{H}$  spectrum of OErAd in  $d_6$ -DMSO.




Figure S3. Fully labeled  $^1\text{H}$  spectrum of OErPi in  $\text{d}_6$ -DMSO.




Figure S4. Fully labeled  $^1\text{H}$  spectrum of OErSu in  $\text{d}_6$ -DMSO.




Figure S5. Fully labeled  $^1\text{H}$  spectrum of OErAz in  $\text{d}_6\text{-DMSO}$ .




Figure S6. Fully labeled  $^1\text{H}$  spectrum of OErSe in  $\text{d}_6$ -DMSO.




Figure S7. Fully labeled  $^1\text{H}$  spectrum of OErDo in  $\text{d}_6$ -DMSO.




Figure S8. Fully labeled  $^1\text{H}$  spectrum of OErTe in  $\text{d}_6$ -DMSO.