

SUPPORTING INFORMATION FOR

Biobased polyurethanes obtained from ester/amide pseudo-diols of fatty compounds synthesized by thiol-ene coupling

M. Desroches^a, S. Caillol^a, R. Auvergne^a, B. Boutevin^a, G. David^a*

^aInstitut Charles Gerhardt UMR CNRS 5253 Equipe Ingénierie et Architecture
Macromoléculaire, Ecole Nationale Supérieure de Chimie de Montpellier, 8 rue de
l'Ecole Normale, 34296 Montpellier Cedex 05, France.

Corresponding author e-mail address: ghislain.david@enscm.fr




Figure S1. ¹H NMR of MEC₂




Figure S2. ¹³C NMR of MEC₂




Figure S3. ^{13}C NMR of MEC_4




Figure S4. ¹H NMR of MEC₆




Figure S5. ^{13}C NMR of MEC₆




Figure S6. ¹H NMR of DEC₂




Figure S7. ¹³C NMR of DEC₂




Figure S8. ¹H NMR of DEC₄




Figure S9. ¹³C NMR of DEC₄




Figure S10. ¹H NMR of DEC₆




Figure S11. ¹³C NMR of DEC₆




Figure S12. ¹H of MAC₂




Figure S13. ^1H NMR of MAC_5




Figure S14. ^{13}C NMR of MAC_5




Figure S15. ^1H NMR of DAC_8




Figure S16. ^{13}C NMR of DAC_8




Figure S17. Thermograms recorded under nitrogen of polyurethanes obtained from monoester containing pseudo-telechelic diols (left) and from diester containing pseudo-telechelic diols (right)