

Abietane diterpenoid of *Salvia Sahendica* Boiss and Buhse potently inhibits MCF-7 breast carcinoma cells by suppression of PI3K/AKT pathway

Vala Kafil^{a, b, c†}, Morteza Eskandani^{a, b†}, Yadollah Omid^{a, c}, Hossein Nazemiyeh^{a, c *}, Jaleh Barar^{a, c **}

^a Research Center for Pharmaceutical Nanotechnology, Tabriz University of Medical Sciences, Tabriz, Iran

^b Student Research Committee, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran


^c Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran

[†] *These authors contributed equally to this work and should be considered as first authors.*


Corresponding authors:

* Hossein Nazemiyeh, Professor of Pharmacognosy, Nazemiyehh@tbzmed.ac.ir, Tel: +98 41 333367014, Fax: +98 41 33367929

** Jaleh Barar, Associate Professor of Biomedicine, jbarar@tbzmed.ac.ir, Tel: +98 41 33367914, Fax: +98 41 33367929




Supplementary 1.1. ¹H-NMR of Ketoethiopinone (1)




Supplementary 1.2. ^{13}C -NMR of Ketoethiopinone (1)

Dr.Nazemiyeh

CDC13 91/10/2

Supplementary 2.1. ¹H-NMR of Ortho-diacetate aethiopinone (2)



Supplementary 2.2. ^{13}C -NMR of Ortho-diacetate aethiopinine (2)