

Zone analysis by two-dimensional electrophoresis with accelerator mass spectrometry of *in vivo* protein bindings of idiosyncratic hepatotoxicants troglitazone and flutamide bioactivated in chimeric mice with humanized liver

Hiroshi Yamazaki,^{*a} Shunji Kuribayashi,^b Tae Inoue,^c Tomohiro Honda,^d Chise Tateno,^c Ken Oofusa,^e Shinichi Ninomiya,^f Toshihiko Ikeda,^g Takashi Izumi^d and Toru Horie^h

^a Showa Pharmaceutical University, Machida, Japan. E-mail: hyamazak@ac.shoyaku.ac.jp

^b Otsuka Pharmaceutical Factory, Inc., Naruto, Japan

^c PhoenixBio, Co., Higashi-Hiroshima, Hiroshima 739-0046, Japan

^d Daiichi Sanyo, Co., Shinagawa-ku, Japan

^e Idea Consultants Inc., Suminoe-ku, Osaka 559-8519, Japan

^f Sekisui Medical Co., Chuo-ku, Japan

^g Yokohama College of Pharmacy, Totsuka-ku, Yokohama 245-0066, Japan

^h Drug Discovery and Development Institute, Tsukuba, Japan

Toxicol. Res., 2015, **4**, 106–111, DOI: [10.1039/C4TX00068D](https://doi.org/10.1039/C4TX00068D)

We would like to confirm that there is no Electronic supplementary information (ESI) associated with this article and apologise for any inconvenience experienced.