


Colorimetric Cyanide Detection Using an Azobenzene Acid in Aqueous Solutions


Xiaoding Lou, Jingui Qin and Zhen Li *

Department of Chemistry, Hubei Key Laboratory on Organic and Polymeric Optoelectronic Materials, Wuhan University, Wuhan, 430072, China. Fax: +86-27-6875-6757; Phone: +86-27-6225-4108; E-mail: lizhen@whu.edu.cn.

Equation S1: the reaction between Cu^{2+} and CN^-



Equation S2: the reaction between CuCN and CN^-



The stability constants (K), $[\text{Cu}(\text{CN})_2]^-$: $K = 1.00 \times 10^{24}$; $[\text{Cu}(\text{CN})_4]^{3-}$: $K = 2.00 \times 10^{30}$.




Figure S1. UV-vis spectra of compound **I** with increasing amounts of Cu^{2+} in water at pH 6.7. The concentration of **I** was 3.0×10^{-5} mol/L.




Figure S2. UV-vis spectra of compound **I** with increasing amounts of Cu^{2+} in water at pH 6.7. The concentration of **I** was 1.5×10^{-5} mol/L.




Figure S3. UV-vis spectra of the solution of compound **I** (3.0×10^{-5} mol/L) and Cu^{2+} (3.0×10^{-5} mol/L) with increasing amounts of CN^- in water at pH 6.7.




Figure S4. UV-vis spectra of the solution of compound **I** (1.5×10^{-5} mol/L) and Cu^{2+} (7.0×10^{-6} mol/L) with increasing amounts of CN^- in water at pH 6.7.




Figure S5. UV-vis spectra of complex **I** (A) at the presence of different cations mixture (1.0×10^{-5} mol/L) in water at pH 7.4 (20 mM HEPES, 135 mM NaCl). The concentration of compound **I** was 6.0×10^{-5} mol/L, and that of Cu^{2+} was 1.0×10^{-5} mol/L.




Figure S6. UV-vis spectra of complex **I** (A) at the presence of different cations mixture (1.0×10^{-5} mol/L) in water at pH 7.4 (20 mM HEPES, 135 mM NaCl). The concentration of compound **I** was 6.0×10^{-5} mol/L (Cu^{2+} : 1.0×10^{-5} mol/L; CN^- : 6.0×10^{-5} mol/L).



Figure S7. Different solutions of compound **I** and copper ions (1.0×10^{-4} and 5.0×10^{-5} mol/L, respectively) in water at pH 7.4 (20 mM HEPES, 135 mM NaCl) in the presence of different concentrations of CN^- . From left to right ($\times 10^{-5}$ mol/L): no copper ions and no cyanide, 0, 1.0, 2.0, 3.0, 4.0, 5.0, 6.0, 7.0, 8.0 and 9.0.



Figure S8. From left to right: compound **I** (1.0×10^{-4} mol/L), compound **I** (1.0×10^{-4} mol/L)+ Cu^{2+} (5.0×10^{-5} mol/L), compound **I** (1.0×10^{-4} mol/L)+ Cu^{2+} (5.0×10^{-5} mol/L)+ CN^- (6.0×10^{-5} mol/L) at pH 7.4 (20 mM HEPES, 135 mM NaCl).



Figure S9. Different solutions of compound **I** and copper ions in water at pH 7.4 [20 mM HEPES, 135 mM NaCl] (1.0×10^{-4} and 5.0×10^{-5} mol/L, respectively) in the presence of anion (6.0×10^{-5} mol/L). From left to right: no copper ions and no anion; no anion, CN^- , SCN^- , F^- , ClO_4^- , HSO_4^- , Br^- , H_2PO_4^- , NO_2^- , SO_4^{2-} , I^- , Cl^- , IO_3^- , the mixture.