Analyst Electronic Supplementary Material

Hydride abstraction in positive-ion electrospray interface: oxidation of 1,4-dihydropyridines in electrospray ionization mass spectrometry

Yunfeng Chai, Hezhi Sun, Jieping Wan, Yuanjiang Pan* and Cuirong Sun*

panyuanjiang@zju.edu.cn

Contents

• Figure S1. Full-scan APCI-IT mass spectrum of compound 1.

• **Figure S2**. Full-scan ESI-IT mass spectra of compound **1** with the solvent as (a) CH₃CN, (b) CH₃OH, (c) CH₃CN-H₂O (1: 1, v/v), (d) CH₃OH-H₂O (1: 1, v/v).

• Figure S3. HPLC chromatogram (a) and EIC chromatograms of ions at m/z 338 (b) and m/z 336 (c) of compound 1 determined by HPLC-MS method.

• Figure S4. Full-scan ESI-FTICR mass spectrum of compound 1.

• **Figure S5**. Cyclic voltammogram (scan rate 0.1 V s⁻¹) of compound **1** (1 mmol L⁻¹) obtained in acetonitrile using 0.1 M (TBA)PF₆ as the supporting electrolyte.

• Figure S6. CID mass spectra of the $[M + H]^+$ ions of (a) compound 1, (b) compound 2, (c) compound 3.

• Figure S7. Full-scan ESI-IT mass spectra of (a) compound 2, (b) compound 3.



Figure S1. Full-scan APCI-IT mass spectrum of compound 1.



Figure S2. Full-scan ESI-IT mass spectra of compound 1 with the solvent as (a) CH₃CN, (b) CH₃OH, (c) CH₃CN-H₂O (1: 1, v/v), (d) CH₃OH-H₂O (1: 1, v/v).



Figure S3. HPLC chromatogram (a) and EIC chromatograms of ions at m/z 338 (b) and m/z 336 (c) of compound 1 determined by HPLC-MS method.



Figure S4. Full-scan ESI-FTICR mass spectrum of compound 1.



Figure S5. Cyclic voltammogram (scan rate 0.1 V s⁻¹) of compound **1** (1 mmol L⁻¹) obtained in acetonitrile using 0.1 M (TBA)PF₆ as the supporting electrolyte.



Figure S6. CID mass spectra of the $[M + H]^+$ ions of (a) compound 1, (b) compound 2, (c) compound 3.



Figure S7. Full-scan ESI-IT mass spectra of (a) compound 2, (b) compound 3.