

Electronic Supplementary Information

Simple, sensitive and selective detection of dopamine using dithiobis(succinimidylpropionate)-modified gold nanoparticles as colorimetric probes

Lin Liu^{*a}, Sujuan Li^a, Leilei Liu^a, Dehua Deng^a, Ning Xia^{*b}

^aCollege of Chemistry and Chemical Engineering, Anyang Normal University, Anyang, Henan 455000, People's Republic of China

^bCollege of Chemistry and Chemical Engineering, Central South University, Changsha, Hunan 410083, People's Republic of China

^{*}Corresponding author. Tel: 86-732-2900040; *E-mail address:* liulin82414@yahoo.com.cn (L. Liu), xianing82414@csu.edu.cn (N. Xia)




Fig. S1 UV/Vis spectra of the DA- Fe^{3+} complex. The concentration ratio of DA and Fe^{3+} kept at 1:1.





Fig. S2 Dependence of absorbance change of DA-anchored DSP-AuNPs on the Fe^{3+} concentration at different pH. The concentration of DA kept at 1 μM .

A pH 6.4



B pH 7.4




Fig. S3 Color change of DSP-AuNPs in different systems: DA + metal ions and DA/interferences + Fe³⁺.




Fig. S4 Effect of BSA on the A₆₅₀/A₅₂₀ ratio at different concentrations of DA. The concentration of BSA kept at 0.5 μM.