

Supplementary Information:

The evaluation of Coulombic interaction in the oriented-attachment growth of colloidal nanorods

Weidong He^{1,2*}, Junhao Lin³, Xiao Lin^{4*}, Ning Lu⁵, Ming Zhou⁶, and Kelvin HL Zhang⁷

¹ Vanderbilt Interdisciplinary Program in Materials Science, Vanderbilt University, Nashville, Tennessee 37234-0106, USA

² Vanderbilt Institute of Nanoscale Science and Engineering, Vanderbilt University, Nashville, Tennessee 37234-0106, USA

³ Department of Physics and Astronomy, Vanderbilt University, Nashville, Tennessee 37234-0106, USA

⁴ Institute of Physics, Chinese Academy of Sciences, P.O. Box 603, Beijing 100190, People's Republic of China

⁵ Department of Breast Cancer Medical Oncology, Breast Cancer Prevention Key Laboratory of Ministry of Education, Tianjin Medical University Cancer Institute and Hospital, Tianjin 300060, People's Republic of China

⁶ Department of Chemistry, The University of Hong Kong, Pokfulam Road, Hong Kong

⁷ Department of Chemistry, Chemistry Research Laboratory, University of Oxford, Mansfield Road, Oxford OX1 3TA, United Kingdom

* Email: weidong.he@vanderbilt.edu; dr.linxiao@gmail.com




Figure S-1. The plots of $\text{Log}(CI/kT)$ versus charge density for nanorods with radii of 1, 5, 25, and 50 nm.