Reliable SERS Substrate Fabricated by Assembling DNA-embedded Au-Ag Coreshell Nanoparticles on Silicon Slide

Zhong Zhang, Sha Zhang, Mengshi Lin*

Division of Food Systems & Bioengineering, University of Missouri, Columbia, MO, USA 65211-5160

Correspondence: M. Lin, Division of Food System & Bioengineering, University of Missouri, Columbia, MO, USA. 65211; Tel: (573) 884-6718; fax (573) 884-7964. (E-mail: linme@missouri.edu)

Table of Content

Figure S1 TEM images of core-satellite structures using 40 nm Au-DNA NPs as core and DNA embedded Au-Ag core-shell NPs with different Ag coating as satellites: (a) 0 nm Ag shell on 17 nm Au NPs; (b) 1.0 nm Ag shell; (c) 2.1 nm Ag shell; (d) 3.1 nm Ag shell; and (f) 27 nm Au NPs as satellite NPs.

Figure S2 TEM image of core-satellite structures: (a) before the plasma treatment; (b) after the plasma treatment.

Figure S3 (a) Raman spectra of 4-MBA (10^{-6} M) acquired from three different assembled substrates using a 785 nm laser; (b) Stability test of the assembled substrate after 4-month storage using a 785 nm laser.

Figure S4 Raman spectra of 4-MBA (10⁻⁶ M) acquired from the assembled substrates and the Klarite using 5-s acquisition time at the excitation of 633 nm without baseline correction.

Figure S5 Raman spectra of 4-MBA (10^{-6} M) for the Klarite using 10-s acquisition time at the excitation of 633 nm.

Figure S1 TEM images of core-satellite structures using 40 nm Au-DNA NPs as core and DNA embedded Au-Ag core-shell NPs with different Ag coating as satellites: (a) 0 nm Ag shell on 17 nm Au NPs; (b) 1.0 nm Ag shell; (c) 2.1 nm Ag shell; (d) 3.1 nm Ag shell; and (f) 27 nm Au NPs as satellite NPs.



Figure S2 TEM image of core-satellite structures: (a) before the plasma treatment; (b) after the plasma treatment.



Figure S3 (a) Raman spectra of 4-MBA (10^{-6} M) acquired from three different assembled substrates using a 785 nm laser; (b) Stability test of the assembled substrate after 4-month storage using a 785 nm laser.



Figure S4 Raman spectra of 4-MBA (10^{-6} M) acquired from the assembled substrates and the Klarite using 5-s acquisition time at the excitation of 633 nm without baseline correction.



Figure S5 Raman spectra of 4-MBA (10^{-6} M) for the Klarite using 10-s acquisition time at the excitation of 633 nm.

