

An Attenuated Total Reflection (ATR) and Raman spectroscopic investigation into the effects of chloroquine on *Plasmodium falciparum*-infected red blood cells

M. Kozicki¹, D. J. Creek², A. Sexton², B. J. Morahan⁴, A. Wesełucha-Birczyńska¹ and B. R. Wood^{3*}

¹ Faculty of Chemistry, Ingardena 3, Kraków, Poland

² Monash Institute of Pharmaceutical Sciences, Monash University, Parkville, VIC 3052, Australia

³ Centre for Biospectroscopy and School of Chemistry, Monash University, Wellington Rd., VIC. 3800, Australia

⁴ Department of Microbiology, Monash University, Clayton, VIC 3800, Australia

Electronic Supplementary Information




Fig S1. Raman spectra of untreated (Plf untreated) and CQ-treated (Plf+CQ) hemoglobin part of *P. falciparum*-iRBCs using 532 nm excitation for 2nd independent trial. The insets show curves fitted spectral ranges (left 3300-2600 cm^{-1} ; right 1700-1300 cm^{-1}).

Table S1. Band ratio's for a curve-fitting approach and calculated integrated areas under the oxy/deoxy maker bands both in the fundamental and overtone region for 2nd independent trial.

	ν_4 oxy / ν_4 deoxy ~ 1377/1361	ν_{37} oxy / ν_{37} deoxy ~ 1587/1575	$2\nu_4$ oxy / $2\nu_4$ deoxy ~ 2754/2722	$2\nu_{37}$ oxy / $2\nu_{37}$ deoxy ~ 3175/3150
CQtreated	2,12±0,12	3,11±0,15	2,19±0,26	2,24±0,45
untreated	0,72±0,08	0,83±0,17	0,68±0,04	0,75±0,07