

Electronic Supplementary Information

Determination **of** aristolochic acids in rat serum by high performance liquid chromatography-Q TOF tandem mass spectrometry

Jiangping Gu^a, Pinjie Zhang^a, Feifang Zhang^{a*}, HeQing Shen^b, Bingcheng Yang^{a*},

Xinmiao Liang^a, Xiaogang Chu^c

^aSchool of Pharmacy, East China University of Science and Technology, Shanghai, 200237, China

^bInstitute of Urban Environment, Chinese Academy of Sciences, Xiamen, 361021, China

^cInstitute of Food Safety, Chinese Academy of Inspection and Quarantine, Beijing 100123, China

Correspondence: Dr. Feifang Zhang, School of Pharmacy, East China University of Science and Technology, Shanghai 200237, China

Email: zhangff@ecust.edu.cn

Tel/Fax: 0086-21-64250627

SI- Table 1. MS operation conditions for four AAs and IS (Naproxen)

Parent ion (m/z)	Daughter ion (m/z)	Scan time (msec)	Analyte	Declustering potential (DP)	Entrance potential (EP)	Collision energy (CEP)	Collision energy (CE)	Collision Cell Exit Potential (CXP)
340	46	50	AAI	-25	-2	-18	-16	-4
310	46	50	AAII	-25	-1	-26	-14	-4
326	236	50	AAIIIa	-40	-4	-18	-26	-2
356	266	50	AAIVa	-40	-305	-20	-30	-2
229	73	50	IS	-20	-6	-22	-42	-2