

Sensitively electrochemical detection of copper ions based on the copper (II) ion assisted etching of Au@Ag nanoparticles

Tianxiang Wu, Teng Xu, and Zhanfang Ma*

Department of Chemistry, Capital Normal University, Beijing 100048, China

Email: mazhanfang@cnu.edu.cn




Fig. S1. The UV-Vis spectra of AuNPs and Au@AgNPs




Fig. S2. The XPS spectra of prepared Au@AgNPs nanoparticles

Table S1. Comparison of this work with other published methods.

Ref.	Analytes	Detection limit of Cu ²⁺ (M)	Linear range of Cu ²⁺ (M)
1	Cu ²⁺	2.0×10 ⁻⁶	1.0×10 ⁻⁶ - 5.0×10 ⁻²
2	Cu ²⁺	7.8×10 ⁻⁶	1.0×10 ⁻⁵ - 5.0×10 ⁻¹
3	Cu ²⁺	3.5×10 ⁻⁹	7.9×10 ⁻⁹ - 3.2×10 ⁻²
4	Cu ²⁺	1.6×10 ⁻⁶	2.0×10 ⁻⁶ - 1.0×10 ⁻¹
5	Cu ²⁺	8.1×10 ⁻⁸	2.0×10 ⁻⁷ - 1.0×10 ⁻¹
6	Cu ²⁺ , Pb ²⁺	2.1×10 ⁻⁹	5.0 ×10 ⁻⁹ - 1.0×10 ⁻⁴
This work	Cu ²⁺	3.0×10 ⁻¹⁰	1 ×10 ⁻⁹ - 1.0×10 ⁻⁴

References

- [1] S. Yoshimoto, H. Mukai, T. Kitano, Y. Sohrin, Copper(II)-selective membrane electrode based on hydrotris(3-isopropylpyrazolyl)methane in a poly(vinyl chloride) matrix, *Anal. Chim. Acta* 494 (2003) 207-213.
- [2] V.K. Gupta, R.N. Goyal, N. Bachheti, L.P. Singh, S. Agarwal, A copper-selective electrode based on bis(acetylacetone)propylenediamine, *Talanta* 68 (2005) 193-197.
- [3] M.H. Mashhadizadeh, K. Eskandari, A. Foroumadi, A. Shafiee, Copper(II) modified carbon paste electrodes based on self-assembled mercapto compounds-gold-nanoparticle, *Talanta* 76 (2008) 497-502.
- [4] A.K. Jain, V.K. Gupta, L.P. Singh, J.R. Raisoni, Chelating ionophore based membrane sensors for copper(II) ions, *Talanta* 66 (2005) 1355-1361.
- [5] S. Chandra, C.K. Singh, H. Agarwal, R.K. Agarwal, A copper(II)-selective PVC membrane electrode based on a macrocyclic ligand, 1,2,5,6,8,11-hexaazacyclododeca-7,12-dione-2,4,8,10-tetraene, *Anal. Sci.* 23 (2007) 683-687.
- [6] M. Bagherzadeh, M. Pirmoradian, F. Riahi, Electrochemical detection of Pb and Cu by using DTPA functionalized magnetic nanoparticles, *Electrochim. Acta* 115 (2014) 573-580.