Electronic Supplementary Information (ESI)

for

Poly(thymine)-templated fluorescent copper nanoparticles for label-free detection of N-acetylcysteine in pharmaceutical samples

Hai-Bo Wang,*^a Hong-Ding Zhang,^a Ying Chen,^a Li-Juan Ou,*^b and

Yan-Ming Liu^a

^aCollege of Chemistry and Chemical Engineering, Xinyang Normal University, Xinyang 464000, PR China

^bCollege of Material and Chemical Engineering, Hunan Institute of Technology, Hengyang 421002, PR China

Tel.: +86 376 6391172; Fax: +86 376 6391172.

E-mail address: wanghaibohn@163.com (H.-B. Wang); oulijuanann@aliyun.com (L.-J. Ou).



Fig. S1 The chemical structure of N-acetylcysteine (NAC) and captopril.



Fig. S2 Fluorescence excitation (a) and emission (b) spectra of the poly T templated copper nanoparticles.



Fig. S3 (A) Typical TEM image of poly T-templated Cu NPs. (B) TEM image of poly T-templated Cu NPs in the presence of of 100 μ M NAC.



Fig. S4 (A) The effect of Cu^{2+} concentration on the fluorescence intensity of poly T templated Cu NPs. (B) The effect of ascorbic acid concentration on the fluorescence intensity of the poly T templated Cu NPs. (C) The effect of incubation time of ascorbic acid on the fluorescence intensity of the poly T templated Cu NPs. (D) The effect of incubation time of NAC on the fluorescence intensity of the poly T templated Cu NPs.