

Ultraviolet-Visible Spectroscopic Characterisation of Automotive Window Tints for Forensic Purposes

Karin J. van der Pal,^{a, b} Mark Maric,^{a, b} Wilhelm van Bronswijk^b and Simon W. Lewis^{a, b*}

^aNanochemistry Research Institute, GPO Box U1987, Perth, Western Australia 6845

^bDepartment of Chemistry, Curtin University, GPO Box U1987, Perth, Western Australia 6845, Australia.

E-mail: S.Lewis@curtin.edu.au; Tel: +61 8 9266 2484.

Supporting Information

Table S1 :List of window tints used in study

Manufacturer	Tint	Transmission
Altrex	Dark Smoke	20%
John Hughes	UltraPlus Black	20%
John Hughes	UltraPlus Black	35%
John Hughes	UltraPlus Security	35%
Tint-A-Car	UV	5%
Tint-A-Car	VLT	17%
Tint-A-Car	Octane	20%
Tint-A-Car	Octane	35%
Tint-A-Car	EMI	35%
Tint-A-Car	Security	41%
DIY Home Tint	Black	5%
DIY Home Tint	Dark Black	8%
DIY Home Tint	Medium Black	15%
DIY Home Tint	Black	35%
DIY Home Tint	Light Black	35%
Protint	Black	5%
Protint	Black	10%
Protint	Black	15%
Protint	Black	30%
Protint	Black	35%
Protint	Black	48%
Hanita Pacific	Night Owl	5%
Hanita Pacific	Natura	15%
Hanita Pacific	Natura	28%

Fig. S1. Variance plot for Window Tint PCA

