Structure transformation and remarkable site-distribution modulation of Eu³⁺ ions in CaMoO₄:Eu³⁺ nanocrystals

under high pressure

Zhi Zhao,^a Zhilei Sui,^b Xiantao Wei,^c Jian Zuo,^a Xianwen Zhang,^{*d} Rucheng Dai,^c Zengming Zhang^{*c} and Zejun Ding^{a,b}

^aHefei National Laboratory for Physical Sciences at the Microscale, University of Science and Technology of China, Hefei, 230026, China

^bDepartment of Physics, University of Science and Technology of China, Hefei, 230026, China

^cThe Center of Physics Experiment, School of Physical Sciences, University of Science and Technology of China, Hefei 230026, China

^dInstitute of Advanced Energy Technology and Equipment, Hefei University of Technology, Hefei, 230009, China

*Corresponding authors: xianwen.zhang@hfut.edu.cn, zzm@ustc.edu.cn

Experimental results:




Fig. S1 The SEM image of CaMoO₄:Eu³⁺ calcined at 1000 °C for 6 hours.




Fig. S2 The XRD patterns of CaMoO₄:Eu³⁺ calcined at 1000 °C for 6 hours.




Fig. S3 (a) Raman spectra of CaMoO₄: Eu³⁺ bulk at different pressures. Asterisks refer to emissions from the ${}^{5}D_{0} \rightarrow {}^{7}F_{3}$ transition of Eu³⁺ ions. (b)

Pressure dependence of the shifts of Raman modes of CaMoO₄:Eu³⁺ bulk.




Fig. S4 UV-vis absorbance spectra of $CaMoO_4$: Eu³⁺ nanocrystals (a) and bulk (b) at ambient pressure.




Fig. S5 Photoluminescence spectra of CaMoO₄: Eu^{3+} bulk at different pressures. The inset is the enlarged emission spectra of ${}^{5}D_{0} \rightarrow {}^{7}F_{0}$ transition

of CaMoO₄:Eu³⁺ bulk at ambient pressure.