

Supporting Information

Halogen bonding. The role of polarizability of the electron-pair donor

Darío J. R. Duarte,^a Gladis L. Sosa,^b Nélide M. Peruchena,^a and Ibon Alkorta^c

^aLaboratorio de Estructura Molecular y Propiedades, Área de Química Física, Departamento de Química, Facultad de Ciencias Exactas y Naturales y Agrimensura, Universidad Nacional del Nordeste, Avda. Libertad 5460, (3400) Corrientes, Argentina.

^bLaboratorio de Química Teórica y Experimental, Departamento de Química. Facultad Regional Resistencia, Universidad Tecnológica Nacional, French 414, (3500) Resistencia, Chaco, Argentina.

^cInstituto de Química Médica, CSIC, Juan de la Cierva, 3, E-28006 Madrid, Spain.

* Corresponding author:

Darío J. R. Duarte, e-mail: djr_duarte@hotmail.com

Contents

Pg. S2 Figure S1. Correlations between the binding energy and polarizabilities perpendicular, α_{\perp} (a), parallel, α_{\parallel} (b), and spherical, α_s (c).

Pg. S2 Figure S2. Envelopes of the function $L(\mathbf{r})=0.0016$ au for F–Br \cdots Br–F complex.

Figure S1. Correlations between the binding energy and polarizabilities perpendicular, α_{\perp} (a), parallel, α_{\parallel} (b), and spherical, α_S (c).




Figure S2. Envelopes of the function $L(\mathbf{r})=0.0016$ au for F-Br \cdots Br-F complex.

