## **Supporting Information**

## Molecular determinants for selective $C_{25}$ -hydroxylation of vitamins $D_2$ and $D_3$ by fungal peroxygenases

Fátima Lucas,<sup>1,a\*</sup> Esteban D. Babot,<sup>1,b</sup> Marina Cañellas,<sup>1,a,c</sup> José C. del Río,<sup>b</sup> Lisbeth Kalum,<sup>d</sup> René Ullrich,<sup>e</sup> Martin Hofrichter,<sup>e</sup> Victor Guallar,<sup>a,f</sup> Angel T. Martínez<sup>g</sup> and Ana Gutiérrez<sup>b\*</sup>

The Supporting Information shows the main interactions of cholecalciferol and ergocalciferol at the peroxygenase heme access channel (**Fig. S1** and **S2**, respectively), the effect of side-chain structure on the conversion rates of five sterols by the *A. aegerita* and *C. cinerea* peroxygenases (**Table S1**), and distributions of estimated Fe=O-H and O-H-C angles in the peroxygenase reactions with cholecalciferol and ergocalciferol (**Fig. S3**).

<sup>&</sup>lt;sup>1</sup> These three authors equally contributed to this work

<sup>&</sup>lt;sup>a</sup> Joint BSC-CRG-IRB Research Program in Computational Biology, Barcelona Supercomputing Center, Jordi Girona 29, E-08034 Barcelona, Spain, E-mail: fati.lucas@gmail.com

<sup>&</sup>lt;sup>b</sup> Instituto de Recursos Naturales y Agrobiología de Sevilla, CSIC, Reina Mercedes 10, E-41012 Seville, Spain. Email: anagu@irnase.csic.es; Fax: +32 954624002; Tel: +32 954624711

<sup>&</sup>lt;sup>c</sup> Anaxomics Biotech, Balmes 89, E-08008 Barcelona, Spain

<sup>&</sup>lt;sup>d</sup> Novozymes A/S, Krogshoejvej 36, 2880 Bagsvaerd, Denmark

<sup>&</sup>lt;sup>e</sup> TU Dresden, Department of Bio- and Environmental Sciences, Markt 23, 02763 Zittau, Germany

<sup>&</sup>lt;sup>f</sup> ICREA, Passeig Lluís Companys 23, E-08010 Barcelona, Spain

g Centro de Investigaciones Biológicas, CSIC, Ramiro de Maeztu 9, E-28040 Madrid, Spain





**Fig. S1** Main interactions (below 3 Å) for cholecalciferol with the peroxygenases of *A. aegerita* (**A**) and *C. cinerea* (**B**).



**Fig. S2** Main interactions (below 3 Å) for ergocalciferol with the peroxygenases of *A. aegerita* (**A**) and *C. cinerea* (**B**).

		Conversion (%)	
		A. aegerita peroxygenase	<i>C. cinerea</i> peroxygenase
A	21 $22$ $24$ $25$ $27C_{19}H_{30}O 26 27$	64	100
В	21 $22$ $24$ $25$ $27$ $21$ $20$ $23$ $23$ $25$ $27$ $21$ $23$ $25$ $27$ $21$ $23$ $23$ $26$	47	30
С	21 $22$ $24$ $25$ $27$ $27$ $21$ $23$ $25$ $27$ $21$ $23$ $24$ $25$ $27$ $27$ $21$ $23$ $26$ $27$	10	6
D	21 $22$ $24$ $25$ $27$ $27$ $20$ $23$ $25$ $27$ $23$ $24$ $25$ $27$ $27$ $23$ $23$ $26$	13	6
Ε	21 $22$ $24$ $25$ $27$ $23$ $23$ $24$ $25$ $27$ $23$ $23$ $23$ $26$ $27$	2	2

**Table S1** Conversion degree of steroids with different side-chains (*A-E*: cholesterol, campesterol, ergosterol, sitosterol and stigmasterol, respectively) by *A. aegerita* and *C. cinerea* peroxygenases<sup>a</sup>

<sup>a</sup>From E. D. Babot, J. C. del Río, M. Cañellas, F. Sancho, F. Lucas, V. Guallar, L. Kalum, H. Lund, G. Gröbe, K. Scheibner, R. Ullrich, M. Hofrichter, A. T. Martínez, and A. Gutiérrez. 2015. Steroid hydroxylation by basidiomycete peroxygenases: A combined experimental and computational study. *Appl. Environ. Microbiol.* 81:4130-4142



**Fig. S3** Relative distributions of computed Fe=O-H and O-H-C angles for the reactions of *A. aegerita* (**A**, **C**) and *C. cinerea* (**B**, **D**) peroxygenases at the C<sub>24</sub>, C<sub>25</sub>, C<sub>26</sub> and C<sub>27</sub>, positions of cholecalciferol (blue) and ergocalciferol (green) and the C<sub>28</sub> position of ergocalciferol. Structures were filtered by energy and distance between the pertinent hydrogen atom and the haem compound I oxygen (below 3 Å). QM/MM studies for P450<sub>cam</sub><sup>b</sup> show that in a pre-arranged reactive position camphor adopts angles of 130° for Fe=O-H and 170° for O-H-C. While for that latter no correlation is found to the theoretical predictions, in the case of Fe=O-H<sub>24</sub> angles between 120 and 140° are favoured in C<sub>24</sub> for cholecalciferol reacting with *A. aegerita* (**A**) in

agreement with the experimentally observed formation of 21% product in this position. In the case of the remaining 3 reactions, few structures are found with angles below 145°. In the case of Fe=O-H<sub>25</sub> (as expected) all compounds present favourable angles. Analogous to the case of the distance distribution analysis, no conclusions can be made for the reactivity of C<sub>26</sub>, C<sub>27</sub> and C<sub>28</sub>. <sup>b</sup>J. C. Schöneboom, S. Cohen, H. Lin, S. Shaik and W. Thiel, J. Am. Chem. Soc., 2004, 126, 4017-4034