

Electronic Supporting Information

Aromatic Hydroxylation by an Oxo-bridged Diiron(III) Complex: A Bio-inspired Functional Model of *Toluene Monooxygenases*

Ambica Kejriwal,^a and Pinaki Bandyopadhyay^a and Achintesh N. Biswas^{*b}

^a Department of Chemistry, University of North Bengal, Raja Rammohunpur, Siliguri 734013, India. Fax: 91 353 2699001; Tel: 91 353 277684; E-mail: pbchem@rediffmail.com

^b Department of Chemistry, Siliguri College, Siliguri 734001, India; E-mail: achintesh@gmail.com

Contents

Figure S1a. ESIMS spectrum obtained during the course of benzene hydroxylation by **2**/H₂O₂.

Figure S1b. Experimental and theoretical isotope distribution patterns. The peak at $m/z = 342$ has been assigned to $[(\text{bpmen})\text{Fe}^{\text{III}}(\text{O})]^+$ whereas the peak at $m/z = 361$ can be assigned to the species $[\{(\text{bpmen})\text{Fe}^{\text{III}}(\text{O})(\text{OH}_2)\} + \text{H}]^+$. The peak at $m/z = 478$ can be assigned to $[(\text{bpmen})\text{Fe}^{\text{III}}(\text{OPh})(\text{OH})(\text{MeCN}) + \text{H}]^+$.

Figure S2. Rate of formation of phenolate as a function of benzene concentration

Figure S3. Electronic changes observed after the addition of additional H₂O₂ (10 mM) to the final diiron(III) complex (black line). The final spectrum corresponding to the phenolate bound iron(III) species is shown in green.

Figure S4. Rate of formation of phenolate as a function of benzene concentration.

Figure S4. X-band EPR spectrum obtained after the addition of 10 mM H₂O₂ to an acetonitrile solution of **2** (1.0 mM) at 298 K followed by rapid freezing of the solution (red line). EPR spectrum obtained after the addition of 10 mM H₂O₂ to an acetonitrile solution of **2** (1.0 mM) and benzene (300 mM) at 298 K followed by rapid freezing of the solution (black line).




Figure S1a. ESIMS spectrum obtained during the course of benzene hydroxylation by **2**/H₂O₂.




Figure S1b. Experimental and theoretical isotope distribution patterns. The peak at $m/z = 342$ has been assigned to $[(\text{bpmen})\text{Fe}^{\text{III}}(\text{O})]^+$ whereas the peak at $m/z = 361$ can be assigned to the species $[\{(\text{bpmen})\text{Fe}^{\text{III}}(\text{O})(\text{OH}_2)\}+\text{H}]^+$. The peak at $m/z = 478$ can be assigned to $[(\text{bpmen})\text{Fe}^{\text{III}}(\text{OPh})(\text{OH})(\text{MeCN})+\text{H}]^+$.




Figure S2. Rate of formation of phenolate as a function of benzene concentration.




Figure S3. Electronic changes observed after the addition of additional H₂O₂ (10 mM) to the final diiron(III) complex (black line). The final spectrum corresponding to the phenolate bound iron(III) species is shown in green.




Figure S4. X-band EPR spectrum obtained after the addition of 10 mM H₂O₂ to an acetonitrile solution of **2** (1.0 mM) at 298 K followed by rapid freezing of the solution (red line). EPR spectrum obtained after the addition of 10 mM H₂O₂ to an acetonitrile solution of **2** (1.0 mM) and benzene (300 mM) at 298 K followed by rapid freezing of the solution (black line).