Electronic Supplementary Information

Assembly of 1D, 2D and 3D lanthanum(III) coordination polymers with perchlorinated benzenedicarboxylates: positional isomeric effect, structural transformation and ring-opening polymerisation of glycolide

Sheng-Chun Chen,^{a,b} An-Qi Dai,^b Kun-Lin Huang,*^c Zhi-Hui Zhang,^b Ai-Jun Cui,^b Ming-Yang He,*^b and Qun Chen,*^{a,b}

^a School of Chemical Engineering, Nanjing University of Science and Technology, Nanjing 20094, P. R. China. E-mail: chenqunjpu@yahoo.com.

^a School of Petrochemical Engineering, Key Laboratory of Fine Petro-chemical Technology, Changzhou University, Changzhou 213164, P. R. China. E-mail: hemingyangjpu@yahoo.com.

^b College of Chemistry, Chongqing Normal University, Chongqing 401331, P. R. China

Contents

(1) Va	rious coordinat	tion modes for	r isomeric to	bdc ligands	in 1–8 (Scl	heme S1)	•••••	3
(2)	The	3D sup	oramolecular	archi	tecture	of	1	(Fig.
S1)3								
(3) The	e 2D supramole	ecular networ	k of 2 (Fig.	S2)				4
(4) The	e 3D supramole	ecular networ	k of 2 (Fig.	S3)				4
(5) The	e 2D supramole	ecular networ	k of 3 (Fig.	S4)				5
(6) The intermolecular weak O–H…Cl and Cl…Cl interactions in 4 (Fig. S5)								5
(7) The	e intermolecula	ar weak O–H…	\cdot Cl, C–H $\cdot \cdot \cdot$ C	and Cl…Cl in	nteractions in	n 5 (Fig. So	6)	6
(8) The	e intermolecula	ar weak C–H··	\cdot Cl and Cl \cdots C	Cl interactions	in 6 (Fig. S	57)		6
(9) The intermolecular weak C–H···Cl and Cl···Cl interactions in 7 (Fig. S8)								7
(10) TI	he intermolecu	lar weak Cl…	Cl interaction	s in 8 (Fig. S	9)			7
(11)	A schematic	representati	on of 1D	shape of	f channel	for 4–8	(Fig.	S10)
		8						
(12) Simulated and experimental X-ray powder diffraction patterns for 1–8 (Fig. S11)9								
(13) T	GA curves of c	complexes 1-8	8 (Fig. S12)					10



Scheme S1 Various coordination modes for isomeric tcbdc ligands in 1–8.



Fig. S1 The 3D supramolecular architecture of **1** (Cl···Cl interactions are shown as bright green dashed lines; irrelevant hydrogen atoms are omitted for clarity).



Fig. S2 The 2D supramolecular network of **2** with hydrogen-bonding interactions highlighted in the bright turquoise dashed lines (irrelevant hydrogen atoms are omitted for clarity).



Fig. S3 The 3D supramolecular network of **2** with hydrogen-bonding interactions highlighted in dashed lines (irrelevant hydrogen atoms are omitted for clarity).



Fig. S4 The 2D supramolecular network of 3 with C–H \cdots Cl hydrogen-bonding interactions highlighted in green dashed lines (irrelevant hydrogen atoms are omitted for clarity)



Fig. S5 The intermolecular weak O–H…Cl (turquoise dashed lines)and Cl…Cl (green dashed lines) interactions in **4** (irrelevant hydrogen atoms are omitted for clarity).



Fig. S6 The intermolecular weak O–H…Cl (turquoise dashed lines), C–H…Cl (violet dashed lines) and Cl…Cl (green dashed lines) interactions in **5** (irrelevant hydrogen atoms are omitted for clarity).



Fig. S7 The intermolecular weak $C-H\cdots Cl$ (turquoise dashed lines) and $Cl\cdots Cl$ (green dashed lines) interactions in 6 (irrelevant hydrogen atoms are omitted for clarity).



Fig. S8 The intermolecular weak C–H···O (turquoise dashed lines) and Cl···Cl (green dashed lines) interactions in **7** (irrelevant hydrogen atoms are omitted for clarity).



Fig. S9 The intermolecular weak Cl····Cl interactions (green dashed lines) in **8** (irrelevant hydrogen atoms are omitted for clarity).



Fig. S10 A schematic representation of 1D shape of channel for **4** (a), **5** (b), **6** (c) and **7** (d).



Fig. S 11 Experimental (black) and simulated (red) PXRD patterns of 1-8.



Fig. S12 TGA curves of 1–8.