

Supporting information


Glycine catalyzed diastereoselective domino-synthesis of 6-imino-2,7-dioxabicyclo[3.2.1]octane-4,4,5-tricarbonitriles in water

O. V. Ershov^{a*}, M. Yu. Ievlev^a, V.A. Tafeenko^b and O.E. Nasakin^a


^aChuvash State University, Moskovskiy pr. 15, Cheboksary, Russia.

^bLomonosov Moscow State University, Leninskie gory 1, Moscow, Russia


*E-mail: oleg.ershov@mail.ru.




Pic. 1. ¹H NMR spectrum of **2a** (DMSO-d₆, 299K, 500.13 MHz)




Pic. 2. ¹H NMR spectrum of **2b** (DMSO-d₆, 299K, 500.13 MHz)




Pic. 3. ¹H NMR spectrum of **2c** (DMSO-d₆, 299K, 500.13 MHz)




Pic. 4. ¹H NMR spectrum of **2d** (DMSO-d₆, 299K, 500.13 MHz)




Pic. 5. ¹H NMR spectrum of **2e** (DMSO-d₆, 299K, 500.13 MHz)



Pic. 6. ¹H NMR spectrum of **2f** (DMSO-d₆, 299K, 500.13 MHz)



Pic. 7. ¹H NMR spectrum of **2g** (DMSO-d₆, 299K, 500.13 MHz)



Pic. 8. ¹H NMR spectrum of **2h** (DMSO-d₆, 299K, 500.13 MHz)