

Electronic Supplementary Material (ESI) for New Journal of Chemistry

Iridium-Catalyzed Transfer Hydrogenation of Nitroarenes to Anilines


Shu-jie Chen, Guo-ping Lu, and Chun Cai*

School of Chemical Engineering, Nanjing University of Science and


Technology, Nanjing 210094, China. Fax: +86(25)84315030

c.cai@mail.njust.edu.cn


4-Chloroaniline (Table 2, entry 8) (^1H NMR, CDCl_3 , 500 MHz)




4-Bromoaniline (Table 2, entry 9) (^1H NMR, CDCl_3 , 500 MHz)



4-Methoxyaniline (Table 2, entry 11) (^1H NMR, CDCl_3 , 500 MHz)




3,5-Dichloroaniline (Table 2, entry 12) (¹H NMR, CDCl₃, 500 MHz)




4-Chloro-3-methylaniline (Table 2, entry 13) (^1H NMR, CDCl_3 , 500 MHz)




3-Bromo-4-methylaniline (Table 2, entry 14) (^1H NMR, CDCl_3 , 500 MHz)




2-Chloro-5-methoxyaniline (Table 2, entry 15) (^1H NMR, CDCl_3 , 500 MHz)



2-(1H-Pyrrol-1-yl)aniline (Table 2, entry 19) (^1H NMR, CDCl_3 , 500 MHz)



Quinolin-6-amine (Table 2, entry 21) (^1H NMR, CDCl_3 , 500 MHz)



1-Methyl-1H-indol-5-amine (Table 2, entry 23) (^1H NMR, CDCl_3 , 500 MHz)

