

SUPPLEMENTARY INFORMATION

Naphthyl "capped" triazole-linked calix[4]arene hosts as fluorescent chemosensors towards Fe^{3+} and Hg^{2+} : an experimental and DFT computational study

Shofiur Rahman,*^a Yousif Assiri,^a Abdullah N. Alodhayb,^b

Luc Y. Beaulieu,^b Ahmed K. Oraby,^a and Paris E. Georghiou,*^a

^a Department of Chemistry, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador, Canada A1B3X7.

E-mail: parisg@mun.ca ; msrahman@mun.ca.

^b Department of Physics and Physical Oceanography, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador, Canada A1B3X7. E-mail: lbeaulieu@mun.ca.

Table of Contents

| | <u>Page</u> |
|---|-------------|
| Computational study | S3 |
| Table S1 DFT Calculated binding energies for 6a-d complexes | S4 |
| Table S2 Selected parameter distances for 6a complexes | S5 |
| Table S3 Selected parameter distances for 6b complexes | S5 |
| Table S4 Selected parameter distances for 6c complexes | S6 |
| Table S5 Selected parameter distances for 6d complexes | S6 |
| Figure SI-1: Ball-and-stick structures of 6a and 6a ⊃ Fe^{3+} complex | S7 |
| Figure SI-2: Space fill structures of 6a and 6a ⊃ Fe^{3+} complex | S7 |
| Figure SI-3: Ball-and-stick structures of 6a and 6a ⊃ Hg^{2+} complex | S8 |
| Figure SI-4: Space fill structures of 6a and 6a ⊃ Hg^{2+} complex | S8 |
| Figure SI-5: Ball-and-stick structures of 6b and 6b ⊃ Fe^{3+} complex | S9 |
| Figure SI-6: Space fill structures of 6b and 6b ⊃ Fe^{3+} complex | S9 |
| Figure SI-7: Ball-and-stick structures of 6b and 6b ⊃ Hg^{2+} complex | S10 |
| Figure SI-8: Space fill structures of 6b and 6b ⊃ Hg^{2+} complex | S10 |
| Figure SI-9: Ball-and-stick structures of 6c and 6c ⊃ Fe^{3+} complex | S11 |
| Figure SI-10: Space fill structures of 6c and 6c ⊃ Fe^{3+} complex | S11 |
| Figure SI-11: Ball-and-stick structures of 6c and 6c ⊃ Hg^{2+} complex | S12 |
| Figure SI-12: Space fill structures of 6c and 6c ⊃ Hg^{2+} complex | S12 |