

SUPPLEMENTARY MATERIAL for

Clerodane Diterpenes: Sources, Structures, and Biological Activities

Rongtao Li^{a,b}, Susan L. Morris-Natschke^b, and Kuo-Hsiung Lee^{b,c*}*

CONTENT:

Tables 2–32: Compound Structures arranged by Chemical Classifications


Abbreviations of Functional Groups

Structure Classifications and Sources of Clerodane Diterpenes

1. Type I with an Acyclic Side Chain at C-9

1.1. Type I Subtype I with an *O*-Containing Five-membered Ring at C-18 and C-19

1.1.1. Type I Subtype Ia with the Isozuelanin Skeleton (Table 2)*



| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | Source | Ref. |
|-----|------------------|-----------------|----------------|----------------|------------------|-----------------|----------------|-------------------------------|--|
| 10 | balanspene A | — | — | — | — | — | — | <i>Casearia balansae</i> | <i>J. Nat. Prod.</i> , 2014, 77 , 2182–2189 |
| 11 | balanspene B | =O | αOBu | αOAc | H | H | H | | |
| 12 | balanspene C | αY ₁ | αOBu | αOAc | H | H | H | | |
| 13 | balanspene D | αOBu | αOBu | αOAc | αOMe | H | H | | |
| 14 | balanspene E | αY ₁ | αOBu | αOAc | αOMe | H | H | | |
| 15 | balanspene F | αY ₁ | αOMe | αOAc | αOMe | H | H | | |
| 16 | balanspene G | βOAc | αOMe | αOAc | αOMe | H | H | | |
| 17 | zuelaguidin A | αOH | αOAc | αOAc | αOCin | H | H | <i>Zuelania guidonia</i> | <i>J. Nat. Prod.</i> , 2014, 77 , 455-463 |
| 18 | zuelaguidin B | βOCin | αOAc | αOAc | αOH | H | H | | |
| 19 | zuelaguidin C | αOH | αOAc | αOAc | αX ₁₃ | H | H | | |
| 20 | zuelaguidin D | αOH | αOAc | αOAc | αX ₁₃ | H | αOH | | |
| 21 | casearupestrin A | αOH | αOAc | βOAc | αX ₁ | βOH | H | <i>Casearia rupestris</i> | <i>J. Nat. Prod.</i> , 2011, 74 776–781 |
| 22 | casearupestrin B | αOH | αOAc | βOAc | αOH | βX ₁ | H | | |
| 23 | casearupestrin C | αOH | αOMe | βOAc | αOH | βX ₁ | H | | |
| 24 | casearupestrin D | αOH | αOAc | βOAc | αOAc | βX ₁ | H | | |
| 25 | corymbulosin A | X ₁ | OAc | OAc | OH | H | H | <i>Laetia</i> | <i>Phytochemistry</i> , |

| | | | | | | | | | |
|----|---|-------------------------|--------------|--------------|----------------|-------------|-----|----------------------------|---|
| 26 | corymbulosin B | OH | OAc | OAc | ODc | H | H | <i>corymbulosa</i> | 2000, 55 , 233-236 |
| 27 | corymbulosin C | OH | OAc | OAc | ODc | H | H | | |
| 28 | <i>ent</i> -6 β -hydroxyisozuelanin-2 β -(2-methyl)-butanoate | α Y ₁ | α OAc | β OAc | α OH | H | H | <i>Casearia corymbosa</i> | <i>Phytochemistry</i> , 1990, 29 , 3591-3595 |
| 29 | <i>ent</i> -6 β -methoxyisozuelanin-2 β -(2-methyl)-butanoate | α Y ₁ | α OAc | β OAc | α OMe | H | H | | |
| 30 | <i>ent</i> -2 β -(2-methyl)butoxy-3,4-dihydro-4 α -isozuelanin | α Y ₁ | α OAc | β OAc | H | H | H | | |
| 31 | <i>ent</i> -6 β -hydroxyisozuelanin-2 β -(2-methyl)-propanoate | α OiBu | α OAc | β OAc | α OH | H | H | | |
| 32 | <i>ent</i> -6 β -methoxyisozuelanin-2 β -(2-methyl)-propanoate | α OiBu | α OAc | β OAc | α OMe | H | H | | |
| 33 | <i>ent</i> -2 β -hydroxy-3,4-dihydro-4 α -isozuelanin | α OH | α OAc | β OAc | H | H | H | | |
| 34 | <i>ent</i> -2 β -acetoxy-3,4-dihydro-4 α -isozuelanin | α OAc | α OAc | β OAc | H | H | H | | |
| 35 | intrapetacin A | β Z ₁ | α OMe | α OAc | α OH | H | H | <i>Licania</i> | <i>J. Nat. Prod.</i> , |
| 36 | intrapetacin B | β Z ₁ | α OAc | α OAc | α OH | H | H | <i>intrapetiolaris</i> | 2001, 64 , 497-501 |
| 37 | caseanigrescen A | α OBu | α OAc | α OAc | α OH | β OAc | H | <i>Casearia nigrescens</i> | <i>J. Nat. Prod.</i> , |
| 38 | caseanigrescen B | α OBu | α OAc | α OAc | α OH | β OH | H | | |
| 39 | caseanigrescen C | α OBu | α OAc | α OAc | α OAc | β OH | H | | |
| 40 | caseanigrescen D | α OBu | α OAc | α OAc | α OH | H | H | | |
| 41 | argutin F | α X ₁ | α OAc | α OAc | OH | β H | OOH | <i>Casearia arguta</i> | <i>J. Nat. Prod.</i> , |
| 42 | argutin G | α X ₁ | α OAc | α OAc | OH | β OH | OOH | | |
| 43 | argutin H | α OH | α OAc | α OAc | X ₁ | β OH | OOH | | |
| 44 | <i>rel</i> -2 β -acetoxyisozuelanin-6 β -cinnamate | β OAc | β OAc | α OAc | β OCin | H | H | <i>Zuelania guidonia</i> | <i>Phytochemistry</i> , |
| 45 | isozuelanin** | H | β OAc | α OAc | H | H | H | | |
| 46 | <i>rel</i> -18(<i>S</i>),19(<i>R</i>)-diacetoxy-18,19-epoxy-6(<i>R</i>)-methoxy-2(<i>S</i>)-(2 ζ -methylbutanoyloxy)-5(<i>R</i>),8(<i>S</i>),9(<i>S</i>),10(<i>R</i>)-cleroda-3,13(16),14-triene | β Y ₁ | β OAc | β OAc | β OMe | H | H | <i>Casearia tremula</i> | <i>Phytochemistry</i> , |
| 47 | <i>rel</i> -18(<i>S</i>),19(<i>R</i>)-diacetoxy-18,19-epoxy-2(<i>S</i>)-(2 ζ -methylbutanoyloxy)-5(<i>R</i>),8(<i>S</i>),9(<i>S</i>),10(<i>R</i>)-cleroda-3,13(16),14-triene | β Y ₁ | β OAc | β OAc | β H | H | H | | |
| 48 | <i>rel</i> -18(<i>S</i>),19(<i>R</i>)-diacetoxy-18,19-epoxy-6(<i>R</i>)-hydroxy-2(<i>S</i>)-(2 ζ -methylbutanoyloxy)-5(<i>R</i>),8(<i>S</i>),9(<i>S</i>),10(<i>R</i>)-cleroda-3,13(16),14-triene | β Y ₁ | β OAc | β OAc | β OH | H | H | | |
| 49 | <i>rel</i> -18(<i>S</i>),19(<i>R</i>)-diacetoxy-18,19-epoxy-6(<i>R</i>)-hydroxy-2(<i>S</i>)-undecanoyloxy-5(<i>R</i>),8(<i>S</i>),9(<i>S</i>),10(<i>R</i>)-cleroda-3,13(16),14-triene | β X ₂ | β OAc | β OAc | β OH | H | H | | |
| 50 | <i>rel</i> -18(<i>S</i>),19(<i>R</i>)-diacetoxy-18,19-epoxy-6(<i>R</i>)-hydroxy-2(<i>S</i>)-octanoyloxy-5(<i>R</i>),8(<i>S</i>),9(<i>S</i>),10(<i>R</i>)-cleroda-3,13(16),14-triene | β OOct | β OAc | β OAc | β OH | H | H | | |
| 51 | <i>rel</i> -18(<i>S</i>),19(<i>R</i>)-diacetoxy-18,19-epoxy-6(<i>R</i>)-hydroxy-2(<i>S</i>)-(3 ζ -hydroxyoctanoyloxy)-5(<i>R</i>),8(<i>S</i>),9(<i>S</i>),10(<i>R</i>)-cleroda-3,13(16),14-triene | β X ₃ | β OAc | β OAc | β OH | H | H | | |
| 52 | <i>rel</i> -18(<i>S</i>),19(<i>R</i>)-diacetoxy-18,19-epoxy-2(<i>R</i>)-hexanoyloxy-5(<i>R</i>),8(<i>S</i>),9(<i>S</i>),10(<i>R</i>)-cleroda-3,13(16),14-triene | α X ₄ | β OAc | β OAc | H | H | H | | |


| | | | | | | | | | |
|----|---|-------------------------|-------------|--------------|-------------|-------------|-------------|--|---|
| 53 | casearlucin B | α OAc | β OAc | β OAc | β OMe | H | H | <i>Casearia lucida</i> | <i>J. Nat. Prod.</i> , 2002, 65 , 100-107 |
| 54 | casearlucin D | β Y ₁ | β OAc | β OAc | β OAc | H | H | | |
| 55 | casearlucin H | β Y ₁ | β OAc | β OAc | β OH | H | α OH | | |
| 56 | casearlucin I | β Y ₁ | β OAc | β OAc | β OH | H | β OH | | |
| 57 | casearlucin M | β Y ₁ | β OAc | β OAc | β OMe | H | H | | |
| 58 | casearlucin J | α Y ₁ | β OAc | β OAc | β OH | H | α OH | | |
| 59 | casearlucin K | α Y ₁ | β OAc | β OAc | β OH | H | β OH | | |
| 60 | caseamembrin A | β Y ₁ | β OBu | β OAc | β OH | H | H | <i>Casearia membranacea</i> | <i>J. Nat. Prod.</i> , 2004, 67 , 316-321 |
| 61 | caseamembrin B | β Y ₁ | β OMe | β OAc | β OH | H | H | | |
| 62 | caseamembrin C | β Y ₁ | β OBu | β OAc | β OH | β OH | H | | |
| 63 | caseamembrin D | β Y ₁ | β OBu | β OAc | β OH | β OAc | H | | |
| 64 | caseamembrin E | α Y ₁ | β OAc | β OAc | β OH | H | H | | |
| 65 | caseamembrin M | β Y ₁ | β OAc | β OAc | β OBu | H | H | | |
| 66 | caseamembrin N | β Y ₁ | β OAc | β OAc | β OH | β OAc | H | | |
| 67 | caseamembrin O | β Y ₂ | β OAc | β OAc | β OH | β OAc | H | <i>Chem. Pharm. Bull.</i> , 2004, 52 , 108-110 | |
| 68 | caseamembrin B | α Y ₁ | β OAc | β OAc | β OH | H | β OH | | |
| 69 | <i>rel</i> -2(<i>R</i>),18(<i>S</i>),19(<i>R</i>)-triacetoxy-18,19-epoxy-4(<i>S</i>),5(<i>S</i>),9(<i>S</i>),10(<i>R</i>)-clerodan-13(16),14-dien-6-one | α OAc | β OAc | β OAc | =O | H | H | <i>Casearia grayi</i> | <i>Nat. Prod. Commun.</i> , 2006, 1 , 441-448 |
| 70 | <i>rel</i> -2(<i>R</i>),18(<i>S</i>),19(<i>R</i>)-tri-acetoxy-18,19-epoxy-4(<i>S</i>),5(<i>R</i>),8(<i>S</i>),9(<i>S</i>),10(<i>R</i>)-clerodan-13(16),14-diene | α OAc | β OAc | β OAc | H | H | H | | |
| 71 | <i>rel</i> -18(<i>S</i>),19(<i>R</i>)-diacetoxy-18,19-epoxy-2(<i>R</i>)-(2 ξ -methylbutanoyl)-4(<i>S</i>),-5(<i>S</i>),9(<i>S</i>),10(<i>R</i>)-clerodan-13(16),14-dien-6-one | α Y ₁ | β OAc | β OAc | =O | H | H | | |
| 72 | <i>rel</i> -18(<i>S</i>),19(<i>R</i>)-diacetoxy-18,19-epoxy-2(<i>R</i>)-isobutanoyl-4(<i>S</i>),5(<i>R</i>),9(<i>S</i>),-10(<i>R</i>)-clerodan-13(16),14-diene | α OiBu | β OAc | β OAc | H | H | H | | |
| 73 | <i>rel</i> -18(<i>S</i>),19(<i>R</i>)-diacetoxy-18,19-epoxy-2(<i>R</i>)-(2 ξ -methylbutanoyl)-4(<i>S</i>),5(<i>R</i>),9(<i>S</i>),10(<i>R</i>)-clerodan-13(16),14-diene | α Y ₁ | β OAc | β OAc | H | H | H | | |
| 74 | caseargrewiin A | β Y ₂ | β OAc | β OAc | β OMe | H | H | <i>Casearia grewifolia</i> | <i>J. Nat. Prod.</i> , 2005, 68 , 183-188 |
| 75 | 6 β -hydroxyisozuelanin-2 β -benzoate | β OBz | β OAc | α OAc | β OH | H | H | <i>Zuelania guidonia</i> | <i>Phytochemistry</i> , 1990, 29 , 2939-2942 |
| 76 | esculentin A | =O | β OAc | β OAc | H | H | H | <i>Casearia</i> | <i>Indian J. Chem.</i> , |

| | | | | | | | | | | |
|-----------|-----------------------|------------------------|-------------|-----|-----|---|---|--|------------------------------|--|
| | | | | | | | | | <i>esculenta</i> | 2002, 41B , 2706-2708 |
| 77 | caseabalansin C | α OAc | OEt | OAc | OMe | — | — | | <i>Casearia balansae</i> | <i>J. Nat. Prod.</i> , 2013, 76 , 1573-1579 |
| 78 | 2-epicaseabalansin C | β OAc | OEt | OAc | OMe | — | — | | | |
| 79 | caseabalansin D | β Y ₁ | OEt | OAc | OH | — | — | | | |
| 80 | caseabalansin E | β Y ₁ | OMe | OH | OMe | — | — | | | |
| 81 | caseabalansin F | =O | OAc | OAc | OMe | — | — | | | |
| 82 | caseabalansin G | =O | OBu | OAc | OMe | — | — | | | |
| 83 | caseabalansin B | α OH | — | — | — | — | — | | | |
| 84 | 2-epicaseabalansin B | β OH | — | — | — | — | — | | | |
| 85 | caseabalansin A | — | β OH | — | — | — | — | | | |
| 86 | 18-epicaseabalansin A | — | α OH | — | — | — | — | | | |

* It should be noted that the absolute stereochemistry has not been determined in all cases. For example, esculentin A (**76**) has been reported in both *ent-neo*¹⁶ and *neo*³² configurations. In addition, caseargrewiin A (**74**) was shown as a *neo*-clerodane, while the co-isolated caseargrewiins B–D (**1256–1258**) were shown as *ent-neo*-clerodanes, with the absolute configuration of C-2 in **1258** established by a modified Mosher's method, NMR coupling constants, and NOESY correlations.³¹ Generally, this review has focused on relative configurations only.

** Not an isolated compound


1.1.2. Type I Subtype Ib with the Zuelanin Skeleton (Table 3)



| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | Source | Ref. |
|-----|-------------------|-----------------|----------------|----------------|-----------------|----------------|----------------|-----------------------------|---|
| 87 | casearvestrin A | βOiBu | αOAc | αOAc | αOH | H | — | <i>Casearia sylvestris</i> | <i>J. Nat. Prod.</i> , 2002, 65 , 95-99 |
| 88 | casearvestrin B | βY ₁ | αOAc | αOAc | αOH | H | — | | |
| 89 | casearvestrin C | βX ₄ | αOAc | αOAc | αOH | H | — | | |
| 90 | argutin A | αX ₁ | αOAc | αOAc | OH | H | — | <i>Casearia arguta</i> | <i>J. Nat. Prod.</i> , 2010, 73 , 2013-2018 |
| 91 | argutin B | αOH | αOAc | αOAc | X ₁ | H | — | | |
| 92 | argutin C | αX ₁ | αOAc | αOAc | OH | OH | — | | |
| 93 | argutin D | αOH | αOAc | αOAc | X ₁ | OH | — | | |
| 94 | argutin E | αX ₁ | αOAc | αOAc | H | OH | — | | |
| 95 | esculentin B | αY ₂ | βOAc | βOAc | βOH | αOH | — | <i>Casearia esculenta</i> | <i>Indian J. Chem.</i> , 2002, 41B , 2706-2708 |
| 96 | casearborin A | αZ ₁ | αOAc | αOAc | αH | H | — | <i>Casearia arborea</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 657-661 |
| 97 | casearborin B | αZ ₂ | αOAc | αOAc | αH | H | — | | |
| 98 | casearborin C | αOH | αOAc | αOAc | αZ ₁ | H | — | | |
| 99 | casearborin D | αZ ₁ | αOAc | αOAc | αOH | H | — | | |
| 100 | casearborin E | αOAc | αOAc | αOAc | αZ ₁ | H | — | | |
| 101 | laetiaprocerine A | αY ₁ | βOAc | βOAc | αOH | H | — | <i>Laetia procera</i> | <i>Bioorg. Med. Chem. Lett.</i> , 2005, 15 , 5065-5070 |
| 102 | laetiaprocerine B | αOiBu | βOAc | βOAc | αOBz | H | — | | |
| 103 | laetiaprocerine C | αY ₁ | βOAc | βOAc | αOBz | H | — | | |
| 104 | caseargrewiin E | αOBu | αOAc | αOAc | H | H | — | <i>Casearia grewiifolia</i> | <i>J. Nat. Prod.</i> , 2007, 70 , 1122-1126 |
| 105 | caseargrewiin F | αOBu | αOAc | αOAc | αOH | H | — | | |
| 106 | caseargrewiin G | αOBu | αOMe | αOAc | αOH | H | — | | |
| 107 | caseargrewiin H | αX ₄ | αOAc | αOAc | H | H | — | | |
| 108 | caseargrewiin I | αX ₄ | αOAc | αOAc | αOH | H | — | | |

| | | | | | | | | | |
|-----|--|---------------|--------------|--------------|--------------|---------------|--------------------|-----------------------------|---|
| 109 | caseargrewiin J | αX_4 | αOMe | αOAc | αOH | H | — | | |
| 110 | caseargrewiin K | αX_4 | αOMe | αOAc | αOH | βOH | — | | |
| 111 | caseargrewiin L | αY_2 | αOAc | αOAc | αOH | H | — | | |
| 112 | <i>rel-2α-hydroxyzuelanin-6β-cinnamate</i> | αOH | βOAc | αOAc | $\beta OCin$ | H | Me | <i>Zuelania guidonia</i> | <i>Phytochemistry</i> , 1990, 29 , 1609-1614 |
| 113 | <i>rel-6β-hydroxyzuelanin-2α-cinnamate</i> | $\alpha OCin$ | βOAc | αOAc | βOH | H | Me | | |
| 114 | <i>rel-2β-hydroxyzuelanin-6β-cinnamate</i> | βOH | βOAc | αOAc | $\beta OCin$ | H | Me | | |
| 115 | casearlucin A | βY_1 | βOAc | βOAc | βOH | H | Me | <i>Casearia lucida</i> | <i>J. Nat. Prod.</i> , 2002, 65 , 100-107 |
| 116 | casearlucin C | βY_1 | βOAc | βOAc | βOAc | H | Me | | |
| 117 | casearlucin F | βY_1 | βOAc | βOAc | H | H | Me | | |
| 118 | casearlucin G | αY_1 | βOAc | βOAc | H | H | Me | | |
| 119 | caseamembrol A | αY_1 | βOAc | βOAc | βOH | H | Me | <i>Casearia membranacea</i> | <i>Chem. Pharm. Bull.</i> , 2004, 52 , 108-110 |
| 120 | <i>6β-hydroxyzuelanin-2α-acetate</i> | αOAc | βOAc | αOAc | βOH | H | Me | <i>Zuelania guidonia</i> | <i>Phytochemistry</i> , 1990, 29 , 2939-2942 |
| 121 | <i>6β-hydroxyzuelanin-2α-<i>n</i>-octacetate</i> | $\alpha OOct$ | βOAc | αOAc | βOH | H | Me | | |
| 122 | <i>6β-hydroxyzuelanin-2β-<i>n</i>-octacetate</i> | $\beta OOct$ | βOAc | αOAc | βOH | H | Me | | |
| 123 | <i>6β-hydroxyzuelanin-2β-<i>n</i>-benzoate</i> | βOBz | βOAc | αOAc | βOH | H | Me | | |
| 124 | <i>2α-hydroxyzuelanin-6β-<i>n</i>-benzoate</i> | αOH | βOAc | αOAc | βOBz | H | Me | | |
| 125 | <i>2α-hydroxyzuelanin-6β-<i>n</i>-(3-hydroxy)-octanoate</i> | αOH | βOAc | αOAc | βX_8 | H | Me | | |
| 126 | <i>zuelanin-2β-benzoate</i> | βOBz | βOAc | αOAc | H | H | Me | | |
| 127 | caseobliquin A | αOAc | βOAc | αOAc | βZ_1 | H | Me | <i>Casearia obliqua</i> | <i>J. Nat. Prod.</i> , 2009, 72 , 1847-1850 |
| 128 | caseobliquin B | βOAc | βOAc | αOAc | $\beta OCin$ | H | Me | | |
| 129 | <i>rel-18(S),19(R)-diacetoxo-18,19-epoxy-2(R),7(S),16-trihydroxy-6(S)-myristoyloxy-5(R),8(R),9(S),10(R)-cleroda-3,12,14-triene</i> | βOH | βOAc | βOAc | $\beta OMyr$ | αOH | CH ₂ OH | <i>Laetia procera</i> | <i>Phytochemistry</i> , 1996, 43 , 635-638 |
| 130 | <i>rel-18(S),19(R)-diacetoxo-18,19-epoxy-2(R),7(S),16-trihydroxy-6(S)-palmitoyloxy-5(R),8(R),9(S),10(R)-cleroda-3,12,14-triene</i> | βOH | βOAc | βOAc | $\beta OPal$ | αOH | CH ₂ OH | | |
| 131 | <i>rel-18(S),19(R)-diacetoxo-18,19-epoxy-2(R),6(S),16-trihydroxy-7(S)-myristoyloxy-5(R),8(R),9(S),10(R)-cleroda-3,12,14-triene</i> | βOH | βOAc | βOAc | βOH | $\alpha OMyr$ | CH ₂ OH | | |
| 132 | <i>rel-18(S),19(R)-diacetoxo-18,19-epoxy-2(R),7(S),16-trihydroxy-6(S)-palmitoyloxy-5(R),8(R),9(S),10(R)-cleroda-3,12,14-triene</i> | βOH | βOAc | βOAc | βOH | $\alpha OPal$ | CH ₂ OH | | |
| 133 | bucidasarin A | $\beta OiBu$ | βOAc | βOAc | βOH | H | Me | <i>Bucida buceras</i> | <i>Bioorg. Med. Chem. Lett.</i> , 2002, 12 , 345-348 |
| 134 | bucidasarin B | βY_1 | βOAc | βOAc | βOH | H | Me | | |
| 135 | bucidasarin C | $\beta OiBu$ | βOAc | βOAc | H | H | Me | | |




1.1.3. Type I Subtype Ic with Other Skeletons (Table 4)



| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | Source | Ref. |
|-----|--|----------------|----------------|----------------|----------------------|----------------------|----------------|----------------------------|---|
| 136 | 15-malonyloxy- <i>ent</i> -cleroda-3,13 <i>E</i> -dien-18,19-olide | H | =O | H | Me | CH ₂ OMal | Me | <i>Olearia teretifolia</i> | <i>Phytochemistry</i> , 1992, 31 , 1703-1711 |
| 137 | 18,19-epoxy-18β-methoxy- <i>ent</i> -cleroda-3,13 <i>E</i> -dien-15-oic acid | H | βOMe | H | Me | CO ₂ H | Me | | |
| 138 | 18,19-epoxy-18α-methoxy- <i>ent</i> -cleroda-3,13 <i>E</i> -dien-15-oic acid | H | αOMe | H | Me | CO ₂ H | Me | | |
| 139 | <i>ent</i> -clerod-3-en-15-oic acid-18,19-olide | H | =O | H | CO ₂ H | — | — | | |
| 140 | 15-malonyloxy- <i>ent</i> -clerod-3-en-18,19-olide | H | =O | H | CH ₂ OMal | — | — | | |
| 141 | 7α-hydroxy- <i>ent</i> -clerod-3-en-15-oic acid-18, 19-olide | H | =O | OH | CO ₂ H | — | — | | |
| 142 | 15-malonyloxy-7α-hydroxy- <i>ent</i> -clerod-3-en-18,19-olide | H | =O | OH | CH ₂ OMal | — | — | | |
| 143 | 2α-hydroxy- <i>ent</i> -cleroda-3-en-15-oic acid-18,19-olide | αOH | =O | H | CO ₂ H | — | — | | |
| 144 | 2β-hydroxy- <i>ent</i> -clerod-3-en-15-oic acid-18,19-olide | βOH | =O | H | CO ₂ H | — | — | | |
| 145 | 15-hydroxy- <i>ent</i> -clerod-3-en-18,19-olide | H | =O | H | CH ₂ OH | — | — | | |
| 146 | 7α,15-dihydroxy- <i>ent</i> -clerod-3-en-18,19-olide | H | =O | OH | CH ₂ OH | — | — | | |
| 147 | 18,19-epoxy-18β-methoxy- <i>ent</i> -clerod-3-en-15-oic acid | H | βOMe | H | CO ₂ H | — | — | | |
| 148 | 18,19-epoxy-18α-methoxy- <i>ent</i> -clerod-3-en-15-oic acid | H | αOMe | H | CO ₂ H | — | — | | |
| 149 | 4β-hydroxy- <i>ent</i> -clerod-2-en-15-oic acid-18,19-olide | — | — | — | — | — | — | | |
| 150 | baclinal | — | — | — | — | — | — | <i>Baccharis linearis</i> | <i>Phytochemistry</i> , 1996, 41 , 1123-1127 |
| 151 | 13- <i>epi</i> -baclinepoxide | — | — | — | — | — | — | | |
| 152 | baclinepoxide | — | — | — | — | — | — | <i>Zuelania guidonia</i> | <i>J. Nat. Prod.</i> , 2014, 77 , 455-463 |
| 153 | zuelaguidin E | OH | OCin | H | — | — | — | | |
| 154 | zuelaguidin G | Dc | OXYl | βOH | — | — | — | | |
| 155 | zuelaguidin H | Dc | OXYl | αOH | — | — | — | | |

1.2. Type I Subtype II with a Double Bond Between C-3 and C-4 or Another Position

1.2.1. Type I Subtype IIa with a Double Bond between C-3 and C-4 (Table 5)



| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | R ₇ | R ₈ | Source | Ref. |
|-----|--|----------------|--------------------|----------------|----------------|----------------|----------------|----------------|----------------|---------------------------|---|
| 156 | 2-oxo-18-hydroxy-10 α ,17 α ,19 α ,20 β -(-)-cleroda-3,13(16),14-triene | =O | CH ₂ OH | α Me | H | α Me | — | — | — | <i>Casearia corymbosa</i> | <i>Phytochemistry</i> , 1990, 29 , 3591-3595 |

| | | | | | | | | | | | |
|-----|---|-------------------------|---------------------|------------------------------|-------------------|-----------------------------|-----------------------------|--------------------|----|---|---|
| 157 | 2-oxo-18,19-diacetoxy-10 α ,17 α ,19 α ,20 β -(-)-cleroda-3,13(16),14-triene | =O | CH ₂ OAc | α CH ₂ OAc | H | α Me | — | — | — | | |
| 158 | balanspene H | α Y ₁ | CHO | α CHO | α OMe | α Me | — | — | — | <i>Casearia balansae</i> | <i>J. Nat. Prod.</i> , 2014, 77 , 2182–2189 |
| 159 | terpentetriene | H | Me | β Me | H | α Me | — | — | — | <i>Streptomyces lividans</i> transformant | <i>J. Bacteriol.</i> , 2001, 183 , 6085-6094 |
| 160 | terpentecin | — | — | — | — | — | — | — | — | <i>Streptomyces griseolosporeus</i> | |
| 161 | ent-clerod-3,13(16),14-triene | H | Me | α Me | H | α Me | — | — | — | <i>Jungermannia infusca</i> | <i>Phytochemistry</i> , 1998, 49 , 601-608 |
| 162 | ent-clerod-3,13(16),14-triene-17-ol | H | Me | α Me | H | α CH ₂ OH | — | — | — | | |
| 163 | caseamembrin F | β Y ₁ | CHO | β CHO | β OH | β Me | — | — | — | <i>Casearia membranacea</i> | <i>J. Nat. Prod.</i> , 2004, 67 , 316-321 |
| 164 | (-)-3,12Z,14-cis-clerodatrien-18-oic acid | CO ₂ H | β Me | H | Me | — | — | — | — | <i>Schistochila acuminata</i> | <i>J. Chin. Chem. Soc.</i> , 1992, 39 , 263-266 |
| 165 | heteroscyphic acid A | Me | α Me | H | CO ₂ H | — | — | — | — | <i>Heteroscyphus planus</i> | <i>Phytochemistry</i> , 1994, 37 , 1263-1268 |
| 166 | heteroscyphic acid B | Me | α Me | OAc | CO ₂ H | — | — | — | — | | |
| 167 | heteroscyphol | — | — | — | — | — | — | — | — | | <i>Phytochemistry</i> , 1995, 38 , 119-127 |
| 168 | 6 α -hydroxy-3,12E,14-clerodatriene | H | Me | α Me | α OH | α Me | — | — | — | <i>Heteroscyphus billardieri</i> | <i>Chem. Pharm. Bull.</i> , 2004, 52 , 556-560 |
| 169 | (-)-3,12E,14-cis-clerodatrien-18-oic acid | H | CO ₂ H | β Me | H | α Me | — | — | — | <i>Schistochila acuminata</i> | <i>J. Chin. Chem. Soc.</i> , 1992, 39 , 263-266 |
| 170 | bucidasin D | =O | CH ₂ OAc | β CH ₂ OAc | H | β Me | — | — | — | <i>Bucida buceras</i> | <i>Bioorg. Med. Chem. Lett.</i> , 2002, 12 , 345-348 |
| 171 | 3,12E-kolavadien-15-oic acid-16-al | — | — | — | — | — | — | — | — | <i>Polyalthia viridis</i> | <i>Phytochemistry</i> , 1993, 34 , 457-460 |
| 172 | 18,19-dihydroxy-ent-cleroda-3,13E-dien-15-oic acid | H | CH ₂ OH | α CH ₂ OH | H | H | α Me | CO ₂ H | Me | <i>Olearia teretifolia</i> | <i>Phytochemistry</i> , 1992, 31 , 1703-1711 |
| 173 | platypodiol | H | Me | α Me | β OH | H | α CH ₂ OH | CH ₂ OH | Me | <i>Baccharis platypoda</i> | <i>Tetrahedron Lett.</i> , 2014, 55 , 4898-4900 |
| 174 | 17-hydroxy-3,13E-clerodadien-15-al | H | Me | α Me | H | H | α CH ₂ OH | CHO | Me | <i>Jungermannia infusca</i> | <i>Chem. Pharm. Bull.</i> , 2000, 48 , 1818-1821 |
| 175 | 6 α ,7 α -dihydroxykolavenol | H | Me | α Me | α OH | α OH | α Me | CH ₂ OH | Me | <i>Ptychopetalum olacoides</i> | <i>Nat. Prod. Commun.</i> , 2011, 6 , 327-332 |
| 176 | 2 β -acetoxy-19-carboxymethyl-cleroda-3,13-dien-15-oic acid | β OAc | CO ₂ Me | α Me | H | H | α Me | CO ₂ H | Me | <i>Scapania bolandeli</i> | <i>Phytochemistry</i> , 1999, 52 , 1551-1553 |
| 177 | deserticolic acid | H | Me | β CH ₂ OH | H | H | α Me | CO ₂ H | Me | <i>Haplopappus deserticola</i> | <i>Phytochemistry</i> , 1999, 52 , 1531-1533 |
| 178 | — | =O | Me | α Me | H | α OAc | α Me | CO ₂ Me | Me | <i>Solidago altissima</i> | <i>Phytochemistry</i> , 1999, 52 , 487-493 |


| | | | | | | | | | | | |
|-----|--|--------------------|---------------------|---------------------|---------------------|-------------------|--------------------|--------------------|--------------------|----------------------------------|--|
| 179 | — | βOH | Me | αMe | H | αOAc | αMe | CO ₂ Me | Me | | |
| 180 | — | αOH | Me | αMe | H | H | αMe | CO ₂ Me | Me | | |
| 181 | — | βOH | Me | αMe | H | H | αMe | CO ₂ Me | Me | | |
| 182 | diastereoisomer of kolavenol | H | Me | αMe | H | H | βMe | CH ₂ OH | Me | <i>Entada abyssinica</i> | <i>J. Ethnopharmacol.</i> , 1998, 61 , 179-183 |
| 183 | 16-oxo- <i>ent</i> -cleroda-3,13Z-diene-15,17-dioic acid | H | Me | αMe | H | H | αCO ₂ H | CO ₂ H | CHO | <i>Diplostephium floribundum</i> | <i>Phytochemistry</i> , 1992, 31 , 213-216 |
| 184 | 15-oxo- <i>ent</i> -cleroda-3,13Z-diene-16,17-dioic acid | H | Me | αMe | H | H | αCO ₂ H | CHO | CO ₂ H | | |
| 185 | 2α-acetoxy-15-oxo- <i>ent</i> -cleroda-3,13Z-diene-16,17-dioic acid | αOAc | Me | αMe | H | H | αCO ₂ H | CHO | CO ₂ H | | |
| 186 | 7-oxo- <i>ent</i> -clerodan-3,13E-dien-15-oic acid | H | Me | αMe | H | =O | αMe | CO ₂ H | Me | <i>Platychaete aucheri</i> | <i>Phytochemistry</i> , 1990, 29 , 985-987 |
| 187 | 6α-hydroxy-7-oxo- <i>ent</i> -clerodane-3,13E-dien-15-oic acid | H | Me | αMe | αOH | =O | αMe | CO ₂ H | Me | | |
| 188 | crotonic acid | H | CH ₂ OAc | βMe | H | H | αMe | CO ₂ H | Me | <i>Croton chilensis</i> | <i>Bol. Soc. Chil. Quim.</i> , 1995, 40 , 157-162 |
| 189 | polyalthialdoic acid | H | Me | αMe | H | H | αMe | CO ₂ H | CHO | <i>Polyalthia longifolia</i> | <i>Planta Med.</i> , 1991, 57 , 380-383 |
| 190 | 18-hydroxy-5,10- <i>trans</i> -cleroda-3,13E-dien-15-oic acid methyl ester | H | CH ₂ OH | αMe | H | H | αMe | CO ₂ Me | Me | <i>Heteroscyphus planus</i> | <i>Phytochemistry</i> , 1996, 41 , 581-587 |
| 191 | pilosanol A | H | Me | αCH ₂ OH | αOH | H | αMe | CH ₂ OH | CH ₂ OH | <i>Portulaca pilosa</i> | <i>Phytochemistry</i> , 1991, 30 , 4075-4077 |
| 192 | pilosanol B | H | CH ₂ OH | αCH ₂ OH | αOH | H | αMe | CH ₂ OH | CH ₂ OH | | |
| 193 | methyl(13E)-2-oxo-neocleroda-3,13-dien-15-oate | =O | Me | αMe | H | H | αMe | CO ₂ Me | Me | <i>Amoora yunnanensis</i> | <i>Helv. Chim. Acta</i> , 2004, 87 , 1279-1286. |
| 194 | (13E)-2-oxoneocleroda-3,13-dien-15-ol | =O | Me | αMe | H | H | αMe | CH ₂ OH | Me | | |
| 195 | (-)-(5R,8S,9S,10R)-cleroda-3,13E-dien-15-oic acid | H | Me | αMe | H | H | βMe | CO ₂ H | Me | <i>Hymenaea courbaril</i> | <i>Phytochemistry</i> , 2001, 58 , 1153-1157 |
| 196 | methyl (5S,8S,9S,10R)-cleroda-3,13E-dien-15-oate | H | Me | βMe | H | H | βMe | CO ₂ Me | Me | | |
| 197 | 5α,8α-2-oxokolavenic acid | βMe | — | — | — | — | — | — | — | <i>Detarium microcarpum</i> | <i>J. Nat. Prod.</i> , 2006, 69 , 768-773 |
| 198 | (13E)-2-oxo-5α- <i>cis</i> -17α,20α-cleroda-3,13-diene-15-oic acid | αMe | — | — | — | — | — | — | — | <i>Eperua purpurea</i> | <i>Phytochemistry</i> , 1991, 30 , 3474-3475 |
| 199 | portulene acetal | CH ₂ OH | CH ₂ OH | — | — | — | — | — | — | <i>Portulaca grandiflora</i> | <i>J. Nat. Prod.</i> , 1997, 60 , 912-914 |
| 200 | solidagocanin A | H | αMe | βOTig | αMe | CO ₂ H | Me | — | — | <i>Solidago canadensis</i> | <i>Helv. Chim. Acta</i> , 2012, 95 , 1121-1125 |
| 201 | solidagocanin B | H | αMe | βOAng | αMe | CO ₂ H | Me | — | — | | |
| 202 | 17-hydroxy-3,13Z-clerodadien-15-al | H | αMe | H | αCH ₂ OH | CHO | Me | — | — | <i>Jungermannia infusca</i> | <i>Chem. Pharm. Bull.</i> , 2000, 48 , 1818-1821 |

| | | | | | | | | | | | |
|-----|--|--------------|-----------------------------|-------------------------------|-----------------------------|-------------------|-------------|-------------------------|---------------------|---------------------------------|--|
| 203 | PMS-1 | H | α Me | H | α CH ₂ OH | Me | Me | — | — | <i>Brazilian propolis</i> | <i>Anticancer Res.</i> , 1996, 16 , 2669-2672 |
| 204 | 16-oxo-cleroda-3,13(14) <i>E</i> -diene-15-oic acid | H | α Me | H | α Me | CO ₂ H | CHO | — | — | <i>Polyalthia longifolia</i> | <i>Fitoterapia</i> , 2005, 76 , 336-339 |
| 205 | 2 α -hydroxy- <i>cis</i> -cleroda-3,13(<i>Z</i>),8,17-trien-15-oic acid | α OH | β Me | H | =CH ₂ | CO ₂ H | Me | — | — | <i>Haplopappus foliosus</i> | <i>Planta Med.</i> , 2003, 69 , 675-677 |
| 206 | 2 α -acetoxy- <i>cis</i> -cleroda-3,13(<i>Z</i>),8,17-trien-15-oic acid | α OAc | β Me | H | =CH ₂ | CO ₂ H | Me | — | — | | |
| 207 | trinerdiol | H | CH ₂ OH | α Me | H | H | α Me | CH ₂ OH | Me | <i>Baccharis trinervis</i> | <i>Phytochemistry</i> , 1993, 34 , 1377-1384 |
| 208 | 15-acetyl-trinerdiol | H | CH ₂ OH | α Me | H | H | α Me | CH ₂ OA c | Me | | |
| 209 | 18-acetyl-trinerdiol | H | CH ₂ OAc | α Me | H | H | α Me | CH ₂ OH | Me | | |
| 210 | 18-methylmalonyl-trinerdiol | H | CH ₂ OMe- Mal | α Me | H | H | α Me | CH ₂ OH | Me | | |
| 211 | 15,18-diacetyl-triner-triol | H | CH ₂ OAc | α Me | H | H | α Me | CH ₂ OA c | CH ₂ OH | | |
| 212 | 14,15,18-triacetyl-trinertetrol | H | CH ₂ OAc | α Me | H | α OH | α Me | CH ₂ OA c | CH ₂ OAc | | |
| 213 | 15,16-diacetyl-trinertriol | H | CH ₂ OH | α Me | H | H | α Me | CH ₂ OA c | CH ₂ OAc | | |
| 214 | 15-hydroxy-16-acetoxy- <i>ent</i> -clerod-3- <i>en</i> -18-oic acid | H | CO ₂ H | α Me | H | H | α Me | CH ₂ OH | CH ₂ OAc | <i>Baccharis gaudichaudiana</i> | <i>Chem Pharm Bull</i> , 2007, 55 , 1532-1534 |
| 215 | 19-hydroxykovalic acid | H | CO ₂ H | α CH ₂ OH | H | H | α Me | CO ₂ H | Me | <i>Olearia teretifolia</i> | <i>Phytochemistry</i> , 1992, 31 , 1703-1711 |
| 216 | 18-oxo-19-seneciyl-oxy- <i>ent</i> -clerod-3- <i>en</i> -15-oic acid | H | CHO | α CH ₂ OSen | H | H | α Me | CO ₂ H | Me | | |
| 217 | 18-oxo- <i>ent</i> -clerod-3- <i>en</i> -15,19-dioic acid | H | CHO | α CO ₂ H | H | H | α Me | CO ₂ H | Me | | |
| 218 | 15,19-dihydroxy- <i>ent</i> -clerod-3- <i>en</i> -18-oic acid | H | CO ₂ H | α CH ₂ OH | H | H | α Me | CH ₂ OH | Me | | |
| 219 | 15,18,19-trihydroxy- <i>ent</i> -clerod-3- <i>ene</i> | H | CH ₂ OH | α CH ₂ OH | H | H | α Me | CH ₂ OH | Me | | |
| 220 | 18,19-dihydroxyclerod-3- <i>en</i> -15-oic acid | H | CH ₂ OH | α CH ₂ OH | H | H | α Me | CO ₂ H | β Me | <i>Chrysocoma comaurea</i> | <i>Phytochemistry</i> , 1991, 30 , 607-609 |
| 221 | 7-oxo-13,14-dihydrokolavenic acid | H | Me | α Me | H | =O | α Me | CO ₂ H | Me | <i>Pteronia camphorata</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |
| 222 | 6 α -hydroxy-7-oxo-13,14-dihydrokolavenic acid | H | Me | α Me | α OH | =O | α Me | CO ₂ H | Me | | |
| 223 | 6 α -angloyloxy-7-oxo-13,14-dihydrokolavenic acid | H | Me | α Me | α OAng | =O | α Me | CO ₂ H | Me | | |

| | | | | | | | | | | | |
|-----|---|--------------|--------------------------------|-------------|--------------------------------|--------------|-------------|--------------------|------------------|-----------------------------------|--|
| 224 | 6 α -isobutyryloxy-7-oxo-13,14-dihydrokolavenic acid | H | Me | α Me | α OiBu | =O | α Me | CO ₂ H | Me | | |
| 225 | <i>epi</i> -populifolic acid | H | Me | β Me | H | H | α Me | CO ₂ H | Me | <i>Aristolochia cymbifera</i> | <i>Phytochemistry</i> , 1992, 31 , 3277-3279 |
| 226 | 18-acetoxy- <i>cis</i> -cleroda-3-en-15-oic acid | H | CH ₂ OAc | β Me | H | H | β Me | CO ₂ H | Me | <i>Haplopappus uncinatus</i> | <i>J. Ethnopharmacol.</i> , 2006, 103 , 297-301 |
| 227 | (13 <i>S</i>)- <i>ent</i> -7 β -hydroxy-3-cleroden-15-oic acid | H | Me | α Me | H | α OH | α Me | CO ₂ H | β Me | | |
| 228 | <i>ent</i> -7 β -hydroxy-2-oxo-3-cleroden-15-oic acid | =O | Me | α Me | H | α OH | α Me | CO ₂ H | β Me | <i>Nuxia sphaerocephala</i> | <i>Phytochemistry</i> , 2006, 67 , 444-451 |
| 229 | <i>ent</i> -2,7-dioxo-3-clero-den-15-oic acid | =O | Me | α Me | H | =O | α Me | CO ₂ H | β Me | | |
| 230 | <i>ent</i> -18-(<i>E</i>)-caffeoyloxy-7 β -hydroxy-3-cleroden-15-oic acid | H | CH ₂ Z ₄ | α Me | H | α OH | α Me | CO ₂ H | β Me | | |
| 231 | 2 β -(formyloxy)clerod-3-en-15-oic acid | β OCHO | Me | α Me | H | H | α Me | CO ₂ H | Me | <i>Clausena dunniana</i> | <i>Helv. Chim. Acta.</i> , 2003, 86 , 3187-3193 |
| 232 | 2 β -(acetyloxy)clerod-3-en-15-oic acid | β OAc | Me | α Me | H | H | α Me | CO ₂ H | Me | | |
| 233 | ethyl clerod-3-en-15-oate | H | Me | α Me | H | H | α Me | CO ₂ Et | Me | | |
| 234 | premnone A | H | Me | α Me | α OCin (<i>trans</i>) | β OH | α Me | CO ₂ Me | =CH ₂ | <i>Premna tomentosa</i> | <i>Phytochemistry</i> , 2006, 67 , 1243-1248 |
| 235 | premnone B | H | Me | α Me | α OCin (<i>cis</i>) | β OH | α Me | CO ₂ Me | =CH ₂ | | |
| 236 | premnone C | H | Me | α Me | α OH | β OCin | α Me | CO ₂ Me | =CH ₂ | | |
| 237 | 7-oxo-kolavelool | H | Me | α Me | H | =O | α Me | α Me | α OH | <i>Ptychopetalum olacoides</i> | <i>Nat. Prod. Commun.</i> , 2011, 6 , 327-332 |
| 238 | 7 α -hydroxykolavelool | H | Me | α Me | H | α OH | α Me | α Me | α OH | | |
| 239 | (-)-13- <i>epi</i> -2-oxo-kolavelool | =O | Me | α Me | H | H | α Me | α Me | β OH | <i>Aristolochia chamissonis</i> | <i>Phytochemistry</i> , 1999, 50 , 455-461 |
| 240 | (-)-2 β -hydroxykolavelool | β OH | Me | α Me | H | H | α Me | α Me | β OH | | |
| 241 | (-)-2 β -hydroperoxykolavelool | β OOH | Me | α Me | H | H | α Me | α Me | β OH | | |
| 242 | 13- <i>epi</i> -roseostachenol | α OH | Me | α Me | H | H | α Me | α Me | β OH | | |
| 243 | roseostachenol | α OH | Me | α Me | H | H | α Me | α Me | α OH | <i>Stachys rosea</i> | <i>Phytochemistry</i> , 1994, 37 , 501-503 |
| 244 | roseostachenone | =O | Me | α Me | H | H | α Me | α Me | OH | | <i>Phytochemistry</i> , 1992, 31 , 3147-3149 |
| 245 | 13-hydroxy- <i>cis-ent</i> -cleroda-3,14-diene | H | Me | β Me | H | H | α Me | α Me | OH | <i>Adelanthus lindenbergianus</i> | <i>Phytochemistry</i> , 2004, 65 , 127-137 |
| 246 | (-)-2-oxo-13-hydroxy-3,14-clerodan-diene | =O | Me | α Me | H | H | α Me | α Me | β OH | <i>Guarea trichilioides</i> | <i>Phytochemistry</i> , 1996, 41 , 1159-1161 |
| 247 | <i>cis</i> -3,14-clerodadien-13-ol | H | Me | β Me | H | H | α Me | α Me | OH | <i>Jungermannia</i> | <i>J. Nat. Prod.</i> , 2001, 64 , 1309-1317 |

| | | | | | | | | | | | |
|-----|---|---------------------|---------------------|-----|---|---|-----|-----|-----|-----------------------------------|---|
| 248 | <i>ent</i> -kolavelool | H | Me | βMe | H | H | βMe | βMe | αOH | <i>infusca</i> | |
| 249 | 15,16-diacetoxy- <i>ent</i> -cleroda-3,12 <i>E</i> -diene | CH ₂ OAc | CH ₂ OAc | — | — | — | — | — | — | <i>Linaria saxatilis</i> | <i>Phytochemistry</i> , 1993, 33 , 631-633 |
| 250 | <i>ent</i> -cleroda-3,12 <i>E</i> -diene-15,16-dial | CHO | CHO | — | — | — | — | — | — | | |
| 251 | 15,16-diacetoxy- <i>ent</i> -cleroda-3,12 <i>Z</i> -diene | CH ₂ OAc | CH ₂ OAc | — | — | — | — | — | — | | |
| 252 | <i>ent</i> -cleroda-3,12 <i>Z</i> -diene-15,16-dial | CHO | CHO | — | — | — | — | — | — | | |
| 253 | palmadorin C | — | — | — | — | — | — | — | — | <i>Austrodoris kerguelenensis</i> | <i>J. Nat. Prod.</i> , 2010, 73 , 416–421 |

1.2.2. Type I Subtype IIb with a Double Bond at Another (or No) Position (Table 6)




| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | R ₇ | Source | Ref. |
|-----|--|----------------|----------------|----------------------------|--------------------|--------------------|----------------|----------------|-------------------------------|---|
| 254 | 3 α ,4 β ,13E-neoclerod-13-ene-3,4,15-triol | α OH | β OH | α Me | CH ₂ OH | Me | — | — | <i>Amora stellatosquamosa</i> | <i>Helv. Chim. Acta</i> , 2004, 87 , 1279-1286 |
| 255 | 3 α ,4 β ,13E-4-ethoxy-neoclerod-13-ene-3,15-diol | α OH | β OEt | α Me | CH ₂ OH | Me | — | — | | |
| 256 | crolechinol | H | H | β CH ₂ OH | CH ₂ OH | CH ₂ OH | — | — | <i>Croton lechleri</i> | <i>Phytochemistry</i> , 1993, 32 , 755-760 |
| 257 | 3R,4R-dihydroxyclerod-13E-en-15-al | α OH | β OH | α Me | CHO | Me | — | — | <i>Jungermannia hyalina</i> | <i>Phytochemistry</i> , 1995, 40 , 209-212 |

| | | | | | | | | | | |
|-----|--|-------------|--------------------|------------------|----------------------------|--------------------|--------------------|----|--|---|
| 258 | (-)-7 β -hydroxy cleroda-8(17),13E-dien-15-oic acid | — | — | — | — | — | — | — | <i>Eperua leucantha</i> | <i>J. Nat. Prod.</i> , 1992, 55 , 845-850 |
| 259 | 3R,4R-dihydroxyclerod-13Z-en-15-al | OH | OH | H | H | H | CHO | H | <i>Jungermannia hyalina</i> | <i>Phytochemistry</i> , 1995, 40 , 209-212 |
| 260 | sypirensin B | H | H | OH | OH | OH | CH ₂ OH | OH | <i>Teucrium chamaedrys</i> ssp. <i>sypirensis</i> | <i>J. Nat. Prod.</i> , 1996, 59 , 457-460 |
| 261 | 2 β -hydroxy-3-oxo- <i>cis</i> -clerodan-15-oic acid | β OH | =O | H | β Me | β Me | H | — | <i>Chrysocoma comaaurea</i> | <i>Phytochemistry</i> , 1991, 30 , 607-609 |
| 262 | 6 α -angeloyloxy-4 α -hydroxy-3,4,13,14-tetrahydrokolavenic acid | H | H | α OH | β Me | α Me | α OTig | — | <i>Pteronia camphorata</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |
| 263 | 4 α ,18-dihydroxyclerodan-15-oic acid | H | H | α OH | β CH ₂ OH | α Me | H | — | <i>Clausena dunniana</i> | <i>Helv. Chim. Acta.</i> , 2003, 86 , 3187-3193 |
| 264 | 4 β -hydroxyclerodan-15-oic acid | H | H | β OH | α Me | α Me | H | — | | |
| 265 | 3 α ,4 α -dihydroxy clerodan-15-oic acid | H | α OH | α OH | β Me | α Me | H | — | | |
| 266 | 3,4-dihydroxyclerodan-13E-en-15-oic acid | — | — | — | — | — | — | — | <i>Detarium microcarpum</i> | <i>J. Nat. Prod.</i> , 2006, 69 , 768-773 |
| 267 | 3,4-dihydroxyclerodan-13Z-en-15-oic acid | — | — | — | — | — | — | — | | |
| 268 | roseotetrol | α OH | α OH | β OH | — | — | — | — | <i>Stachys rosea</i> | <i>Phytochemistry</i> , 1994, 37 , 501-503 |
| 269 | roseostachone | H | =O | H | — | — | — | — | | <i>Phytochemistry</i> , 1992, 31 , 3147-3149 |
| 270 | 13-hydroxycleroda-7,14-diene | — | — | — | — | — | — | — | <i>Carex distachya</i> | <i>Nat. Prod. Commun.</i> , 2010, 5 , 1539-1542 |
| 271 | 15-hydroxycleroda-7,13-diene | — | — | — | — | — | — | — | | |
| 272 | (-)-cleroda-7,13E-dien-15-oic acid | — | — | — | — | — | — | — | <i>Eperua purpurea</i> | <i>J. Nat. Prod.</i> , 1993, 56 , 1586-1589 |
| 273 | — | — | — | — | — | — | — | — | <i>Schistochila aligera</i> | <i>Phytochemistry</i> , 1991, 30 , 849-851 |
| 274 | laetiaprocerine D | — | — | — | — | — | — | — | <i>Laetia procera</i> | <i>Bioorg. Med. Chem. Lett.</i> , 2005, 15 , 5065-5070 |
| 275 | (3 α ,4 β -14RS)-neo-clerod-13(16)-ene-3,4,14,15-tetrol | — | — | — | — | — | — | — | <i>Amoora stellatosquamosa</i> | <i>Helv. Chim. Acta.</i> , 2004, 87 , 1279-1286 |
| 276 | leojaponin A | — | — | — | — | — | — | — | <i>Leonurus japonicus</i> | <i>Chin. Chem. Lett.</i> , 2014, 25 , 677-679 |
| 277 | tinosporaclerodanol | — | — | — | — | — | — | — | <i>Tinospora cordifolia</i> | <i>Nat. Prod. Res.</i> , 2010, 24 , 926-934 |
| 278 | pentandranoic acid C | α OH | CO ₂ H | =CH ₂ | =O | — | — | — | <i>Callicarpa pentandra</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 1062-1065 |
| 279 | 3 β -hydroxy-clerod-4(18)-en-15-oic acid | β OH | CO ₂ H | Me | H | — | — | — | <i>Clausena dunniana</i> | <i>Helv. Chim. Acta.</i> , 2003, 86 , 3187-319 |
| 280 | ethyl-clerod-4(18)-en-15-oate | H | CO ₂ Et | Me | H | — | — | — | | |
| 281 | 6 α ,16-dihydroxy cleroda-4(18),13-dien-15-oic acid | — | — | — | — | — | — | — | | |
| 282 | — | α OH | Me | α OAc | CO ₂ Me | Me | — | — | <i>Solidago altissima</i> | <i>Phytochemistry</i> , 1999, 52 , 487-493 |
| 283 | — | β OH | Me | α OAc | CO ₂ Me | Me | — | — | | |
| 284 | — | =O | Me | H | CO ₂ Me | Me | — | — | | |
| 285 | cleroda-4(18),13(14)E-dien-15-oic acid | H | Me | H | CO ₂ H | Me | — | — | <i>Polyalthia cheliensis</i> | <i>Phytochemistry</i> , 1995, 39 , 447-448 |
| 286 | 3-hydroxy-cleroda-4(18),13Z-dien-15-oic acid | β OH | Me | H | CO ₂ H | Me | — | — | <i>Cyathocalyx zeylanica</i> | <i>Phytochemistry</i> , 1995, 39 , 443-445 |
| 287 | porwenin A | H | CO ₂ H | H | CH ₂ OH | CH ₂ OH | — | — | <i>Portulaca okinawensis</i> | <i>J. Nat. Prod.</i> , 2001, 64 , 804-805 |

| | | | | | | | | | | |
|-----|--------------|-----|---|---|---|---|---|---|---------------------------------------|--|
| 288 | palmadorin A | OH | — | — | — | — | — | — | <i>Austrodoris kerguelenensis</i> | <i>J. Nat. Prod.</i> , 2010, 73 , 416–421 |
| 289 | palmadorin B | OAc | — | — | — | — | — | — | | |

1.3. Type I Subtype III with an Epoxy Ring (Table 7)




| No. | Compound Name | Source | Ref. |
|-----|---|------------------------------|---|
| 290 | <i>ent</i> -3 β ,4 β -epoxyclerod-13 <i>E</i> -en-15-ol | <i>Jungermannia hyalina</i> | <i>Phytochemistry</i> , 1995, 40 , 209-212 |
| 291 | 3,4-epoxyclerodan-13 <i>E</i> -en-15-oic acid | <i>Detarium microcarpum</i> | <i>J. Nat. Prod.</i> , 2006, 69 , 768-773 |
| 292 | heteroscyphic acid C | <i>Heteroscyphus planus</i> | <i>Phytochemistry</i> , 1994, 37 , 1263-1268 |
| 293 | <i>ent</i> -3 β ,4 β -epoxy-clerod-14-en-13 ζ -ol | <i>Jungermannia paroica</i> | <i>Phytochemistry</i> , 1992, 31 , 1420-1421 |
| 294 | 3 α ,4 α -epoxyroseostachenol | <i>Stachys glutinosa</i> | <i>J. Nat. Prod.</i> , 2015, 78 , 69-76 |
| 295 | 4 α ,18 β -epoxy-16-hydroxyclerod-13-en-15-oic acid | <i>Polyalthia longifolia</i> | <i>J. Nat. Prod.</i> , 2009, 72 , 1960-1963 |
| 296 | ajugacumbin J | <i>Ajuga decumbens</i> | <i>Nat. Prod. Res.</i> , 2014, 28 , 196-200 |

2. Type II with a 2-Ethylfuran-based Side Chain at C-9

2.1. Type II with Various *O*-Containing Rings

2.1.1. Type II Subtype Ia with Various *O*-Containing Rings (Table 8)




| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | Source | Ref. |
|-----|---|----------------|-------------------|----------------|---------------------|----------------|---------------------------------|---|
| 297 | nasimalun A | H | H | H | αCO ₂ Me | H | <i>Barringtonia racemosa</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 410-411 |
| 298 | 2α-hydroxy-7α-acetoxy-12-oxo-15:16-epoxy-neoclerodan-3,13(16),14-trien-18:19-olide | H | OH | αOAc | αMe | =O | <i>Salvia urolepis</i> | <i>Phytochemistry</i> , 1995, 38 , 171-174 |
| 299 | — | H | H | βOAc | αMe | OH | <i>Salvia miniata</i> | <i>Phytochemistry</i> , 2011, 72 , 265-275 |
| 300 | ent-(5R,9S,10S)-7S-acetoxy-15,16-epoxy-1S,2S,12ζ-trihydroxycleroda-3,13(16),14-trien-18,19-olide | αOH | βOH | αOAc | βMe | OH | <i>Salvia haenke</i> | <i>Tetrahedron</i> , 1997, 53 , 14719-14728 |
| 301 | raspailol | — | — | — | — | — | <i>Raspailia</i> species | <i>Aust. J. Chem.</i> , 1998, 51 , 1097-1101 |
| 302 | teumassilenin B | — | — | — | — | — | <i>Teucrium massiliense</i> | <i>J. Nat. Prod.</i> , 1998, 61 , 1242-1247 |
| 303 | gaudichaudone | — | — | — | — | — | <i>Baccharis gaudichaudiana</i> | <i>J. Nat. Prod.</i> , 1994, 57 , 801-807 |
| 304 | dugesin G | — | — | — | — | — | <i>Salvia dugesii</i> | <i>Nat. Prod. Bioprospect.</i> , 2011, 1 , 81-86 |
| 305 | teuctomin | — | — | — | — | — | <i>Teucrium tomentosum</i> | <i>Nat. Prod. Res.</i> , 2010, 24 , 7-12 |
| 306 | plaunol E | — | — | — | — | — | <i>Croton stellatopilosus</i> | <i>J. Nat. Med.</i> , 2013, 67 , 174-181 |
| 307 | 12-O-acetylteugnaphalodin | — | — | — | — | — | <i>Teucrium oxylepis</i> | <i>Phytochemistry</i> , 1991, 30 , 4079-4082 |
| 308 | (2aβ,3α,5aβ,6β,7R,8αα)-6-[2-(3-furanyl)ethyl]-2a,3,4,5,5a,6,7,8,8a,8b-decahydro-2a,3-dihydroxy-6,7,8b-trimethyl-2H-naphtho[1,8-bc]furan-2-one | — | — | — | — | — | <i>Tinospora rumphii</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 509-511 |
| 309 | dihydrolinguifolide | — | — | — | — | — | <i>Demotarisia linguifolia</i> | <i>Phytochemistry</i> , 1990, 29 , 3229-3231 |
| 310 | linguifolide | H | H | H | =O | — | | |
| 311 | 1β-acetoxy-12-hydroxy-15,16-epoxy-cis-cleroda-3,13(16),14-triene-18,16-olide | βOAc | H | H | OH | — | <i>Jamesoniella autumnalis</i> | <i>Phytochemistry</i> , 1995, 39 , 859-868 |
| 312 | 1β-acetoxy-7,12-dihydroxy-15,16-epoxy-cis-cleroda-3,13(16),14-triene-18,6-olide | βOAc | βOH | H | OH | — | | <i>Phytochemistry</i> , 1998, 48 , 681-685 |
| 313 | 17-acetoxy-1β,12-dihydroxy-15,16-epoxy-cis-ent-cleroda-3,13(16),14-triene-6α,18-olide* | βOH | H | OAc | OH | — | | <i>J. Nat. Prod.</i> , 1992, 55 , 111-121 |
| 314 | tepolin A | OH | CO ₂ H | OH | — | — | <i>Teucrium polium</i> | <i>Him. Prir. Soedin.</i> , 1992, 5 , 503-508 |
| 315 | tepolin B | H | CO ₂ H | OH | — | — | | |
| 316 | difuranofruticol | — | — | — | — | — | <i>Teucrium fruticans</i> | <i>Phytochemistry</i> , 2005, 66 , 2298-2303 |
| 317 | syrapolin II | — | — | — | — | — | <i>Teucrium polium</i> | <i>Jordan J. Chem.</i> , 2011, 6 , 339-345 |
| 318 | sacacarin | — | — | — | — | — | <i>Croton cajucara</i> | <i>Phytochemistry</i> , 1998, 49 , 823-828 |

| | | | | | | | | |
|-----|--|------------|-------------|----|-------------|------------|--------------------------|---|
| 319 | — | — | — | — | — | — | <i>Salvia miniata</i> | <i>Phytochemistry</i> , 2011, 72 , 265-275 |
| 320 | 6 β - <i>O</i> -acetyl-3 β -hydroxyteucroxylepin | β OH | β OAc | H | =O | OH | <i>Teucrium yemense</i> | <i>Phytochemistry</i> , 1995, 40 , 1737-1741 |
| 321 | teucroxylepin | H | β OH | H | =O | β OH | <i>Teucrium oxylepis</i> | <i>Phytochemistry</i> , 1991, 30 , 4079-4082 |
| 322 | montanin H | H | α OH | =O | α OH | OAc | <i>Teucrium montanum</i> | <i>Phytochemistry</i> , 1992, 31 , 4029-4030 |
| 323 | pteroneeniol | — | — | — | — | — | <i>Pteronia eenii</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |

*Compound's name (6 α ,18-olide) and structure (18,6 α -olide) are shown as given in reference 121.


2.1.2. Type II Subtype Ib with Other *O*-Containing Rings (Table 9)



| No. | Compound Name | R ₁ | R ₂ | Source | Ref. |
|-----|--|----------------|----------------|--------------------------------|---|
| 324 | 3 α ,12-dihydroxy-4 α ,20,15,16-bisepoxy-8 β ,10 β H-ent-cleroda-13(16),14-diene | — | — | <i>Microglossa pyrropappa</i> | Phytochemistry, 1990, 29 , 3233-3241 |
| 325 | 12,20-dihydroxy-3 α ,4 α ,15,16-bisepoxy-8 β ,10 β H-ent-cleroda-13(16),14-diene | — | — | | |
| 326 | teucosin B | — | — | <i>Teucrium cossonii</i> | Phytochemistry, 1992, 31 , 3957-3960 |
| 327 | ptychonolide | =O | — | <i>Ptychopetalum olacoides</i> | J. Nat. Prod., 2008, 71 , 1760-1763 |
| 328 | 20- <i>O</i> -methylptychonolide | β OMe | — | | |
| 329 | ptychonolide hemiacetal | β OH | — | | |
| 330 | methyl dodonate A | H | — | <i>Dodonaea viscosa</i> | Tetrahedron, 2001, 57 , 2981-2989 |

| | | | | | |
|-----|--|--------------|----|-----------------------------------|---|
| 331 | methyl dodonate B | OH | — | | |
| 332 | methyl dodonate C | — | — | | |
| 333 | furocrotinsulolide B | — | — | <i>Croton insularis</i> | <i>Helv. Chim. Acta.</i> , 2005, 88 , 2654-2660 |
| 334 | teucrolin A | — | — | <i>Teucrium oliverianum</i> | <i>J. Nat. Prod.</i> , 1993, 56 , 830-842 |
| 335 | teucrolin E | — | — | | <i>Phytochemistry</i> , 2002, 59 , 409-414 |
| 336 | teumassilenin C | — | — | <i>Teucrium massiliense</i> | <i>J. Nat. Prod.</i> , 1998, 61 , 1242-1247 |
| 337 | anastreptin | — | — | <i>Adelanthus lindenbergianus</i> | <i>Phytochemistry</i> , 2004, 65 , 127-137 |
| 338 | 8 β ,12:15,16-diepoxy- <i>cis-ent</i> -cleroda-13(16),14-dien-18 α ,6 α -olide | — | — | | |
| 339 | bafoudiosbulbin A | — | — | <i>Dioscorea bulbifera</i> | <i>Phytochemistry</i> , 2006, 67 , 1957-1963 |
| 340 | bafoudiosbulbin D | — | — | | <i>Helv. Chim. Acta</i> , 2007, 90 , 1599-1605 |
| 341 | bafoudiosbulbin E | — | — | | |
| 342 | Apiciflorin | — | — | <i>Cleidion spiciflorum</i> | <i>Phytochemistry</i> , 2006, 67 , 1029-1033 |
| 343 | scaparvin A | — | — | <i>Scapania parva</i> | <i>Org. Lett.</i> , 2010, 12 , 4404-4407 |
| 344 | scaparvin B | =O | — | | |
| 345 | scaparvin C | α OH | — | | |
| 346 | scaparvin D | α OAc | — | | |
| 347 | parvitexin A | H | — | <i>Scapania parvitexta</i> | <i>Biosci. Biotechnol. Biochem.</i> , 2007, 71 , 2751-2758 |
| 348 | parvitexin B | β OAc | — | | |
| 349 | parvitexin C | β OH | — | | |
| 350 | scaparvin E | OAc | — | <i>Scapania parva</i> | <i>Org. Lett.</i> , 2010, 12 , 4404-4407 |
| 351 | stephanialide A | =O | OH | <i>Scapania stephaniai</i> | <i>Phytochemistry</i> , 2014, 105 , 85-91 |
| 352 | stephanialide B | β OH | OH | | |
| 353 | stephanialide C | β OAc | OH | | |
| 354 | stephanialide D | =O | OH | | |
| 355 | stephanialide E | β OAc | OH | | |

2.1.3. Type II Subtype Ic with a Simple Epoxy Ring (Table 10)




| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | R ₇ | Source | Ref. |
|-----|--|----------------|----------------|--------------------|---------------------|---------------------|----------------|----------------|------------------------------------|---|
| 356 | parvitexin D | OAc | — | — | — | — | — | — | <i>Scapania parvixesta</i> | <i>Biosci. Biotechnol. Biochem.</i> , 2007, 71 , 2751-2758 |
| 357 | parvitexin E | OH | — | — | — | — | — | — | | |
| 358 | epoxychiromodine | H | H | CO ₂ Me | Me | =O | — | — | <i>Croton megalocarpus</i> | <i>Phytochemistry</i> , 1992, 31 , 2055-2058 |
| 359 | eluterin C | H | αOAc | Me | CH ₂ OH | H | — | — | <i>Croton eluteria</i> | <i>J. Agric. Food Chem.</i> , 2002, 50 , 5131-5138 |
| 360 | eluterin D | H | αOH | Me | CH ₂ OAc | H | — | — | | |
| 361 | eluterin E | H | αOAc | Me | CH ₂ OH | βOH | — | — | | |
| 362 | eluterin F | αOAc | αOAc | Me | CH ₂ OAc | H | — | — | | |
| 363 | eluterin G | αOAc | αOAc | Me | CHO | H | — | — | | |
| 364 | 3β,4β:15,16-diepoxy-13(16),14-clerodadiene | H | — | — | — | — | — | — | <i>Thysananthus spathulistipus</i> | <i>Chem. Pharm. Bull.</i> , 2006, 54 , 1046-1049 |
| 365 | thysaspathone | =O | — | — | — | — | — | — | | |
| 366 | teucossin A | H | OAc | OH | H | CH ₂ OAc | H | OAc | <i>Teucrium cossonii</i> | <i>Phytochemistry</i> , 1992, 31 , 3957-3960 |
| 367 | teugracilin E | βOAc | OAc | OAc | H | CH ₂ OAc | H | OAc | <i>Teucrium gracile</i> | <i>Phytochemistry</i> , 1992, 31 , 3531-3534 |
| 368 | teugracilin C | βOH | OAc | OH | H | Me | H | OAc | | <i>Phytochemistry</i> , 1991, 30 , 3693-3697 |
| 369 | teucrolivin D | βOAc | OAc | OH | βOAc | Me | H | H | <i>Teucrium oliverianum</i> | <i>Phytochemistry</i> , 1991, 30 , 1603-1606 |
| 370 | teucrolivin E | βOAc | OAc | OH | =O | Me | H | H | | |
| 371 | 7β-hydroxyfruticolone | H | OAc | =O | βOH | Me | H | H | <i>Teucrium fruticans</i> | <i>Phytochemistry</i> , 2004, 65 , 387-392 |
| 372 | 11-hydroxyfruticolone | H | OAc | =O | H | Me | OH | H | | |
| 373 | deacetylfruticolone | H | OH | =O | H | Me | H | H | | |
| 374 | deoxyfruticolone | H | OAc | =O | H | Me | H | H | | |

| | | | | | | | | | | |
|-----|----------------------|---|---|---|---|---|---|---|--------------------------|--|
| 375 | didehydrofruticolone | — | — | — | — | — | — | — | | |
| 376 | phlomeoic acid | — | — | — | — | — | — | — | <i>Phlomis bracteosa</i> | <i>Nat. Prod. Commun.</i> , 2011, 6 , 171-173 |

2.2. Type II with or without a C=C Double Bond in the Decalin Moiety

2.2.1. Type II Subtype IIa with One or More Decalin C=C Double Bonds (Table 11)




| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | Source | Ref. |
|-----|-------------------------|--------------------|----------------|----------------|----------------|--------------------|---------------------|--------------------------------|---|
| 377 | vishautriwaic acid | — | — | — | — | — | — | <i>Dodonaea viscosa</i> | <i>Z. Naturforsch. B</i> , 2010, 65 , 83-86 |
| 378 | hautriwaic acid | — | — | — | — | — | — | <i>Eremocarpus setigerus</i> | <i>Indian J. Chem. B</i> , 1991, 30 , 1054-1055 |
| 379 | nasimalun B | CO ₂ Me | Me | H | H | CO ₂ Me | Me | <i>Barringtonia racemosa</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 410-411 |
| 380 | ptychonal | Me | Me | H | αOH | Me | CHO | <i>Ptychopetalum olacoides</i> | <i>J. Nat. Prod.</i> , 2008, 71 , 1760-1763 |
| 381 | eluterin K | Me | Me | αOAc | αOAc | Me | CH ₂ OAc | <i>Croton eluteria</i> | <i>J. Agric. Food Chem.</i> , 2003, 51 , 6970-6974 |
| 382 | 6α,7α-dihydroxyannonene | Me | Me | αOH | αOH | Me | Me | <i>Ptychopetalum</i> | <i>Bioorg. Med. Chem. Lett.</i> , 2009, 19 , |


| | | | | | | | | | |
|-----|--|--------------------|-------------------|------------------------------|--------------|---------------------|--------------------|---------------------------------|---|
| 383 | 7 α ,20-dihydroxyannonene | Me | Me | H | α OH | Me | CH ₂ OH | <i>olacoides</i> | 882–886 |
| 384 | 7 α -acetoxybacchotricuneatin D | CH ₂ OH | Me | H | α OAc | Me | Me | <i>Copaifera sp.</i> | <i>Phytochemistry</i> , 1996, 42 , 1653-1656 |
| 385 | 6 α -hydroxyannonene | Me | Me | α OH | H | Me | Me | <i>Croton sonderianus</i> | <i>Phytochemistry</i> , 1994, 36 , 1457-1463 |
| 386 | 6 α ,7 β -dihydroxyannonene | Me | Me | α OH | β OH | Me | Me | | |
| 387 | 6 α ,7 β -diacetoxyannonene | Me | Me | α OAc | β OAc | Me | Me | | |
| 388 | 7 α -hydroxyhardwickiic acid | CO ₂ H | Me | H | α OH | Me | Me | <i>Conyza hypoleuca</i> | <i>Phytochemistry</i> 1991, 30 , 575-581 |
| 389 | 15,16-epoxy-8R-(benzoyloxy)methyl-cleroda-3,13(16),14-trien-18-oic acid | CO ₂ H | Me | H | H | CH ₂ OBz | Me | <i>Dodonaea polyandra</i> | <i>J. Nat. Prod.</i> , 2011, 74 , 650-657 |
| 390 | 15,16-epoxy-8R-(benzoyloxy)methyl-2R-hydroxycleroda-3,13(16),14-trien-18-oic acid | α OH | CO ₂ H | Me | H | CH ₂ OBz | Me | | |
| 391 | 15,16-epoxy-2R-benzoyloxycleroda-3,13(16),14-trien-18-oic acid | α OBz | CO ₂ H | Me | H | Me | Me | | |
| 392 | 15,16-epoxy-8R-(benzoyloxy)methyl-2-oxocleroda-3,13(16),14-trien-18-oic acid | =O | CO ₂ H | Me | H | CH ₂ OBz | Me | | |
| 393 | 15,16-epoxy-3,13(16)-clerodatriene-2-one | =O | Me | α Me | H | Me | Me | <i>Croton ururucana</i> | <i>Phytochemistry</i> , 1998, 49 , 171-174 |
| 394 | cajucarín A | =O | Me | CHO | H | Me | CO ₂ Me | <i>Croton cajucara</i> | <i>Chem. Pharm. Bull.</i> , 1990, 38 , 701-705 |
| 395 | chromiargyne | =O | Me | Me | H | Me | CO ₂ Me | <i>Croton hemiargyreus</i> | <i>Nat. Prod. Lett.</i> , 1998, 12 , 41-46 |
| 396 | 7-acetoxycromiargyne | =O | Me | Me | OAc | Me | CO ₂ Me | | |
| 397 | raspailenone | — | — | — | — | — | — | <i>Raspailia</i> species | <i>Aust. J. Chem.</i> , 1998, 51 , 1097-1101 |
| 398 | laevigatbenzoate | β OBz | CO ₂ H | β Me | H | H | H | <i>Croton laevigatus</i> | <i>J. Nat. Med.</i> , 2011, 65 , 391–394 |
| 399 | 6 β -hydroxy-15,16-epoxy-5 β ,8 β ,9 β ,10 α -cleroda-3,13(16),14-trien-18-oic acid | H | CO ₂ H | β Me | β OH | H | H | <i>Duranta repens</i> | <i>Helv. Chim. Acta</i> , 2001, 84 , 649-655 |
| 400 | 2 β -hydroxy-15,16-epoxy-5 β ,8 β ,9 β ,10 α -cleroda-3,13(16),14-trien-18-oic acid | β OH | CO ₂ H | β Me | H | H | H | | |
| 401 | (5R,8S,9R,10S)-15,16-epoxy- <i>cis</i> -cleroda-3,13(16),14-trien-18-al | H | CHO | α Me | H | H | H | <i>Gottschelia schizopleura</i> | <i>Planta Med.</i> , 2009, 75 , 1597-1601 |
| 402 | (+)-7 β -acetoxy-15,16-epoxy-3,13(16),14-clerodatrien-18-oic acid | H | CO ₂ H | β Me | H | α OAc | H | <i>Sindora sumatrana</i> | <i>Chem. Pharm. Bull.</i> , 1994, 42 , 1202-1207 |
| 403 | scaparvin F | H | Me | α CH ₂ OAc | β OH | H | β OAc | <i>Scapania</i> | <i>Phytochem. Lett.</i> , 2012, 5 , 535–540 |

| | | | | | | | | | |
|-----|--|-------------|--------------------|---------------------------|---------------------|----|---|----------------------------------|--|
| | | | | | | | | <i>parva</i> | |
| 404 | crotonolide G | — | — | — | — | — | — | <i>Croton laui</i> | <i>J. Nat. Prod.</i> , 2014, 77 , 1013-1020 |
| 405 | divinatorin A | α OH | CO ₂ H | α Me | Me | H | — | <i>Salvia divinorum</i> | <i>J. Nat. Prod.</i> , 2003, 66 , 1242-1244 |
| 406 | divinatorin B | α OH | CO ₂ Me | α Me | CH ₂ OH | H | — | | |
| 407 | divinatorin C | H | CO ₂ H | α Me | CH ₂ OAc | H | — | | |
| 408 | (-)-hardwickiic acid | H | CO ₂ H | α Me | Me | H | — | | |
| 409 | divinatorin D | α OH | CO ₂ Me | α Me | CH ₂ OAc | H | — | <i>Salvia divinorum</i> | <i>Bioorg. Med. Chem.</i> , 2005, 13 , 5635-5639 |
| 410 | divinatorin E | α OH | CO ₂ Me | α Me | CHO | H | — | <i>Conyza hypoleuca</i> | <i>Phytochemistry</i> 1991, 30 , 575-581 |
| 411 | 12-hydroxyhardwickiic acid | H | CO ₂ H | α Me | Me | OH | — | | |
| 412 | 12,17-dihydroxyhardwickiic acid methyl ester | H | CO ₂ Me | α Me | CH ₂ OH | OH | — | <i>Chrysocoma comaurea</i> | <i>Phytochemistry</i> , 1991, 30 , 607-609 |
| 413 | 15,16-epoxy- <i>cis</i> -cleroda-3,13(16),14-trien-19-oic acid | H | Me | β CO ₂ H | Me | H | — | | |
| 414 | crotomembranafuran | H | CO ₂ Me | α Me | CO ₂ Me | =O | — | <i>Croton membranaceus</i> | <i>Nat. Prod. Commun.</i> , 2008, 3 , 1875-1878 |
| 415 | 1,2,11,12E-tetradehydrohardwickiic acid | — | — | — | — | — | — | <i>Conyza hypoleuca</i> | <i>Phytochemistry</i> 1991, 30 , 575-581 |
| 416 | 2 α -hydroxy-11,12E-dehydrohardwickiic acid | α OH | CO ₂ H | — | — | — | — | | |
| 417 | dehydrohardwickiic acid | H | CO ₂ H | — | — | — | — | <i>Eremocarpus setigerus</i> | <i>Indian J. Chem. B</i> , 1991, 30 , 1054-1055 |
| 418 | crotonoligaketone | — | — | — | — | — | — | <i>Croton oligandrum</i> | <i>Z. Naturforsch. C</i> , 2014, 69 , 181-185 |
| 419 | eluterin B | — | — | — | — | — | — | <i>Croton eluteria</i> | <i>J. Agric. Food Chem.</i> , 2002, 50 , 5131-5138 |

2.2.2. Type II Subtype IIb without a Decalin C=C Double Bond (Table 12)




420-427



428

| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | R ₇ | R ₈ | R ₉ | Source | Ref. |
|-----|---|----------------|----------------|----------------|---------------------------|--------------------|----------------|----------------|---------------------|----------------|-----------------------------|---|
| 420 | crotonolide H | H | α OH | β OH | Me | Me | H | H | Me | β OH | <i>Croton laui</i> | <i>J. Nat. Prod.</i> , 2014, 77 , 1013-1020 |
| 421 | 12-deoxycrotonolide H | H | α OH | β OH | Me | Me | H | H | Me | H | | |
| 422 | crolechinic acid | H | H | H | β CO ₂ H | Me | H | H | Me | H | <i>Croton lechleri</i> | <i>Phytochemistry</i> , 1993, 32 , 755-760 |
| 423 | teumassilenin A | H | H | H | β CHO | CH ₂ OH | α OH | H | Me | β OH | <i>Teucrium massiliense</i> | <i>J. Nat. Prod.</i> , 1998, 61 , 1242-1247 |
| 424 | caszarilladione | =O | H | H | α Me | Me | H | H | Me | =O | <i>Croton eluteria</i> | <i>J. Agric. Food Chem.</i> , 2003, 51 , 6970-6974 |
| 425 | eluterin A | H | =O | H | β Me | Me | H | α OAc | CH ₂ OAc | H | | <i>J. Agric. Food Chem.</i> , 2002, 50 , 5131-5138 |
| 426 | 3,12-dioxo-15,16-epoxy-cleroda-13(16),14-dien-9-al | H | =O | H | β Me | Me | H | H | CHO | =O | <i>Croton hovarum</i> | <i>Phytochemistry</i> , 1997, 45 , 379-381 |
| 427 | 3 α ,4 β -dihydroxy-15,16-epoxy-12-oxo-cleroda-13(16),14-dien-9-al | H | α OH | β OH | α Me | Me | H | H | CHO | =O | | <i>Phytochemistry</i> , 1996, 41 , 561-563 |
| 428 | nepetanal | — | — | — | — | — | — | — | — | — | <i>Nepeta juncea</i> | <i>Magn. Reson. Chem.</i> , 2009, 47 , 625-627 |

2.3. Type II Subtype III with a Tetrahydrofuran Ring (Table 13)





| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | R ₇ | Source | Ref. |
|-----|--|--------------------------|----------------|----------------|----------------|----------------|----------------|----------------|---------------------------------|---|
| 429 | trinerolide | αOMe | — | — | — | — | — | — | <i>Baccharis trinervis</i> | <i>Phytochemistry</i> , 1993, 34 , 1377-1384 |
| 430 | 15-epitrinerolide | βOMe | — | — | — | — | — | — | | |
| 431 | 18-acetyl-7α-hydroxy-epimethyl-trineracetal | CH ₂ OAc | H | αOH | H | H | βOMe | H | | |
| 432 | 18-methylmalonyl-7α-hydrox-methyl-trineracetal | CH ₂ OMe-malo | H | αOH | H | H | αOMe | H | | |
| 433 | 18-methylmalonyl-7α-hydroxy-epimethyl-trineracetal | CH ₂ OMe-malo | H | αOH | H | H | βOMe | H | | |
| 434 | 15,16-epoxy-7α,18-dihydroxy-15-methoxy- <i>ent</i> -clerod-3-ene | CH ₂ OH | H | αOH | H | H | OMe | H | <i>Baccharis articulata</i> | <i>Phytochemistry</i> , 1993, 34 , 1087-1090 |
| 435 | 15,16-epoxy-15α-methoxy- <i>ent</i> -clerod-3-en-18-oic acid | COOH | H | H | H | H | αOMe | H | <i>Baccharis gaudichaudiana</i> | <i>J. Nat. Prod.</i> , 2006, 69 , 274-276 |
| 436 | 13- <i>epi</i> -15,16-epoxy-15α-methoxy- <i>ent</i> -clerod-3-en-18-oic acid | COOH | H | H | H | H | βOMe | H | | |
| 437 | visclerodol acid | CO ₂ H | H | H | OH | βOH | βOAc | βOAc | <i>Dodonaea viscosa</i> | <i>Z. Naturforsch. B</i> , 2010, 65 , 83-86 <i>Phytochem. Lett.</i> , 2014, 8 , 10-15 |
| 438 | 13,14-dihydroxy-15,16-dimethoxy-(-)-6α-hydroxy-5α,8α,9α,10α-cleroda-3-en-18-oic acid | CO ₂ H | αOH | H | OH | βOH | βOMe | βOMe | <i>Scutellaria galericulata</i> | <i>Phytochemistry</i> , 1993, 33 , 309-315, <i>Phytochemistry</i> , 1996, 41 , 247-253 |
| 439 | scutegalin B | OH | βOH | — | — | — | — | — | | |
| 440 | scutegalin C | OTig | OH | — | — | — | — | — | | |
| 441 | 14,15,16-triacetoxy-15,16-epoxy- <i>ent</i> -cleroda-4(18),12 <i>E</i> -diene | — | — | — | — | — | — | — | | |
| 442 | 15,16-diacetoxy-12,13-15,16-diepoxy- <i>ent</i> -cleroda-3-ene | — | — | — | — | — | — | — | | |
| 443 | 14,15,16-triacetoxy-15,16-epoxy- <i>ent</i> -cleroda-3,12 <i>Z</i> -diene | βOAc | αOAc | βOAc | — | — | — | — | <i>Linaria saxatilis</i> | <i>Phytochemistry</i> , 1993, 33 , 631-633, <i>Phytochemistry</i> , 1995, 40 , 1307-1309 |
| 444 | 15,16-diacetoxy-15,16-epoxy- <i>neo</i> -cleroda-3,13 <i>Z</i> -diene | OAc | OAc | H | — | — | — | — | | |
| 445 | 15,16-diacetoxy-12,13-15,16-diepoxy-14-hydroxy- <i>neo</i> -cleroda-3-ene | — | — | — | — | — | — | — | | |

3. Type III with a 3-Ethyl-2-butenolide-based Side Chain at C-9

3.1. Type III Subtype I with *O*-Containing Rings

3.1.1. Type III Subtype Ia with Five-Membered Cyclic *O*-Containing Rings (Table 14)






| No. | Compound Name | R ₁ | R ₂ | R ₃ | Source | Ref. |
|-----|--|----------------|----------------|----------------|------------------------------------|---|
| 446 | amphiacrolide A | H | H | =O | <i>Amphiachyris dracunculoides</i> | <i>J. Nat. Prod.</i> , 1990, 53 , 1312-1326, <i>J. Nat. Prod.</i> , 1996, 59 , 463-468, <i>J. Nat. Prod.</i> , 1996, 59 , 5-14 |
| 447 | amphiacrolide B | H | =O | H | | |
| 448 | amphiacrolide C | H | H | OH | | |
| 449 | amphiacrolide L | βOH | =O | H | | |
| 450 | amphiacrolide J | — | — | — | | |
| 451 | amphiacrolide D | H | H | OH | | |
| 452 | amphiacrolide M | βOH | H | βOH | | |
| 453 | amphiacrolide E | H | αOEt | αOH | | |
| 454 | amphiacrolide I | H | αOMe | αOH | | |
| 455 | cleroda-3,13(14)-dien-16,15:18,19-diolide | — | — | — | <i>Solidago virgaurea</i> | <i>Phytochemistry</i> , 2010, 71 , 104–109 |
| 456 | Mkapwanin | H | H | H | <i>Dodonaea angustifolia</i> | <i>Phytochem. Lett.</i> , 2010, 3 , 217–220 |
| 457 | 15-methoxymkapwanin | H | H | OMe | | |
| 458 | 1α,7α-dihydroxyneocleroda-3,13-dien-16,15:18,19-diolide | αOH | αOH | H | <i>Baccharis crispa</i> | <i>J. Nat. Prod.</i> , 1997, 60 , 490-492 |
| 459 | 1α,7α,15-trihydroxyneocleroda-3,13-dien-16,15:18,19-diolide | αOH | αOH | OH | | |
| 460 | 8β-hydroxy-7-oxo-ent-cleroda-3-en-15,18-diacid-16,19-dilactone | — | — | — | <i>Baccharis articulate</i> | <i>Phytochemistry</i> , 1993, 34 , 1087-1090 |
| 461 | gaudichanolide A | OH | — | — | <i>Baccharis gaudichaudiana</i> | <i>J. Nat. Prod.</i> , 2005, 68 , 1121-1124 |
| 462 | gaudichanolide B | H | — | — | | |
| 463 | cephaloziellin C | =O | OH | — | <i>Cephaloziella kiaeri</i> | <i>J. Nat. Prod.</i> , 2013, 76 , 1700-1708 |
| 464 | cephaloziellin D | OH | =O | — | | |
| 465 | amphiacrolide F | =O | H | — | <i>Amphiachyris dracunculoides</i> | <i>J. Nat. Prod.</i> , 1996, 59 , 5-14 |
| 466 | ballatenolide A | αH | — | — | <i>Ballota limbata</i> | <i>Helv. Chim. Acta</i> , 2004, 87 , 682-689 |

| | | | | | | |
|-----|--|--------------|-------------|---|-----------------------------------|---|
| 467 | limbatolide A | β H | — | — | <i>Ostegia limbata</i> | <i>Chem. Pharm. Bull.</i> , 2005, 53 , 378-381 |
| 468 | crispene E | — | — | — | <i>Tinospora crispa</i> | <i>Org. Biomol. Chem.</i> , 2015, 13 , 3882-3886 |
| 469 | 1 α -acetoxy-8 β ,12-epoxy-15-hydroxy- <i>cis-ent</i> -cleroda-13-en-16,15:18 α ,6 α -diolide | α OAc | OH | — | <i>Adelanthus lindenbergianus</i> | <i>Phytochemistry</i> , 2004, 65 , 127–137 |
| 470 | 8 β ,12-epoxy-15 α -hydroxy-transcleroda-13-en-16,15:18 α ,6 α -diolide | H | α OH | — | | |
| 471 | 8 β ,12-epoxy-15 β -hydroxy-transcleroda-13-en-16,15:18 α ,6 α -diolide | H | β OH | — | | |
| 472 | 8 β ,12-epoxy-16 α -hydroxy-transcleroda-13-en-15,16:18 α ,6 α -diolide | α OH | — | — | | |
| 473 | 8 β ,12-epoxy-16 β -hydroxy-transcleroda-13-en-15,16:18 α ,6 α -diolide | β OH | — | — | | |
| 474 | 7 β ,12:8 β ,12-diepoxy-16 α -hydroxy- <i>cis-ent</i> -cleroda-13-en-15,16:18 α ,6 α -diolide | α OH | — | — | | |
| 475 | 7 β ,12:8 β ,12-diepoxy-16 β -hydroxy- <i>cis-ent</i> -cleroda-13-en-15,16:18 α ,6 α -diolide | β OH | — | — | | |
| 476 | 7 β ,12:8 β ,12-diepoxy-15-hydroxy- <i>cis-ent</i> -cleroda-13-en-16,15:18 α ,6 α -diolide | — | — | — | | |
| 477 | 1 β ,12-epoxy-16-hydroxy- <i>cis-ent</i> -cleroda-13-en-15,16:18 α ,6 α -diolide | — | — | — | | |
| 478 | ciliatolide B | OEt | OH | — | <i>Scapania ciliata</i> | <i>Chem. Biodivers.</i> , 2013, 10 , 1606-1612 |
| 479 | ciliatolide C | OMe | OH | — | | |
| 480 | ciliatolide D | OEt | OEt | — | | |

2.3.1.2 Type III Subtype Ib with Other O-Containing Rings (Table 15)




| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | R ₇ | Source | Ref. |
|-----|--|-----------------|----------------|----------------|----------------|----------------|----------------|------------------|-------------------------------|---|
| 481 | scutorientalin C | H | OH | OiBu | H | βOH | βOAc | H | <i>Scutellaria orientalis</i> | <i>Phytochemistry</i> , 1996, 43 , 173-178 |
| 482 | scutalpin H | H | OAc | Y ₁ | βH | βOH | βOAc | H | <i>Scutellaria alpina</i> | <i>Phytochemistry</i> , 1995, 38 , 181-187 |
| 483 | scutalpin I | H | OAc | OBz | βH | βOH | βOAc | H | | |
| 484 | scutalpin L | H | OAc | OBz | βOBz | βOH | βH | H | | |
| 485 | (12 <i>S</i>)-6 <i>α</i> -acetoxy-4 <i>α</i> ,18-epoxy-12-hydroxy-19-tigloyloxy- <i>neo</i> -clerod-13-en-15,16-olide | H | OTig | OAc | H | H | H | βOH | <i>Ajuga ciliata</i> | <i>Phytochem. Lett.</i> , 2012, 5 , 563-566 |
| 486 | scutalpin N | H | OAc | OBz | βOBz | βOH | H | H | <i>Scutellaria alpina</i> | <i>Phytochemistry</i> , 1998, 49 , 2449-2452 |
| 487 | ajugapantin A | βOAc | OAc | OAc | H | H | H | αOAc | <i>Ajuga pantantha</i> | <i>Phytochemistry</i> , 1993, 34 , 1091-1094 |
| 488 | ajugamacrin C | βOiBu | OAc | OAc | H | H | H | αOiBu | | |
| 489 | ajugamacrin D | βOiBu | OAc | OAc | H | H | H | αOY ₁ | | |
| 490 | ajugamacrin E | βY ₁ | OAc | OAc | H | H | H | αOiBu | | |

| | | | | | | | | | | |
|-----|---|----------------|-------------|-----------------|------------------------|------------|-------------|------------------------|------------------------------------|---|
| 491 | ajugamacrin A | β OAc | OAc | OAc | H | H | H | β iBu | <i>Ajuga macrosperma</i> | <i>Phytochemistry</i> , 1993, 33 , 887-889 |
| 492 | ajugamacrin B | β OAc | OAc | OAc | H | H | H | β Y ₁ | | |
| 493 | ajugatakasins A | OTig | OAc | OAc | H | H | H | OTig | <i>Ajuga decumbens</i> | <i>Biosci. Biotechnol. Biochem.</i> , 1997, 61 , 1518-1522 |
| 494 | ajugatakasins B | Y ₁ | OAc | OAc | H | H | H | Y ₁ | | |
| 495 | ajugamarin L2 | H | OTig | OH | H | H | H | H | <i>Ajuga nipponensis</i> | <i>Chin. Chem. Lett.</i> , 1995, 6 , 581-582 |
| 496 | (4S,11S)-11-acetoxy-8 β ,19-dihydroxy-6 α -tigloyloxy-4,18-epoxy- <i>neo</i> -clerod-13-en-15,16-olide | H | OH | OTig | H | β OH | OAc | H | <i>Scutellaria alpina</i> | <i>Yakugaku Zasshi</i> , 1994, 114 , 264-271, <i>Phytochemistry</i> , 1993, 34 , 1589-1594 |
| 497 | scutalpin C | H | OAc | OTig | H | OH | β OH | H | | |
| 498 | scupolin A | H | OH | Y ₁ | H | β OH | β OAc | H | <i>Scutellaria polyodon</i> | <i>J. Nat. Prod.</i> , 1997, 60 , 1229-1235 |
| 499 | scupolin B | H | OAc | OAc | H | β OH | β OAc | H | | |
| 500 | scupolin C | H | OBz | OBz | H | β OH | β OAc | H | | |
| 501 | scupolin D | H | OH | OH | β Y ₁ | β OH | β OAc | H | | |
| 502 | scupolin E | H | OH | Y ₁ | β OH | β OH | β OAc | H | | |
| 503 | scupolin F | — | — | — | — | — | — | — | | |
| 504 | scutorientalin E | H | OAc | OCin | β OAc | β OH | H | H | <i>Scutellaria orientalis</i> | <i>Phytochemistry</i> 1997, 46 , 587-589 |
| 505 | hastifolin A | H | H | OCin | H | β OH | H | H | <i>Scutellaria hastifolia</i> | <i>Phytochemistry</i> , 2010, 71 , 2087-2091 |
| 506 | ajugalide A | β OH | OAc | OAc | H | H | H | α OAc | <i>Ajuga taiwanensis</i> | <i>Chem. Pharm. Bull.</i> , 2005, 53 , 164-167 |
| 507 | ajugalide B | β OAc | OAc | OAc | H | H | H | α OH | | |
| 508 | ajugalide C | H | OAc | OAc | H | H | H | α OH | | |
| 509 | (4S)-19-acetoxy-8 β -hydroxy-6 α -tigloyloxy-4,18-epoxy- <i>neo</i> -cleroda-11,13-dien-15,16-olide | α OTig | H | — | — | — | — | — | <i>Scutellaria alpina</i> | <i>Yakugaku Zasshi</i> , 1994, 114 , 264-271 <i>Phytochemistry</i> , 1995, 38 , 181-187 |
| 510 | scutalpin J | α OBz | H | — | — | — | — | — | | |
| 511 | scutalpin K | α OBz | β OBz | — | — | — | — | — | | |
| 512 | ajugacumbin E | OAc | β OAc | X ₁₂ | H | — | — | — | <i>Ajuga decumbens</i> | <i>Chem. Pharm. Bull.</i> , 1990, 38 , 3167-3168 |
| 513 | 3 α -hydroxyajugamarin F4 | H | α OH | OAc | β Y ₁ | — | — | — | <i>Ajuga reptans</i> | <i>Phytochemistry</i> , 1998, 47 , 1227-1232 |
| 514 | scuterivulactone A | α OBz | H | β OH | β OAc | — | — | — | <i>Scutellaria rivularis</i> | <i>Chem. Pharm. Bull.</i> , 1997, 45 , 152-160 |
| 515 | seguiniilactone A | α OAc | β OAc | H | H | — | — | — | <i>Colquhounia seguinii</i> | <i>J. Integr. Plant Biol.</i> , 2014, 56 , 928-940 |
| 516 | seguiniilactone B | α OAc | — | — | — | — | — | — | | |
| 517 | seguiniilactone C | H | — | — | — | — | — | — | | |
| 518 | amphiacrolide K | — | — | — | — | — | — | — | <i>Amphiachyris dracunculoides</i> | <i>J. Nat. Prod.</i> , 1996, 59 , 463-468 |
| 519 | Conyzalactone | — | — | — | — | — | — | — | <i>Conyza blinii</i> | <i>Heterocycles</i> , 1999, 51 , 605-609 |

| | | | | | | | | | | |
|-----|---|-----|-----|----|---|---|---|---|--------------------------------|--|
| 520 | (12 <i>S</i>)-1 α ,19-epoxy-6 α ,18-diacetoxy-4 α ,12-dihydroxy- <i>neo</i> -clerod-13-en-15,16-olide | — | — | — | — | — | — | — | <i>Ajuga decumbens</i> | <i>Fitoterapia</i> , 2012, 83 , 1409-1414 |
| 521 | ptycholide I | =O | OMe | =O | — | — | — | — | <i>Ptychopetalum olacoides</i> | <i>Nat. Prod. Commun.</i> , 2011, 6 , 327-332 |
| 522 | ptycholide II | OMe | OMe | =O | — | — | — | — | | |
| 523 | ptycholide III | OMe | H | =O | — | — | — | — | | |
| 524 | ptycholide IV | OMe | =O | OH | — | — | — | — | | |
| 525 | Teucrasiolide | — | — | — | — | — | — | — | <i>Teucrium asiaticum</i> | <i>Phytochemistry</i> , 1997, 45 , 383-385 |
| 526 | 6 α -hydroxycleroda-3,13-dien-15(16), 4(7)-diolide | — | — | — | — | — | — | — | <i>Pulicaria wightiana</i> | <i>Helv. Chim. Acta.</i> , 2008, 91 , 2081-2088 |
| 527 | microdon B | — | — | — | — | — | — | — | <i>Gomphostemma microdon</i> | <i>Z. Naturforsch., B: J. Chem. Sci.</i> , 2009, 64 , 443-446 |

3.2.1. Type III Subtype IIa with C3/C4 Double Bond (Table 16)



| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | R ₇ | Source | Ref. |
|-----|--|----------------|--------------------|-----------------------------|----------------|----------------|----------------|----------------|----------------------------------|---|
| 530 | <i>ent</i> -2 β ,18,19-trihydroxycleroda-3,13-dien-16,15-olide | α OH | CH ₂ OH | α CH ₂ OH | H | H | H | — | <i>Crassocephalum bauchiense</i> | <i>Nat. Prod. Res.</i> , 2015, 29 , 1990-1994 |
| 531 | (-)-6 α -hydroxy-5 α ,8 α ,9 α ,10 α -cleroda-3,13-dien-16,15-olid-18-oic acid* | H | CO ₂ H | α Me | α OH | H | H | — | <i>Dodonaea viscosa</i> | <i>Phytochem. Lett.</i> , 2014, 8 , 10-15 |
| 532 | 15-hydroxy-16-oxo-15,16H-hardwickiic acid | H | CO ₂ H | α Me | H | H | OH | — | <i>Grangea maderaspatana</i> | <i>Phytochemistry</i> , 1999, 52 , 1341-1343 |
| 533 | limbatolide B | H | CO ₂ H | β Me | H | H | OMe | — | <i>Otostegia limbata</i> | <i>Chem. Pharm. Bull.</i> , 2005, 53 , 378-381 |
| 534 | limbatolide C | H | CO ₂ H | β Me | H | H | H | — | | |
| 535 | 12(<i>S</i>)-hydroxycleroda-3,13-dien-16,15-olide | H | Me | α Me | H | β OH | H | — | <i>Callicarpa americana</i> | <i>J. Nat. Prod.</i> , 2007, 70 , 372-377 |
| 536 | 12(<i>S</i>),16 ζ -dihydroxycleroda-3,13-dien-15,16-olide | H | Me | β OH | OH | — | — | — | | |
| 537 | 12(<i>S</i>)-hydroxy-16 ζ -methoxycleroda-3,13-dien-15,16-olide | H | Me | β OH | OMe | — | — | — | | |
| 538 | 12(<i>S</i>)-hydroxycleroda-3,13-dien-15,16-olide | H | Me | β OH | H | — | — | — | | |



| | | | | | | | | | | |
|-------|--|----------------------|---------------------|-----|------|------|-----|-----|---|--|
| 539 | 2β-methoxy-cleroda-3,13-dien-18-carboxy-15,16-olide | OMe | CO ₂ H | H | H | — | — | — | <i>Casearia sylvestris</i> | <i>Fitoterapia</i> , 2009, 80 , 404–407 |
| 540 | 16(R)-3,13Z-kolavadien-16,15-olide-2-one | =O | Me | H | αOH | — | — | — | <i>Polyalthia viridis</i> | <i>Phytochemistry</i> , 1990, 29 , 653-655 |
| 541 | 16(S)-3,13Z-kolavadien-16,15-olide-2-one | =O | Me | H | βOH | — | — | — | | |
| 542 | polylongifoliaon A | =O | Me | H | αOMe | — | — | — | <i>P. longifolia</i> var. <i>pendula</i> | <i>RSC Advances</i> , 2014, 4 , 23707-23712 |
| 543 | polylongifoliaon B | =O | Me | H | βOMe | — | — | — | | |
| 544 | 2β-(2-methylbutanoyl)cleroda-3,13-dien-15,16-olid-18-oic acid | Y ₁ | CO ₂ H | H | H | — | — | — | <i>Pulicaria wightiana</i> | <i>Helv. Chim. Acta.</i> , 2008, 91 , 2081-2088 |
| 545 | methyl 6-oxocleroda-3,13-dien-15,16-olid-18-oate | CO ₂ Me | αMe | =O | H | H | — | — | | |
| 546 | methyl 6α-hydroxycleroda-3,13-dien-15,16-olid-18-oate | CO ₂ Me | αMe | αOH | H | H | — | — | | |
| 547 | methyl 6α,7α-dihydroxycleroda-3,13-dien-15,16-olid-18-oate | CO ₂ Me | αMe | αOH | αOH | H | — | — | | |
| 548 | clerodermic acid | CO ₂ H | αMe | H | H | H | — | — | <i>Clerodendrum inerm</i> | <i>Phytochemistry</i> , 1990, 29 , 3671-3673 |
| 549 | (-)-12,16-dihydroxy- <i>cis</i> -cleroda-3,13-dien-15-oic acid-15,16-olide | CO ₂ H | βMe | H | H | OH | — | — | <i>Croton schiedeanus</i> | <i>Phytochemistry</i> , 1999, 51 , 643-649 |
| 550 | conyhypolide A | CO ₂ H | αMe | H | αOH | OH | — | — | <i>Conyza hypoleuca</i> | <i>Phytochemistry</i> 1991, 30 , 575-581 |
| 551 | 16α-hydroxy-cleroda-3,13Z-diene-15,16-olide | Me | αMe | H | H | αOH | — | — | <i>Polyalthia longifolia</i> | <i>Fitoterapia</i> , 2005, 76 , 336-339 |
| 552 | 16α-methoxycleroda-3,13Z-dien-16,15-olide | Me | αMe | H | H | αOMe | — | — | | <i>J. Nat. Prod.</i> , 1992, 55 , 256-258 |
| 553 | 7-oxo- <i>ent</i> -clerodan-3,13-dien-15,16-olide | Me | αMe | H | =O | H | — | — | <i>Platychaete aucheri</i> | <i>Phytochemistry</i> , 1990, 29 , 985-987 |
| 554 | 6-hydroxy-7-oxo- <i>ent</i> -clerodane-3,13-dien-15,16-olide | Me | αMe | αOH | =O | H | — | — | | |
| 555 | thymonin | CH ₂ OH | αCH ₂ OH | αOH | H | H | — | — | <i>Salvia thymoides</i> | <i>Phytochemistry</i> , 1997, 46 , 1249-1254 |
| 556 | 7β-hydroxythymonin | CH ₂ OH | αCH ₂ OH | αOH | βOH | H | — | — | | |
| 557 | souldiol | CH ₂ OH | αCH ₂ OH | H | H | H | — | — | <i>Aster souliei</i> | <i>Chin. Chem. Lett.</i> , 1996, 7 , 619-620 |
| 558 | solidagoic acid H | Me | — | — | — | — | — | — | <i>Solidago virgaurea</i> | <i>Phytochemistry</i> , 2010, 71 , 104–109 |
| 559 | solidagoic acid I | CH ₂ OAng | — | — | — | — | — | — | | |
| 560 | solidagoic acid C | Me | αCO ₂ H | H | H | αMe | βMe | H | | |
| 561 | solidagoic acid D | CH ₂ OAng | αCO ₂ H | H | H | αMe | βMe | H | | |
| 562 | solidagoic acid E | Me | αCO ₂ H | H | H | αMe | βMe | OH | | |
| 563 | solidagoic acid F | CH ₂ OAng | αCO ₂ H | H | H | αMe | βMe | OH | | |
| 564 | solidagoic acid G | Me | αCO ₂ H | H | H | αMe | βMe | OMe | | |
| 565** | methyl 6,15-dihydroxycleroda-3,13-dien-16,15-olid-18-oate | CO ₂ Me | αMe | αOH | H | αMe | αMe | OH | <i>Pulicaria wightiana</i> | <i>Helv. Chim. Acta.</i> , 2008, 91 , 2081-2088 |
| 566 | (-)-6β-hydroxy-5β,8β,9β,10α-cleroda-3,13-dien-16,15-olid-18-oic acid | CO ₂ H | βMe | βOH | H | βMe | βMe | H | <i>Duranta repens</i> | <i>Chem. Pharm. Bull.</i> , 2004, 52 , 785-789 |
| 567 | (+)-3,13-clerodadien-16,15-olid-18-oic acid | CO ₂ H | βMe | H | H | βMe | βMe | H | | |
| 568 | (+)-7β-acetoxy-18-oxo-3,13-clerodadien-16,15-olide | CHO | βMe | H | βOAc | βMe | βMe | H | <i>Sindora sumatrana</i> | <i>Chem. Pharm. Bull.</i> , 1994, 42 , |

| | | | | | | | | | | |
|-----|---|-------------------|------------|-------------|-------------|------------|------------|----|----------------------------|---|
| 569 | (+)-7 β -acetoxy-3,13-clerodadien-16,15-olid-18-oic acid | CO ₂ H | β Me | H | β OAc | β Me | β Me | H | | 1202-1207 |
| 570 | (+)-7 β -acetoxy-16-hydroxy-3,13-clerodadien-16,15-olid-18-oic acid | CO ₂ H | β Me | H | β OAc | β Me | β Me | OH | | |
| 571 | scapanialide A | — | — | — | — | — | — | — | <i>Scapania parva</i> | <i>Phytochem. Lett.</i> , 2012, 5 , 535–540 |
| 572 | scapanialide C | H | β OH | β Me | OH | — | — | — | | |
| 573 | 2-oxo-5 α ,8 α -cleroda-3,13-dien-16,15-olide | =O | H | α Me | H | — | — | — | <i>Vellozia bicolor</i> | <i>Phytochemistry</i> , 1994, 37 , 1115-1117 |
| 574 | scutebata A | — | — | — | — | — | — | — | <i>Scutellaria barbata</i> | <i>J. Nat. Prod.</i> , 2010, 73 , 233-236 |
| 575 | trinerolactone | — | — | — | — | — | — | — | <i>Baccharis trinervis</i> | <i>Phytochemistry</i> , 1993, 34 , 1377-1384 |

* Compound's name indicates 10 α H; however, compound's structure was presented as an *ent*-clerodane with a 10 β H in *Phytochem. Lett.*, 2014, **8**, 10-15.

** C-10 H should be β

3.2.2. Type III Subtype IIb with Double Bonds in Other Positions (Tables 17 & 18)



| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | Source | Ref. |
|-----|--|-------------------------|----------------|----------------|----------------|----------------|----------------|-----------------------------|---|
| 576 | 16 ζ -hydroxycycloroda-3,11(<i>E</i>),13-trien-15,16-olide | H | H | H | OH | – | – | <i>Callicarpa americana</i> | <i>J. Nat. Prod.</i> , 2007, 70 , 372-377 |
| 577 | barbatin C | α OH | β OH | β OH | H | | | <i>Scutellaria barbata</i> | <i>J. Integr. Plant Biol.</i> , 2008, 50 , 699-702, <i>Fitoterapia</i> , 2010, 81 , 737-741, <i>Planta Med.</i> , 2011, 77 , 1536-1541, <i>Phytochemistry</i> , 2006, 67 , 1326-1330, <i>Planta Med.</i> , 2007, 73 , 1217-1220, <i>J. Asian Nat. Prod. Res.</i> , 2010, 12 , 859-864 |
| 578 | barbatin D | α OBz | β OBz | β OH | H | – | – | | |
| 579 | barbatin E | Y ₄ | M ₁ | β OH | H | – | – | | |
| 580 | scutebata I | α OAc | β OH | β OH | H | – | – | | |
| 581 | scutebata J | α OBz | β OH | β OH | H | – | – | | |
| 582 | scutebata K | α Y ₉ | β OH | β OH | H | – | – | | |
| 583 | 6,7-di- <i>O</i> -acetoxybarbatin A | α OAc | β OAc | β OH | H | – | – | | |
| 584 | 6-(2,3-epoxy-2-isopropyl- <i>n</i> -propoxy)barbatin C | α M ₂ | β OH | β OH | H | – | – | | |
| 585 | 6-acetoxybarbatin C | α OAc | β OH | β OH | H | – | – | | |

| | | | | | | | | | |
|-----|--|----------------|--------------|-------------|------------|------------|-----------------------------|--|---|
| 586 | scuterivulactone D | H | – | – | – | | | <i>Scutellaria rivularis</i> | <i>Chem. Pharm. Bull.</i> , 1997, 45 , 152-160 |
| 587 | salvidivin C | OH | =O | – | – | | | <i>Salvia divinorum</i> | <i>J. Nat. Prod.</i> , 2006, 69 , 1782-1786 |
| 588 | salvidivin D | =O | OH | – | – | | | | |
| 589 | (-)-3 α ,16 α -dihydroxycleroda-4(18),13(14)Z-dien-15,16-olide | α OH | H | H | H | H | α OH | <i>Polyalthia longifolia</i> | <i>Nat. Prod. Res.</i> , 2010, 24 , 1687-1694 |
| 590 | 3 β ,16 α -dihydroxycleroda-4(18),13(14)Z-dien-15,16-olide | β OH | H | H | H | H | α OH | <i>Polyalthia barnesii</i> | <i>Phytochemistry</i> 1994, 37 , 1659-1662 |
| 591 | cleroda-4(18),13-dien-16,15-olide* | H | H | H | H | H | H | <i>Polyalthia longifolia</i> | <i>Phytochemistry</i> 1995, 38 , 189-194. |
| 592 | 16-hydroxycleroda-4(18),13-dien-16,15-olide* | H | H | H | H | H | OH (α/β 1:1) | | |
| 593 | — | H | H | H | H | H | β OH | | |
| 594 | 16(R&S)-methoxycleroda-4(18),13-dien-15,16-olide | H | H | H | H | H | OMe | <i>Polyalthia longifolia</i> var. <i>pendula</i> | <i>Molecules</i> , 2014, 19 , 2049-2060 |
| 595 | 3 α ,12(S)-dihydroxy-cleroda-4(18),13-dien-15,16-olide | β OH | H | H | H | β OH | H | <i>Callicarpa americana</i> | <i>J. Nat. Prod.</i> , 2007, 70 , 372-377 |
| 596 | Scutebaicalin | H | α OBz | β OBz | β OH | H | H | <i>Scutellaria baicalensis</i> | <i>Phytochemistry</i> , 1996, 43 , 835-837 |
| 597 | scutebata L | – | – | – | – | – | – | <i>Scutellaria barbata</i> | <i>Planta Med.</i> , 2011, 77 , 1536-1541 |
| 598 | calcicolin A | OiBu | – | – | – | – | – | <i>Glossocarya calcicola</i> | <i>Phytochemistry</i> , 2005, 66 , 2844-2850 |
| 599 | calcicolin B | OTig | – | – | – | – | – | | |
| 600 | calcicolin C | Y ₁ | – | – | – | – | – | | |

*Compounds' names (16,15-olide) and structures (15,16-olide) are shown as given in *Phytochemistry* 1995, **38**, 189-194.


Table 18. Type III Subtype IIb continued


| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | R ₇ | R ₈ | Source | Ref. |
|-----|---|----------------|----------------|--|----------------|----------------------|----------------|----------------|----------------|------------------------|---|
| 601 | (12S)-6 α ,18,19-triacetoxy-4 α ,12-dihydroxy-1 β -tigloyloxy- <i>neo</i> -clerod-13-en-15,16-olide | β OTig | H | β CH ₂ OAc | α OH | CH ₂ OAc | β OAc | β OH | H | <i>Ajuga ciliata</i> | <i>Fitoterapia</i> , 2011, 82 , 1123-1127 |
| 602 | ajugaciliatin I | H | H | β CH ₂ OAc | α OH | CH ₂ OTig | β OH | H | H | | <i>J. Nat. Prod.</i> , 2011, 74 , 1575– 1583 |
| 603 | ajugaciliatin J | H | H | β CH ₂ OTig | α OH | CH ₂ OH | β OH | H | H | | |
| 604 | (12S)-18,19-diacetoxy-4 α ,6 α ,12-trihydroxy-1 β -tigloyloxy- <i>neo</i> -clerod-13-en-15,16-olide | β OTig | H | β CH ₂ OAc | α OH | CH ₂ OAc | β OH | β OH | H | <i>Ajuga decumbens</i> | <i>Planta Med.</i> , 2012, 78 , 1579-1583 |
| 605 | 4 α ,6 α -dihydroxy-18-(4'-methoxy-4'-oxobutyryloxy)-19-tigloyloxy- <i>neo</i> -clerod-13-en-15,16-olide | H | H | β CH ₂ X ₇ | α OH | CH ₂ OTig | β OH | β H | H | | |

| | | | | | | | | | | | |
|-----|---|-------------|----|-----------------------------|-----------------------------|----------------------|-------------|-------------|-------------|--|---|
| 606 | 6 α ,18-diacetoxy-4 α -hydroxy-19-tigloyloxy- <i>neo</i> -clerod-13-en-15,16-olide | H | H | β CH ₂ OAc | α OH | CH ₂ OTig | β OAc | H | H | <i>Ajuga ciliate</i> | <i>Phytochem. Lett.</i> , 2012, 5 , 563-566 |
| 607 | ajugalide D | H | H | H | α CO ₂ Me | Me | α OH | α OH | H | <i>Ajuga taiwanensis</i> | <i>Chem. Pharm. Bull.</i> , 2005, 53 , 164-167 |
| 608 | ajugacumbin F | H | H | β CH ₂ OH | α OH | CH ₂ OTig | α OH | H | H | <i>Ajuga decumbens</i> | <i>Chem. Pharm. Bull.</i> , 1990, 38 , 3167-3168 |
| 609 | 4 β ,16 α -dihydroxy-cleroda-13(14) <i>Z</i> -en-15,16-olide | H | H | β OH | α Me | Me | H | H | α OH | <i>Polyalthia barnesii</i> | <i>Phytochemistry</i> 1994, 37 , 1659-1662 |
| 610 | 16-hydroxycleroda-13-ene-15,16-olide-3-one | H | =O | H | α Me | Me | H | H | OH | <i>Polyalthia longifolia</i> var. <i>pendula</i> | <i>Planta Med.</i> , 2006, 72 , 1344-1347 |
| 611 | methyl(1 α ,4 $\alpha\alpha$,5 α ,6 β ,8 $\alpha\alpha$)-5-[2-(3-furan-3-ene-2-one)ethyl]-1,2,3,4,4a,5,6,7,8,8a-decahydro-1,2-dihydroxy-1-naphthalene carboxylate | - | - | - | - | - | - | - | - | <i>Tinospora rumphii</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 509-511 |
| 612 | dytesinin A | α Me | OH | - | - | - | - | - | - | <i>Cystodytes</i> sp | <i>Tetrahedron</i> 2000, 56 , 7923-7926 |
| 613 | dytesinin B | α Me | H | - | - | - | - | - | - | | |
| 614 | echinoclerodane A | β Me | OH | - | - | - | - | - | - | <i>Echinomuricea</i> sp | <i>Molecules</i> , 2012, 17 , 9443-9450 |

4. Type IV with a 5-(3-Furyl)- δ -valerolactone-based Side Chain at C-9

4.1. Type IV Subtype I with *O*-Containing Rings (Table 19)






| No. | Compound Name | R ₁ | R ₂ | R ₃ | Source | Ref. |
|-----|--|----------------|----------------|----------------|---------------------------|---|
| 615 | tehuanin D | — | — | — | <i>Salvia herbacea</i> | <i>J. Nat. Prod.</i> , 2012, 75 , 951-958 |
| 616 | salvimicrophyllin C | — | — | — | <i>Salvia microphylla</i> | <i>J. Nat. Prod.</i> , 2014, 77 , 1088-1092 |
| 617 | salvimicrophyllin D | — | — | — | | |
| 618 | dugesin E | — | — | — | <i>Salvia dugesii</i> | <i>Nat. Prod. Bioprospect.</i> , 2011, 1 , 81-86 |
| 619 | <i>ent</i> -(5 <i>R</i> ,9 <i>R</i>)-15,16-epoxy-10 <i>S</i> -hydroxy-cleroda-3,13(16),14-triene-17,12 <i>S</i> ;18,19-diolide | αOH | H | — | <i>Salvia haenkei</i> | <i>Tetrahedron</i> , 1997, 53 , 14719-14728 |
| 620 | — | βH | αOH | — | <i>Salvia miniata</i> | <i>Phytochemistry</i> , 2011, 72 , 265-275 |
| 621 | — | βOH | H | — | | |
| 622 | — | — | — | — | | |
| 623 | — | — | — | — | | |
| 624 | <i>ent</i> -(5 <i>R</i> ,9 <i>R</i>)-15,16-epoxy-10 <i>S</i> -hydroxycleroda-3,7,13(16),14-tetraene-17,12 <i>S</i> ,18,19-diolide | — | — | — | <i>Salvia haenkei</i> | <i>Tetrahedron</i> , 1997, 53 , 14719-14728 |
| 625 | <i>ent</i> -(4 <i>S</i> ,5 <i>R</i> ,9 <i>S</i> ,10 <i>R</i>)-15,16-epoxycleroda-1,13(16),14-trien-17,12 <i>S</i> ;18,19-diolide | αH | αH | αH | | |
| 626 | infuscatin | βOH | βOH | αOH | <i>Salvia infuscata</i> | <i>Phytochem. Anal.</i> , 1994, 5 , 302-304 |
| 627 | tehuanin E | OH | — | — | <i>Salvia herbacea</i> | <i>J. Nat. Prod.</i> , 2012, 75 , 951-958 |
| 628 | 1β,10β-epoxysalviarin | H | — | — | | |
| 629 | tehuanin F | OH | βH | — | | |
| 630 | tehuanin G | H | αOH | — | | |
| 631 | tehuanin H | H | H | — | | |
| 632 | 1α,2α-epoxy-3,4α-dihydrolinearolactone | αH | — | — | <i>Salvia reptans</i> | <i>Phytochemistry</i> , 1991, 30 , 2335-2338 |
| 633 | polystachyne D | βH | — | — | <i>Salvia polystachya</i> | <i>Phytochemistry</i> , 2000, 53 , 103-109 |
| 634 | polystachyne E | — | — | — | | |
| 635 | tehuanin A | — | — | — | <i>Salvia herbacea</i> | <i>J. Nat. Prod.</i> , 2012, 75 , 951-958 |

| | | | | | | |
|-----|---|-------------------|-------------------|------------|--------------------------------|---|
| 636 | tehuanin B | H | — | — | | |
| 637 | tehuanin C | OH | — | — | | |
| 638 | 15,16-epoxy,1,3,13(16),14-clerodatetraene-17,12:18,6-diolide | — | — | — | | |
| 639 | 15-carboxy-8 β ,16-dihydroxy-1,3,13E-clerodatriene-17,12:18,6-diolide | CO ₂ H | H | — | <i>Jamesoniella autumnalis</i> | <i>Phytochemistry</i> , 1998, 48 , 681-685 |
| 640 | 15-carboxy-8 β ,16-dihydroxy-1,3,13Z-clerodatriene-17,12:18,6-diolide | H | CO ₂ H | — | | |
| 641 | fibrauretin A | — | — | — | <i>Fibraurea tinctoria</i> | <i>J. Nat. Prod.</i> , 2007, 70 , 1930-1933 |
| 642 | tinospin E | — | — | — | | <i>Chem. Pharm. Bull.</i> , 2012, 60 , 1324-1328 |
| 643 | tinosporin A | — | — | — | <i>Tinospora sagittata</i> | <i>Phytochem. Lett.</i> , 2015, 12 , 173-176 |
| 644 | tinosporin B | H | β OH | β H | | |
| 645 | 6-hydroxycolumbin | α OH | H | β H | <i>Penianthus zenkeri</i> | <i>Phytochemistry</i> , 1991, 30 , 1957-1962 |
| 646 | 6-hydroxyisocolumbin | α OH | H | α H | | |
| 647 | epi-8-hydroxycolumbin | — | — | — | | |
| 648 | fibrauretin B | α H | — | — | <i>Fibraurea tinctoria</i> | <i>Bioorg. Med. Chem.</i> , 2008, 16 , 9603-9609 |
| 649 | fibrauretin C | β OH | — | — | | |
| 650 | antadiosbulbin A | OH | H | α H | <i>Dioscorea antaly</i> | <i>Phytochemistry</i> , 2010, 71 , 1007-1013 |
| 651 | antadiosbulbin B | OH | H | β H | | |
| 652 | bafoudiosbulbin F | H | H | β OH | <i>Dioscorea bulbifera</i> | <i>Phytochemistry</i> , 2008, 69 , 2374-2379 |
| 653 | bafoudiosbulbin G | H | α OAc | β OH | | |
| 654 | 3,4,15,16-diepoxy-cleroda-13(16),14-diene-12,17-olide | β Me | H | — | <i>Croton oblongifolius</i> | <i>Phytochem. Lett.</i> , 2011, 4 , 147-150 |
| 655 | ravidin A | α Me | =O | — | <i>Nannoglottis ravida</i> | <i>Phytochemistry</i> , 2004, 65 , 2533-2537 |

4.2. Type IV Subtype II Other Compounds (Table 20)




| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | Source | Ref. |
|-----|------------------------------------|----------------|----------------|--------------------|----------------|------------------------------|--|
| 656 | microdon A | H | =O | Me | =O | <i>Gomphostemma microdon</i> | <i>Z. Naturforsch., B: J. Chem. Sci.</i> , 2009, 64 , 443-446 |
| 657 | 12-epi-methyl-barbascoate | H | H | CO ₂ Me | =O | <i>Croton ururucana</i> | <i>Phytochemistry</i> , 1998, 49 , 171-174 |
| 658 | 17-oxo-1,2-dihydrowelwitschic acid | H | H | CO ₂ H | =O | <i>Conyza welwitschii</i> | <i>Phytochemistry</i> , 1990, 29 , 2247-2252 |
| 659 | salvinorin C | OAc | OAc | CO ₂ Me | =O | <i>Salvia divinorum</i> | <i>Org. Lett.</i> , 2001, 3 , 3935-3937 |
| 660 | salvinorin D | OAc | OH | CO ₂ Me | =O | | <i>J. Nat. Prod.</i> , 2003, 66 , 703-705 |
| 661 | salvinorin E | OH | OAc | CO ₂ Me | =O | | |
| 662 | salvinorin F | OH | H | CO ₂ Me | =O | | |
| 663 | salvinorin H | OH | OH | CO ₂ Me | =O | | |
| 664 | salvinorin I | OH | OH | CO ₂ Me | βOH | | |
| 665 | salvinorin J | OH | OAc | CO ₂ Me | α,β-OH | | |
| 666 | penianthic acid methyl ester | — | — | — | — | <i>Penianthus zenkeri</i> | <i>Phytochemistry</i> , 1991, 30 , 1957-1962 |
| 667 | 8-epicordatin | — | — | — | — | <i>Croton palanostigma</i> | <i>J. Braz. Chem. Soc.</i> , 2010, 21 , 731-739 |
| 668 | tinocallone A | βMe | βH | — | — | <i>Tinospora capillipes</i> | <i>Chin. Chem. Lett.</i> , 1992, 3 , 185-188 |
| 669 | tinocallone B | αMe | αH | — | — | | |

| | | | | | | | |
|-----|---|--------------|-----------------|--------------------|------------|-----------------------------------|---|
| 670 | tinocallone C | H | =O | H | β H | | |
| 671 | tinocallone D | H | =O | H | α H | | |
| 672 | tincordin | OH | =O | OH | β H | <i>Tinospora cordifolia</i> | <i>Nat. Prod. Res.</i> , 2013, 27 , 1431-1436 |
| 673 | crotonolide E | =O | α Me | — | — | <i>Croton laui</i> | <i>J. Nat. Prod.</i> , 2014, 77 , 1013-1020 |
| 674 | crotonolide F | α OH | CH ₂ | — | — | | |
| 675 | — | α H | =O | CO ₂ Me | — | <i>Croton jimenezii</i> | <i>Ingenieria Y Ciencia Quimica</i> , 2000, 19 , 68-73 |
| 676 | — | β H | =O | CO ₂ Me | — | | |
| 677 | 17-oxo-10-epi-welwitschic acid | α H | =O | CO ₂ H | — | <i>Conyza welwitschii</i> | <i>Phytochemistry</i> , 1990, 29 , 2247-2252 |
| 678 | 17-oxo-welwitschic acid | β H | =O | CO ₂ H | — | | |
| 679 | 17 α -hydroxywelwitschic acid | β H | α OH | CO ₂ H | — | | |
| 680 | 17 β -hydroxywelwitschic acid | β H | β OH | CO ₂ H | — | | |
| 681 | 2-epi-8-epi-salvinorin A | — | — | — | — | the smoke of salvinorin A | <i>Tetrahedron Lett.</i> , 2010, 51 , 5207-5209 |
| 682 | methyl 3-oxo-12-epibarbascoate | =O | H | CO ₂ Me | — | <i>Croton urucurana</i> | <i>J. Braz. Chem. Soc.</i> , 2013, 24 , 609-614. |
| 683 | furocrotinsulolide A | α OH | β OH | α Me | — | <i>Croton insularis</i> | <i>Helv. Chim. Acta.</i> , 2005, 88 , 2654-2660 |
| 684 | salvinicin A | β OMe | β OMe | — | — | <i>Salvia divinorum</i> | <i>Org. Lett.</i> , 2005, 7 , 3017-3020 |
| 685 | salvinicin B | α OMe | α OMe | — | — | | |
| 686 | salvidivin A | =O | OH | — | — | | <i>J. Nat. Prod.</i> , 2006, 69 , 1782-1786 |
| 687 | salvidivin B | OH | =O | — | — | | |
| 688 | cornutin A | — | — | — | — | <i>Cornutia grandifolia</i> | <i>J. Org. Chem.</i> , 1992, 57 , 862-866 |
| 689 | 1 β ,12:15,16-diepoxy-cis-ent-cleroda-13(16),14-dien-18 α ,6 α -olide | — | — | — | — | <i>Adelanthus lindenbergianus</i> | <i>Phytochemistry</i> , 2004, 65 , 127-137 |
| 690 | orcadensin | — | — | — | — | | |
| 691 | salvianduline D | — | — | — | — | <i>Salvia miniata</i> | <i>Phytochemistry</i> , 2011, 72 , 265-275 |

5. Type V with an α -Spiro-attached 4-(3-Furyl)- γ -butyrolactone-based Side Chain at C-9


5.1. Type V Subtype I with Various O-Containing Rings (Table 21)



| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | Source | Ref. |
|-----|---------------------------------------|----------------|----------------|----------------|----------------|---------------------------|---|
| 692 | <i>trans</i> -1,2-dihydrosalvifaricin | — | — | — | — | <i>Salvia fulgens</i> | <i>J. Nat. Med.</i> , 2006, 60 , 58-63 |
| 693 | — | =O | β Me | α OAc | — | <i>Salvia miniata</i> | <i>Phytochemistry</i> , 2011, 72 , 265-275 |
| 694 | — | α OH | α Me | β OAc | — | | |
| 695 | salvifolin | — | — | — | — | <i>Salvia tiliaefolia</i> | <i>J. Org. Chem.</i> , 1990, 55 , 3522-3525 |
| 696 | dugesin F | — | — | — | — | <i>Salvia dugesii</i> | <i>Nat. Prod. Bioprospect.</i> , 2011, 1 , 81-86 |
| 697 | teupolin XII | H | β OMe | β OH | H | <i>Teucrium polium</i> | <i>Phytochemistry</i> , 2011, 72 , 2037-2044 |
| 698 | teucvisin A | OH | =O | α OH | H | <i>Teucrium viscidum</i> | <i>Chem. Pharm. Bull.</i> , 2014, 62 , 472-476 |
| 699 | teuperminB | H | =O | =O | OH | <i>Teucrium perny</i> | <i>Phytochemistry</i> , 1991, 30 , 1963-1966 |
| 700 | teuperminC | H | =O | β OH | OH | | |

| | | | | | | | |
|-----|---|--------------|-----|-----|---|-----------------------------|---|
| 701 | teupernin A | — | — | — | — | | |
| 702 | cephaloziellin A | — | — | — | — | <i>Cephaloziella kiaeri</i> | <i>J. Nat. Prod.</i> , 2013, 76 , 1700-1708 |
| 703 | cephaloziellin B | — | — | — | — | | |
| 704 | teupolin X | α OH | OH | — | — | <i>Teucrium polium</i> | <i>Phytochemistry</i> , 2011, 72 , 2037-2044. |
| 705 | teupolin XI | α OMe | OH | — | — | | |
| 706 | teucvisin B | =O | OAc | — | — | <i>Teucrium viscidum</i> | <i>Chem. Pharm. Bull.</i> , 2014, 62 , 472-476 |
| 707 | 12- <i>epi</i> -montanin D | H | — | — | — | <i>Teucrium maghrebinum</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 1029-1031 |
| 708 | teusandrin C | OH | — | — | — | <i>Teucrium sandrasicum</i> | <i>Phytochemistry</i> , 1997, 45 , 1653-1662 |
| 709 | teusandrin D | H | — | — | — | | |
| 710 | teusandrin E | =O | — | — | — | | |
| 711 | teusandrin F | α OH | — | — | — | | |
| 712 | 4 β -hydroxy-3 α ,6 α ,15,16-bis-epoxy-6 β -methoxy- <i>cis</i> -clerodane-13(16),14-dien-20,12-olide | — | — | — | — | <i>Pteronia eenii</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |
| 713 | crotonolide A | — | — | — | — | <i>Croton laui</i> | <i>J. Nat. Prod.</i> , 2014, 77 , 1013-1020 |
| 714 | crotonolide B | H | OH | H | — | | |
| 715 | isocrotonolide B | OH | H | H | — | | |
| 716 | crotonolide C | H | OH | OAc | — | | |
| 717 | isocrotonolide C | OH | H | OAc | — | | |
| 718 | crotonolide D | H | OH | — | — | | |
| 719 | isocrotonolide D | OH | H | — | — | | |

5.2. Type V Subtype II with 4,18-; 3,4-; or 8,17-Oxirane Moieties (Table 22)




| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | R ₇ | R ₈ | Source | Ref. |
|-----|------------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|-----------------------------|---|
| 720 | teumassin | OH | H | OAc | =O | H | H | =O | H | <i>Teucrium massiliense</i> | <i>Phytochemistry</i> , 1992, 31 , 4366-4367 |
| 721 | 4 α ,18-epoxy-tafricanin A | H | =O | OAc | =O | H | H | =O | H | <i>Teucrium pestalozzae</i> | <i>Phytochemistry</i> , 1990, 29 , 988-989 |
| 722 | 20-oxo-teuflavin | H | =O | OAc | β OH | H | H | =O | H | | |
| 723 | teutridin | H | =O | OAc | =O | β OH | H | =O | H | <i>Teucrium trifidum</i> | <i>Phytochemistry</i> , 1994, 36 , 1549-1550 |
| 724 | 3- <i>O</i> -deacetylteugracilin A | H | β OH | OAc | β OH | H | H | =O | H | <i>Teucrium gracile</i> | <i>Phytochemistry</i> , 1991, 30 , 3693-3697 |
| 725 | teugracilin A | H | β OAc | OAc | β OH | H | H | =O | H | | |
| 726 | teugracilin B | H | β OH | OAc | α OH | H | H | =O | H | | |

| | | | | | | | | | | | |
|-----|------------------------------------|-------------|-------------|--------------|--------------|------------|----|--------------|----|-----------------------------|---|
| 727 | teusandrin A | H | H | OAc | α OAc | H | OH | =O | OH | <i>Teucrium sandrasicum</i> | <i>Phytochemistry</i> , 1997, 45 , 1653-1662 |
| 728 | teusandrin B | H | H | OAc | α OH | H | OH | =O | OH | | |
| 729 | 3-deacetylteucropodine | H | β OAc | OAc | α OH | H | H | =O | H | <i>Teucrium polium</i> | <i>Phytochemistry</i> , 1994, 37 , 1663-1666 |
| 730 | 3,20-bis-diacetylteupyreinidine | H | β OH | OAc | α OAc | H | H | α OH | H | | |
| 731 | 6,20-bis-deacetylteupyreinidine | H | β OAc | OAc | α OH | H | H | α OH | H | | |
| 732 | 3,6,20-tri-deacetyl-teupyreinidine | H | β OH | OAc | α OH | H | H | α OH | H | <i>Teucrium fruticans</i> | <i>Phytochemistry</i> , 2005, 66 , 2298-2303 |
| 733 | 10-hydroxyteucjaponin B | H | H | OAc | β OH | H | H | =O | OH | | |
| 734 | 6-acetyl-10-hydroxyteucjaponin B | H | H | OAc | β OAc | H | H | =O | OH | | |
| 735 | 6-acetylteucjaponin B | H | H | OAc | β OAc | H | H | =O | H | <i>Teucrium nudicaule</i> | <i>Nat. Prod. Res.</i> , 1996, 8 , 189-197 |
| 736 | 12- <i>epi</i> -teupyreinin | H | β OAc | OAc | α OAc | H | H | =O | H | | |
| 737 | teubutilin B | H | H | OAc | α OAc | H | H | β OAc | H | <i>Teucrium abutiloides</i> | <i>Phytochemistry</i> , 1990, 29 , 579-584 |
| 738 | teucrasiatin | H | H | OAc | =O | H | H | α OH | H | <i>Teucrium asiaticum</i> | <i>Phytochemistry</i> , 1996, 43 , 435-438 |
| 739 | teugracilin D | H | β OH | OAc | α OH | H | H | α OAc | H | <i>Teucrium gracile</i> | <i>Phytochemistry</i> , 1992, 31 , 3531-3534 |
| 740 | teucryemin | H | β OH | OH | β OAc | H | H | =O | H | <i>Teucrium yemense</i> | <i>Phytochemistry</i> , 1995, 40 , 1737-1741 |
| 741 | 19- <i>O</i> -acetylteucryemin | H | β OH | OAc | β OAc | H | H | =O | H | | |
| 742 | teucryeminone | β OAc | OAc | =O | H | β H | — | — | — | | |
| 743 | 12- <i>epi</i> -teucjaponin A | H | OAc | β OH | H | β H | — | — | — | <i>Teucrium maghrebinum</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 1029-1031 |
| 744 | sandrasin A | H | OAc | α OAc | OH | β OH | — | — | — | <i>Teucrium</i> | <i>Phytochemistry</i> , |


| | | | | | | | | | | | |
|-----|--|----------------------------|-------------|-----------------------------|--------------|------------|----|-------------|----|------------------------------------|--|
| 745 | 6-deacetylsandrasin A | H | OAc | α OH | OH | β OH | — | — | — | <i>sandrasicum</i> | 1996, 42 , 775-778 |
| 746 | teubrevin C | β OAc | OAc | =O | H | H | — | — | — | <i>Teucrium brevifolium</i> | <i>Tetrahedron</i> , 1995, 51 , 837-848 |
| 747 | teubrevin D | β OAc | OAc | =O | OH | OH | — | — | — | | |
| 748 | teupestalin A | — | — | — | — | — | — | — | — | <i>Teucrium pestalozzae</i> | <i>Phytochemistry</i> , 1990, 29 , 2229-2233 |
| 749 | teupestalin B | — | — | — | — | — | — | — | — | | |
| 750 | teubutilin A | — | — | — | — | — | — | — | — | <i>Teucrium abutiloides</i> | <i>Phytochemistry</i> , 1990, 29 , 579-584 |
| 751 | 3 α -angeloyloxy-4 β ,10 β -dihydroxy-8 β ,17,15,16-bis-epoxy- <i>cis</i> -clerodane-13(16),14-dien-20,12-olide | α OAng | — | — | — | — | — | — | — | <i>Pteronia incana</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |
| 752 | 3 α -acetoxy-4 β ,10 β -dihydroxy-8 β ,17,15,16-bis-epoxy- <i>cis</i> -clerodane-13(16),14-dien-20,12-olide | α OAc | — | — | — | — | — | — | — | | |
| 753 | 3 α ,4 β ,10 β -trihydroxy-8 β ,17,15,16-bis-epoxy- <i>cis</i> -clerodane-13(16),14-dien-20,12-olide | α OH | — | — | — | — | — | — | — | | |
| 754 | korberin A | β CO ₂ Me | α Me | α CO ₂ Me | H | =O | — | — | — | <i>Croton lechleri</i> | <i>Phytochemistry</i> , 1993, 34 , 265-268 |
| 755 | — | β Me | β Me | H | =O | =O | — | — | — | <i>Nardophyllum lanantum</i> | <i>Phytochemistry</i> , 1990, 29 , 1227-1230 |
| 756 | eluterin J | H | α Me | H | α OAc | H | H | H | =O | <i>Croton eluteria</i> | <i>J. Agric. Food Chem.</i> , 2002, 50 , 5131-5138 |
| 757 | eluterin I | H | α Me | H | α OAc | H | H | β OAc | H | | |
| 758 | 3 α ,4 α -epoxy-6-oxo-15,16-epoxy-8 β ,10 β H- <i>cis</i> -clerodane-13(16),14-dien-20,12-olide | H | β Me | =O | H | H | H | =O | H | <i>Pteronia eenii</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |
| 759 | 3 α ,4 α -epoxy-8 β -hydroxy-6-oxo-15,16-epoxy- <i>cis</i> -clerodane-13(16),14-dien-20,12-olide | H | β Me | =O | H | OH | H | =O | H | | |
| 760 | 2 α -hydroxy-3 α ,4 α ,15,16-bis-epoxy-8 β ,10 β H- <i>ent</i> -cleroda-13(16),14-diene-20,12-olide | α OH | β Me | H | H | H | H | =O | H | <i>Microglossa pyrrhopappa</i> | <i>Phytochemistry</i> , 1990, 29 , 3233-3241 |
| 761 | 6 β ,10 β -dihydroxy-3 α ,4 α ,15,16-bis-epoxy-8 β H-cleroda-13(16),14-diene-20,12-olide | H | β Me | β OH | H | H | OH | =O | H | | |
| 762 | 6 β -angeloyloxy-10 β -hydroxy-3 α ,4 α ,15,16-bis-epoxy-8 β H-cleroda-13(16),14-diene-20,12-olide | H | β Me | β OAng | H | H | OH | =O | H | | |

| | | | | | | | | | | | |
|------------|---|---|------------|--------------|---|---|----|----|---|------------------------------------|---|
| 763 | 6 β -[2-methylbutyryloxy]-10 β -hydroxy-3 α ,4 α ,15,16-bis-epoxy-8 β H-cleroda-13(16),14-diene-20,12-olide | H | β Me | β MeBu | H | H | OH | =O | H | | |
| 764 | crotonpene A | — | — | — | — | — | — | — | — | <i>Croton yanhuii</i> | <i>Fitoterapia</i> , 2014, 95 , 229-233 |
| 765 | crotonpene B | — | — | — | — | — | — | — | — | | |
| 766 | ravidin B | — | — | — | — | — | — | — | — | <i>Nannoglottis ravida</i> | <i>Phytochemistry</i> , 2004, 65 , 2533-2537 |
| 767 | 2 α -hydroxy-3 α ,4 α ,15,16-bis-epoxy-8 β H- <i>ent</i> -cleroda-1(10),13(16),14-diene-20,12-olide | — | — | — | — | — | — | — | — | <i>Microglossa pyrrhopappa</i> | <i>Phytochemistry</i> , 1990, 29 , 3233-3241 |


5.3. Type V Subtype III with a C-9-Spiro- γ -lactol Moiety (Table 23)



768,769




770



771-773

| No. | Compound Name | R ₁ | R ₂ | Source | Ref. |
|-----|--------------------|----------------|----------------|-----------------------------|---|
| 768 | heteroscypholide A | β OAc | — | <i>Heteroscyphus planus</i> | <i>Phytochemistry</i> , 1996, 41 , 581-587 |
| 769 | heteroscypholide B | β OH | — | | |
| 770 | heteroscyphone A | — | — | | <i>Phytochemistry</i> , 1995, 38 , 119-127 |
| 771 | heteroscyphone B | H | β OH | | |
| 772 | heteroscyphone C | H | H | | |
| 773 | heteroscyphone D | β OH | β OH | | |

5.4. Type V Subtype IV Other Compounds (Table 24)




| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | R ₇ | R ₈ | Source | Ref. |
|------------------|--|--------------------|--------------------|--------------------|---------------------|----------------|----------------|------------------|----------------|-----------------------------|---|
| 774 | eluterin H | Me | Me | αOAc | βOAc | — | — | — | — | <i>Croton eluteria</i> | <i>J. Agric. Food Chem.</i> , 2002, 50 , 5131-5138 |
| 775 | crotocorylifuran | CO ₂ Me | CO ₂ Me | H | =O | — | — | — | — | <i>Croton haumanianus</i> | <i>Tetrahedron</i> , 1990, 46 , 5199-5202 |
| 776 ^a | teupernin D | CO ₂ Me | H | — | — | — | — | — | — | <i>Teucrium pernyi</i> | <i>Phytochemistry</i> , 1993, 33 , 716-717 |
| 777 | teulolin A | CH ₂ OH | βOH | — | — | — | — | — | — | <i>Teucrium polium</i> | <i>Phytochemistry</i> , 1999, 51 , 921-925 |
| 778 | teulolin B | — | — | — | — | — | — | — | — | | |
| 779 | 3α-angeloyloxy-2α,4β-dihydroxy-15,16-epoxy- <i>cis</i> -clerodane-1(10),13(16),14-dien-20,12-olide | — | — | — | — | — | — | — | — | <i>Pteronia eenii</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |
| 780 | teupolin VI | — | — | — | — | — | — | — | — | <i>Teucrium polium</i> | <i>Phytochemistry</i> , 2011, 72 , 2037-2044 |
| 781 | sandrasin B | H | OH | CH ₂ OH | αCH ₂ OH | =O | H | αMe | αOH | <i>Teucrium sandrasicum</i> | <i>Phytochemistry</i> , 1996, 42 , 775-778 |
| 782 | 3α,4β,8β,10β-tetrahydroxy-15,16-epoxy- <i>cis</i> -clerodane-13(16),14-dien-20,12-olide | αOH | Me | OH | βMe | H | H | αMe | βOH | <i>Pteronia eenii</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |
| 783 | 3α-acetoxy-4β,8β,10β-trihydroxy-15,16-epoxy- <i>cis</i> -clerodane-13(16),14-dien-20,12-olide | αOAc | Me | OH | βMe | H | H | αMe | βOH | | |
| 784 | 3-oxo-4β,8β,10β-trihydroxy-15,16-epoxy- <i>cis</i> -clerodane-13(16),14-dien-20,12-olide | =O | Me | OH | βMe | H | H | αMe | βOH | | |
| 785 | 3α,4β,10β-trihydroxy-15,16-epoxy- <i>cis</i> -clerodane-8(17),13(16),14-dien-20,12-olide | αOH | Me | OH | βMe | H | H | =CH ₂ | βOH | | |

| | | | | | | | | | | | |
|-----|--|---------------|-----------------------------|-----------------------------|-----------------------------|---|---|------------------|------------|--------------------------------------|--|
| 786 | 3-oxo-4 β ,10 β -dihydroxy-15,16-epoxy- <i>cis</i> -clerodane-8(17),13(16),14-dien-20,12-olide | =O | Me | OH | β Me | H | H | =CH ₂ | β OH | | |
| 787 | 3 α ,4 β ,10 β -trihydroxy-15,16-epoxy- <i>cis</i> -clerodane-13(16),14-dien-20,12-olide | α OH | Me | OH | β Me | H | H | α Me | β OH | <i>Pteronia</i> <i>incana</i> | |
| 788 | 3 α -angeloyloxy-4 β ,10 β -dihydroxy-15,16-epoxy- <i>cis</i> -clerodane-8(17),13(16),14-dien-20,12-olide | α OAng | Me | OH | β Me | H | H | =CH ₂ | β OH | | |
| 789 | korberin B | H | β CO ₂ Me | α Me | α CO ₂ Me | — | — | — | — | <i>Croton</i> <i>lechleri</i> | <i>Phytochemistry</i> , 1993, 34 , 265-268 |
| 790 | teuctomentin | β OAc | α CH ₂ OH | α CH ₂ OH | β OH | — | — | — | — | <i>Teucrium</i> <i>tomentosum</i> | <i>Der Pharmacia</i> <i>Lette</i> , 2014, 6 , 295-298 |
| 791 | musabalbisiane A | OH | CO ₂ H | CHO | CHO | — | — | — | — | <i>Musa</i> <i>balbisiana</i> | <i>Phytochemistry</i> , 1992, 31 , 2173-2175 |
| 792 | musabalbisiane B | OH | CHO | CO ₂ H | CH ₂ OH | — | — | — | — | | |
| 793 | musabalbisiane C | OAng | CH ₂ OH | CH ₂ OH | CH ₂ OH | — | — | — | — | | |

^a Compound **1025** in Table 29 was also given the name 'teupernin D'.

6. Type VI with a Furofuran-based Side Chain at C-9

6.1. Type VI Subtype I with a Hexahydrofurofuran-based Side Chain at C-9 (Table 25)




| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | Source | Ref. |
|-----|---------------|-------------------------|----------------|----------------|----------------|----------------|-------------------------|---|
| 794 | lupulin F | H | α OH | OPr | OAc | — | <i>Ajuga lupulina</i> | <i>Indian J. Chem.</i> , 1999, 38B , 743-745 |
| 795 | ajubractin C | H | H | MeBuO | OAc | — | <i>Ajuga bracteosa</i> | <i>J. Nat. Prod.</i> , 2011, 74 , 1036-1041 |
| 796 | ajubractin D | H | α OH | OiBu | OAc | — | | |
| 797 | ajubractin E | H | H | OH | OAc | — | | |
| 798 | areptin A | β OH | α OAc | Y ₁ | OAc | — | <i>Ajuga reptans</i> | <i>Phytochemistry</i> , 1998, 49 , 2443-2447 |
| 799 | ajugavensin A | β OY ₁ | H | H | OAc | — | <i>Ajuga genevensis</i> | <i>Phytochemistry</i> , 1991, 30 , 4083-4085 |

| | | | | | | | | |
|-----|---|---------------|----------------|------------------------|------|--------------|---|---|
| 800 | ajugavensin B | α OTig | H | H | OAc | — | | |
| 801 | ajugavensin C | β OH | H | H | OTig | — | | |
| 802 | 3 β -hydroxyajugavensin B | α OTig | H | OH | OAc | — | <i>Ajuga reptans</i> | <i>Phytochemistry</i> , 1998, 47 , 1227-1232 |
| 803 | ajugorientin | β OTig | H | OH | OAc | — | <i>Ajuga orientalis</i> | <i>Phytochemistry</i> , 1997, 45 , 121-123 |
| 804 | ajugapyrin A | β OH | H | β OTig | H | — | <i>Ajuga pyramidalis</i> | <i>Phytochemistry</i> , 1998, 47 , 303-305 |
| 805 | 14,15-dihydrojodrellin T | β OTig | H | OAc | H | — | <i>Scutellaria</i> | <i>Phytochemistry</i> , 1990, 29 , 1793-1796 |
| 806 | scutegalin A | H | H | OTig | OTig | — | <i>galericulata</i> | <i>Phytochemistry</i> , 1993, 33 , 309-315 |
| 807 | scutecyprin | H | H | β OTig | H | — | <i>Scutellaria cypria</i> var. <i>elatior</i> | <i>Phytochemistry</i> , 1993, 33 , 931-932 |
| 808 | scutecolumnin B | H | H | β Y ₁ | H | — | <i>Scutellaria columnae</i> | <i>Phytochemistry</i> , 1992, 31 , 3639-3641 |
| 809 | scutecolumnin C | H | H | β OH | H | — | | |
| 810 | scutecyprol B | H | H | β OTig | H | OH | <i>Scutellaria cypria</i> var. <i>cypria</i> | <i>Phytochemistry</i> , 1996, 42 , 555-557 |
| 811 | scupolin K | H | β OH | OiBu | H | OH | <i>Scutellaria polyodon</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 1032-1034 |
| 812 | 6 α - <i>O</i> -acetyl-15 β ,19 β -di- <i>O</i> -ethyl-2,19:4,18:11,16:15,16-tetraepoxyneoclerodane-6,15,19-triol | H | H | OEt | H | β OEt | <i>Scutellaria discolor</i> | <i>Chem. Pharm. Bull.</i> , 1996, 44 , 1540-1545 |
| 813 | 6 α - <i>O</i> -acetyl-15 α ,19 β -di- <i>O</i> -ethyl-2,19:4,18:11,16:15,16-tetraepoxyneoclerodane-6,15,19-triol | H | H | OEt | H | α OEt | | |
| 814 | 6 α - <i>O</i> -acetyl-19 β - <i>O</i> -ethyl-2,19:4,18:11,16:15,16-tetraepoxyneoclerodane-6,15,19-triol | H | H | OEt | H | OH | | |
| 815 | 6 α ,19-di- <i>O</i> -acetyl-2,19:4,18:11,16:15,16-tetraepoxy-neoclerodane-6,15,19-triol | H | H | OAc | H | OH | | |
| 816 | scutalbin B | H | H | β Y ₁ | H | OH | <i>Scutellaria albida</i> | <i>Phytochemistry</i> , 1996, 42 , 1059-1064 |
| 817 | scutalbin C | H | H | β OH | H | OH | | |
| 818 | scutalsin | H | H | β OiBu | H | OH | | |
| 819 | 11-episcutecyprin | H | H | β OTig | H | H | <i>Scutellaria columnae</i> | <i>Phytochemistry</i> , 1997, 46 , 955-958 |
| 820 | 11-epi-scutecolumnin C | H | H | β OH | H | H | <i>Scutellaria columnae</i> var. <i>columnae</i> | <i>Phytochemistry</i> , 1998, 49 , 811-815 |
| 821 | 15-epi-lupulin B | α OH | MeBuO | OAc | OAc | α OMe | <i>Ajuga bracteosa</i> | <i>J. Nat. Prod.</i> , 2011, 74 , 1036-1041 |
| 822 | lupulin A | α OH | Y ₁ | OAc | OAc | β OMe | <i>Ajuga lupulina</i> | <i>J. Nat. Prod.</i> , 1996, 59 , 668-670 |
| 823 | lupulin B | H | Y ₁ | OAc | OAc | α OMe | | |
| 824 | (15 <i>S</i>)-14,15-dihydro-15-hydroxyajugachin A | α OH | OiBu | OAc | OAc | β OH | <i>Ajuga salicifolia</i> | <i>Phytochemistry</i> , 1993, 34 , 1173-1175 |
| 825 | (15 <i>R</i>)-14,15-dihydro-15-hydroxyajugachin A | α OH | OiBu | OAc | OAc | α OH | | |
| 826 | 14,15-dihydro-15-oxoajugachin A | α OH | OiBu | OAc | OAc | =O | | |

| | | | | | | | | |
|-----|---|-------------|-----------------|------|-----|--------------|---|---|
| 827 | hativene A | α OH | OiBu | OAc | OAc | β OMe | <i>Ajuga pseudoiva</i> | <i>Fitoterapia</i> , 2000, 71 , 105-112 |
| 828 | hativene B | α OH | OiBu | OAc | OAc | α OMe | | |
| 829 | hativene C | β OH | OiBu | OAc | OAc | α OMe | | |
| 830 | 15-methoxy-14,15-dihydro-3-epicaryoptin | H | OAc | OAc | OAc | β OMe | <i>Clerodendrum inerme</i> | <i>Phytochemistry</i> , 1992, 31 , 338-340 |
| 831 | 14,15-dihydro-15 β -methoxy-3-epicaryoptin | H | OAc | OAc | OAc | OH | | <i>Phytochemistry</i> , 2005, 66 , 643-648 |
| 832 | 15 β -ethoxy-14-hydroclerodin | H | H | OAc | OAc | β OEt | <i>Scutellaria discolor</i> | <i>Chem. Pharm. Bull.</i> , 1996, 44 , 1540-1545 |
| 833 | 15 α -ethoxy-14-hydroclerodin | H | H | OAc | OAc | α OEt | | |
| 834 | 14-hydro-15-hydroxy-6- <i>O</i> -deacetylclerodin | H | H | OAc | OH | OH | | |
| 835 | scutelaterin C | Y1 | H | OAc | OAc | OH | <i>Scutellaria lateriflora</i> | <i>Phytochemistry</i> , 1998, 48 , 687-691 |
| 836 | scutellin A | H | H | OPr | OAc | β OMe | <i>Scutellaria barbata</i> | <i>Yunnan Zhiwu Yanjiu</i> , 2009, 31 , 474-476 |
| 837 | scutalpin O | H | H | OiBu | OAc | OH | <i>Scutellaria alpina</i> | <i>Phytochemistry</i> , 1998, 49 , 2449-2452 |
| 838 | scutecyprol A | H | H | OAc | OAc | OH | <i>Scutellaria cypria</i> var. <i>cypria</i> | <i>Phytochemistry</i> , 1996, 42 , 555-557 |
| 839 | scupontin C | H | OAc | — | — | — | <i>Scutellaria pontica</i> | <i>J. Nat. Prod.</i> , 1997, 60 , 348-355 |
| 840 | scupontin D | OH | OAc | — | — | — | | |
| 841 | scupontin F | OH | OX ₆ | — | — | — | | |
| 842 | scutalpin M | — | — | — | — | — | <i>Scutellaria alpina</i> | <i>Phytochemistry</i> , 1995, 38 , 181-187 |
| 843 | lupulin C | — | — | — | — | — | <i>Ajuga lupulina</i> | <i>J. Nat. Prod.</i> , 1996, 59 , 668-670 |
| 844 | inerme A | H | H | — | — | — | <i>Clerodendrum inerme</i> | <i>Phytochemistry</i> , 2005, 66 , 643-648 |
| 845 | inerme B | OMe/H | H/OMe | — | — | — | | |


6.2. Type VI Subtype II with a Tetrahydrofurofuran-based Side Chain at C-9 (Table 26)



| No. | Compound Name | R ₁ | R ₂ | R ₃ | Source | Ref. |
|-----|-----------------------------|-----------------|-----------------|-----------------|---------------------------------|---|
| 846 | lupulin E | H | αOH | OPr | <i>Ajuga lupulina</i> | <i>Indian J. Chem.</i> , 1999, 38B , 743-745 |
| 847 | ajugachin A | H | αOH | OiBu | <i>Ajuga chamaepitys</i> | <i>Phytochemistry</i> , 1990, 29 , 2931-2933 |
| 848 | ajugachin B | H | αOH | Y ₁₁ | | |
| 849 | clerodendrin B | H | αOH | Y ₃ | <i>Clerodendrum inerme</i> | <i>Phytochemistry</i> , 1993, 34 , 572-574 |
| 850 | clerodendrin C | H | αOH | Y ₁₀ | | |
| 851 | clerodendrin H | H | αOH | Y ₇ | <i>Clerodendrum trichotomum</i> | <i>Phytochemistry</i> , 1998, 49 , 1975-1980 |
| 852 | galericulin | H | αOH | OTig | <i>Scutellaria galericulata</i> | <i>Phytochemistry</i> , 1990, 29 , 1793-1796 |
| 853 | scutelaterin A | H | βOAc | H | <i>Scutellaria laterifora</i> | <i>Phytochemistry</i> , 1998, 48 , 687-691 |
| 854 | scutelaterin B | H | βY ₁ | H | | |
| 855 | ajubractin A | H | H | MeBuO | <i>Ajuga bracteosa</i> | <i>J. Nat. Prod.</i> , 2011, 74 , 1036-1041 |
| 856 | ajubractin B | H | H | iBuO | | |
| 857 | ajugapitin | H | αOH | MeBuO | | |
| 858 | areptin B | βOTig | H | OH | <i>Ajuga reptans</i> | <i>Phytochemistry</i> , 1998, 49 , 2443-2447 |
| 859 | 14,15-dehydroajugareptansin | βY ₁ | H | OH | | <i>Phytochemistry</i> , 1998, 47 , 1227-1232 |
| 860 | jodrellin A | H | H | βOAc | <i>Scutellaria</i> spp. | <i>Phytochemistry</i> , 1991, 30 , 1125-1127 |
| 861 | jodrellin B | H | H | βOiPr | | |
| 862 | scupolin J | H | βOH | OiBu | <i>Scutellaria polyodon</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 1032-1034 |
| 863 | jodrellin T | βOTig | H | βOAc | <i>Scutellaria galericulata</i> | <i>Phytochemistry</i> , 1990, 29 , 1793-1796 |
| 864 | 19-O-deacetyljodrellin A | H | H | OH | <i>Scutellaria discolor</i> | <i>Chem. Pharm. Bull.</i> , 1996, 44 , 1540-1545 |

| | | | | | | |
|------------|-----------------|-----------------|----------------|------------------------|---------------------------------|---|
| 865 | scutegrossin A | H | H | β OTig | <i>Scutellaria grossa</i> | <i>Chem. Pharm. Bull.</i> , 1997, 45 , 1097-1100 |
| 866 | scutalbin A | H | H | β OH | <i>Scutellaria albida</i> | <i>Phytochemistry</i> , 1996, 42 , 1059-1064 |
| 867 | scutecolumnin A | H | H | β Y ₁ | <i>Scutellaria columnae</i> | <i>Phytochemistry</i> , 1992, 31 , 3639-3641 |
| 868 | clerodendrin I | Y ₆ | | — | <i>Clerodendrum trichotomum</i> | <i>Biosci. Biotechnol. Biochem.</i> , 1999, 63 , 1795-1797 |
| 869 | clerodendrin E | Y ₁₀ | | — | | <i>Phytochemistry</i> , 1998, 49 , 1975-1980 |
| 870 | clerodendrin F | Y ₇ | | — | | |
| 871 | clerodendrin G | Y ₁ | | — | | |
| 872 | scupontin A | OH | OAc | — | <i>Scutellaria pontica</i> | <i>J. Nat. Prod.</i> , 1997, 60 , 348-355 |
| 873 | scupontin B | H | OAc | — | | |
| 874 | scupontin E | OH | X ₆ | — | | |


7 Type VII with a 13-Spiro-15,16- γ -lactone Moiety (Table 27)






| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | R ₆ | Source | Ref. |
|-----|---------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|---|---|
| 875 | scutorientalin A | OiBu | H | H | OAc | | | <i>Scutellaria orientalis</i> subsp. <i>pinnatifida</i> | <i>Phytochemistry</i> , 1996, 43 , 173-178 |
| 876 | scutorientalin C | OiBu | H | α OH | OAc | | | | <i>Phytochemistry</i> , 1997, 44 , 121-124 |
| 877 | scutorientalin D | OiBu | H | β OAc | OAc | | | | |
| 878 | scutalpin D | OTig | H | β OAc | OAc | | | <i>Scutellaria alpina</i> subsp. <i>javallambrensis</i> | <i>Phytochemistry</i> , 1993, 34 , 1589-1594 |
| 879 | 11-deacetylscutalpin D | OTig | H | β OH | OAc | | | | <i>Phytochemistry</i> , 1997, 44 , 593-597 |
| 880 | scutalpin A | Y ₁ | β OAc | H | OAc | | | <i>Scutellaria alpina</i> | <i>Phytochemistry</i> , 1993, 34 , 453-456 |
| 881 | scutalpin F | OAc | β OAc | H | OAc | | | | <i>Phytochemistry</i> , 1994, 35 , 1285-1288 |
| 882 | scutalpin G | OBz | H | β OAc | OAc | | | | <i>Phytochemistry</i> , 1995, 38 , 181-187 |
| 883 | hastifolin C | OCin | H | H | H | | | <i>Scutellaria hastifolia</i> | <i>Phytochemistry</i> , 2010, 71 , 2087-2091 |
| 884 | scuterivulactone C ₂ | OH | Me | OH | OBz | OAc | | <i>Scutellaria rivularis</i> | <i>Chem. Pharm. Bull.</i> , 1997, 45 , 152-160 |

| | | | | | | | | | |
|-----|---------------------------------|------|------|--------------------|------|-----|------|--|---|
| 885 | hastifolin E | H | OH | CH ₂ OH | OCin | H | | <i>Scutellaria hastifolia</i> | <i>Phytochemistry</i> , 2010, 71 , 2087-2091 |
| 886 | hastifolin G | OCin | H | H | | | | | |
| 887 | barbatin A | H | OBz | OH | OBz | | | <i>Scutellaria barbata</i> | <i>Phytochemistry</i> , 2006, 67 , 1326-1330 |
| 888 | barbatellarine E | | | | | | | | <i>Helv. Chim. Acta.</i> , 2011, 94 , 643-649 |
| 889 | barbatin B | OBz | OBz | OH | | | | | <i>Phytochemistry</i> , 2006, 67 , 1326-1330 |
| 890 | scutebata D | OBz | OAc | OAc | H | | | | <i>J. Nat. Prod.</i> , 2010, 73 , 233-236 |
| 891 | scutebata E | OiBu | OAc | OAc | H | | | | <i>Chem. Nat. Compds.</i> , 2014, 50 , 256-257 |
| 892 | barbatellarine F | | | | | | | | |
| 893 | scuteselerin | OAc | OH | Y ₁₂ | OAc | | | <i>Scutellaria seleriana</i> | <i>Phytochemistry</i> , 1998, 47 , 135-137 |
| 894 | hastifolin F | H | OCin | H | H | | | <i>Scutellaria hastifolia</i> | <i>Phytochemistry</i> , 2010, 71 , 2087-2091 |
| 895 | hastifolin B | H | H | OCin | H | H | H | | |
| 896 | scutenisin | H | OH | OiBu | OiBu | H | H | <i>Scutellaria orientalis</i> subsp. <i>sintenisii</i> | <i>Phytochemistry</i> , 1998, 49 , 1825-1827 |
| 897 | scutalpin E | H | OAc | OTig | OAc | H | H | <i>Scutellaria alpina</i> | <i>Phytochemistry</i> , 1994, 35 , 1285-1288 |
| 898 | scuteguatemalin | OiBu | H | OAc | H | OAc | OiBu | <i>Scutellaria guatemalensis</i> | <i>Heterocycles</i> , 1997, 45 , 2247-2252 |
| 899 | hastifolin D | H | OH | CH ₂ OH | OCin | H | | <i>Scutellaria hastifolia</i> | <i>Phytochemistry</i> , 2010, 71 , 2087-2091 |
| 900 | scuterivulactone C ₁ | OH | Me | OH | OBz | OAc | | <i>Scutellaria rivularis</i> | <i>Chem. Pharm. Bull.</i> , 1997, 45 , 152-160 |
| 901 | scuterivulactone B | =O | Me | H | OBz | OAc | | | |
| 902 | barbatellarine A | | | | | | | <i>Scutellaria barbata</i> | <i>Bioorg. Med. Chem. Lett.</i> , 2010, 20 , 288-290 |

8. Clerodane Diterpene Glycosides (Table 28)





| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | Source | Ref. |
|-----|---|----------------|----------------|----------------|----------------|----------------------------|---------------------------------------|
| 903 | α -vinyl-1,2,3,4,4a,7,8,8a-octahydro- α ,1,2,4a,5-pentamethyl-1-naphthalenepropanol-4-O- β -glucopyranoside | β -Glc | OH | Me | — | <i>Gleichenia japonica</i> | <i>Chem. Lett.</i> , 1991, 4, 701-704 |

| | | | | | | | |
|-----|---|--|---|-------------|---|----------------------------------|---|
| 904 | α -vinyl-1,2,3,4,4a,7,8,8a-octahydro- α ,1,2,4a,5-pentamethyl-1-naphthalenepropanol-4- <i>O</i> - α -rhamnopyranosyl-(1 \rightarrow 2)- β -glucopyranoside | β -Glc-(2 \rightarrow 1)- α -L-Rha | OH | Me | — | | |
| 905 | (6 <i>S</i> ,13 <i>S</i>)-6- <i>O</i> -[β -D-glucopyranosyl-(1 \rightarrow 4)- α -L-rhamnopyranosyl]-13- <i>O</i> -[α -L-rhamnopyranosyl-(1 \rightarrow 4)- β -D-fucopyranosyl]-cleroda-3,14-diene | α -L-Rha-(4 \rightarrow 1)- β -D-Glc | β -{ β -D-Fuc-(4 \rightarrow 1)- α -L-Rha} | α Me | — | <i>Dicranopteris pedata</i> | <i>Phytochemistry</i> , 1997, 46 , 839-844 |
| 906 | (6 <i>S</i> ,13 <i>S</i>)-cleroda-3,14-diene-6,13-diol-6- <i>O</i> - β -glucopyranosyl-13- <i>O</i> - β -fucopyranosyl-(1 \rightarrow 2)- α -rhamnopyranoside | β -Glc | β -{ α -Rha-(2 \rightarrow 1)- β -Fuc} | α Me | — | | |
| 907 | (5 <i>S</i> ,8 <i>R</i> ,9 <i>S</i> ,10 <i>R</i> ,13 <i>S</i>)-10 β H-13- <i>O</i> - α -L-arabinopyranosyl-2-oxo-17 α ,19 β ,20 α -trimethyl-3,14-clerodadiene | =O | — | — | — | | |
| 908 | (5 <i>S</i> ,8 <i>R</i> ,9 <i>S</i> ,10 <i>R</i> ,13 <i>S</i>)-10 β H-13- <i>O</i> - α -L-arabinopyranosyl-17 α ,19 β ,20 α -trimethyl-3,14-clerodadiene | H | — | — | — | <i>Nannoglottis carpesioides</i> | <i>Fitoterapia</i> , 2014, 93 , 39-46 |
| 909 | (5 <i>S</i> ,8 <i>R</i> ,9 <i>S</i> ,10 <i>R</i> ,13 <i>S</i>)-10 β H-13- <i>O</i> - α -L-arabinopyranosyl-2 β -hydroperoxyl-17 α ,19 β ,20 α -trimethyl-3,14-clerodadiene | OOH | — | — | — | | |
| 910 | (6 <i>S</i> ,13 <i>S</i>)-6- <i>O</i> -[6- <i>O</i> -acetyl- β -D-glucopyranosyl-(1 \rightarrow 4)- α -L-rhamnopyranosyl]cleroda-3,14-dien-13-ol | 6- <i>O</i> -acetyl- β -D-Glc-(1 \rightarrow 4)- α -L-Rha | — | — | — | <i>Dicranopteris dichotoma</i> | <i>J. Nat. Prod.</i> , 2007, 70 , 265-268 |
| 911 | (6 <i>S</i> ,13 <i>S</i>)-6- <i>O</i> -[4- <i>O</i> -acetyl- β -D-glucopyranosyl-(1 \rightarrow 4)- α -L-rhamnopyranosyl]cleroda-3,14-dien-13-ol | 4- <i>O</i> -acetyl- β -D-Glc-(1 \rightarrow 4)- α -L-Rha | — | — | — | | |
| 912 | 6- <i>O</i> -[6- <i>O</i> -acetyl- β -D-glucopyranosyl-(1-4)- α -L-rhamnopyranosyl]-(13 <i>E</i>)-cleroda-3,13-dien-15-ol | 6- <i>O</i> -acetyl- β -D-Glc-(1 \rightarrow 4)- α -L-Rha | — | — | — | | |

| | | | | | | | |
|-----|--|-----------------------------|-----------------------------|---|---|--|---|
| 913 | 6- <i>O</i> -[β-D-glucopyranosyl]-(1→4)-α-L-rhamnopyranosyl-(13 <i>E</i>)-cleroda-3,13-dien-15-ol | β-Glc-(1→4)-α-L-Rha | — | — | — | | |
| 914 | cussoside A | Glc | H | — | — | <i>Cussonia racemosa</i> | <i>Phytochemistry</i> , 2002, 60 , 339-343 |
| 915 | cussoside B | Glc | Glc | — | — | | |
| 916 | cussoside C | Glc | Api | — | — | | |
| 917 | cussoside D | Glc-(6→1)-Api | H | — | — | | |
| 918 | porwenin B | — | — | — | — | <i>Portulaca okinawensis</i> | <i>J. Nat. Prod.</i> , 2001, 64 , 804-805 |
| 919 | pilosanol C | — | — | — | — | <i>Portulaca pilosa</i> | <i>Phytochemistry</i> , 1991, 30 , 4075-4077 |
| 920 | salvigreside A | Me | β-D-Glc | — | — | <i>Salvia greggii</i> | <i>Phytochemistry</i> , 2004, 65 , 2577-2581 |
| 921 | salvigreside B | Me | 6- <i>O</i> -acetyl-β-D-Glc | — | — | | |
| 922 | salvigreside C | CH ₂ OH | 6- <i>O</i> -acetyl-β-D-Glc | — | — | | |
| 923 | salvigreside D | 6- <i>O</i> -acetyl-β-D-Glc | — | — | — | | |
| 924 | 6-hydroxy(-)-hardwickiic acid 2'-β-D-glucopyranosyl benzyl ester | H | — | — | — | <i>Elsholtzia bodinieri</i> | <i>Indian J. Chem.</i> , 2008, 47B , 166-170 |
| 925 | 6,7-dihydroxy(-)-hardwickiic acid 2'-β-D-glucopyranosyl benzyl ester | OH | — | — | — | | |
| 926 | amarisolide | — | — | — | — | <i>Salvia amarissima</i> | <i>Phytochemistry</i> , 1996, 42 , 1105-1108 |
| 927 | borapetoside D | H | Glc-(6→1)-Glc | — | — | <i>Tinospora tuberculata</i> | <i>Liebigs Ann. Chem.</i> , 1993, 491-495 |
| 928 | borapetoside E | H | Glc | — | — | | |
| 929 | rumphioside I | αOH | Glc | — | — | <i>Tinospora rumphii</i> | <i>Phytochemistry</i> , 1996, 42 , 153-158 |
| 930 | <i>cis</i> -cleroda-15,16-dihydroxy-3,13(<i>Z</i>)-dien-18- <i>O</i> -[β-D-galactopyranosyl]-peracetyylester | — | — | — | — | <i>Baccharis sagittalis</i> | <i>Phytochemistry</i> , 2002, 61 , 899-905 |
| 931 | <i>cis</i> -cleroda-3,13(14)-dien-15,16-olide-18- <i>O</i> -[β-D-galactopyranosyl]-peracetyylester | — | — | — | — | | |
| 932 | sagittatayunnanoside B | Glc-(6→1)-Glc | | | — | <i>Tinospora sagittata</i> var. <i>yunnanensis</i> | <i>Planta Med.</i> , 2014, 80 , 419-425 |
| 933 | sagittatayunnanoside A | OH | CH ₂ OGlc | | | | |


| | | | | | | | |
|-----|---|----------------------|----------------------|----------------------------|----|-----------------------------|---|
| 934 | sagittatayunnanoside C | OGlc | CH ₂ OGlc | | | | |
| 935 | sagittatayunnanoside D | OH | CO ₂ Glc | — | — | | |
| 936 | rumphioside A | β-D-Glc | H | =O | — | <i>Tinospora rumphii</i> | <i>Phytochemistry</i> , 1995, 40 , 1729-1736 |
| 937 | rumphioside B | β-D-Glc | Me | =O | — | | |
| 938 | rumphioside Ac-D | Glc(Ac) ₄ | Me | OMe | — | | |
| 939 | rumphioside C | β-D-Glc | — | — | — | | |
| 940 | rumphioside C-1 | β-D-Glc | — | — | — | | |
| 941 | rumphioside E | — | — | — | — | | |
| 942 | rumphioside F | — | — | — | — | | |
| 943 | cordifolide B | — | — | — | — | <i>Tinospora cordifolia</i> | <i>Org. Lett.</i> , 2012, 14 , 2118-2121 |
| 944 | cordifolide C | — | — | — | — | | |
| 945 | fibrauretinioside A | αOGlc | — | — | — | <i>Fibraurea tinctoria</i> | <i>J. Nat. Prod.</i> , 2007, 70 , 1930-1933 |
| 946 | <i>epi</i> -fibrauretinioside A | βOGlc | — | — | — | | |
| 947 | borapetoside A | — | — | — | — | <i>Tinospora crispa</i> | <i>J. Nat. Prod.</i> , 2010, 73 , 541-547 |
| 948 | cordifolide A | — | — | — | — | <i>Tinospora cordifolia</i> | <i>Org. Lett.</i> , 2012, 14 , 2118-2121 |
| 949 | cordioside | — | — | — | — | | <i>Phytochemistry</i> , 1995, 38 , 447-449 |
| 950 | (5 <i>R</i> ,6 <i>R</i> ,8 <i>S</i> ,9 <i>R</i> ,10 <i>S</i> ,12 <i>S</i>)-15,16-epoxy-2-oxo-6- <i>O</i> -(β-D-glucopyranosyl)-cleroda-3,13(16),14-trien-17,12-olid-18-oic acid methyl ester | — | — | — | — | | |
| 951 | (2 <i>R</i> ,5 <i>R</i> ,6 <i>R</i> ,8 <i>S</i> ,9 <i>S</i> ,10 <i>S</i> ,12 <i>S</i>)-15,16-epoxy-2-hydroxy-6- <i>O</i> -{β-D-glucopyranosyl-(1→6)-α-D-xylopyranosyl}-cleroda-3,13(16),14-trien-17,12-olid-18-oic acid methyl ester | αOH | Me | β- {β-D-Glc-(1→6)-α-D-Xyl} | αH | <i>Tinospora crispa</i> | <i>J. Nat. Prod.</i> , 2010, 73 , 541-547 |
| 952 | (2 <i>R</i> ,5 <i>R</i> ,6 <i>R</i> ,8 <i>R</i> ,9 <i>S</i> ,10 <i>S</i> ,12 <i>S</i>)-15,16-epoxy-2-hydroxy-6- <i>O</i> -(β-D-glucopyranosyl)-cleroda-3,13(16),14-trien-17,12-olid-18-oic acid methyl ester | αOH | Me | β-(β-D-Glc) | βH | | |


| | | | | | | | |
|-----|--|-----------|----------------|-------------|-----|------------------------------|---|
| 953 | (5R,6R,8S,9R,10R,12S)-15,16-epoxy-2-oxo-6-O-(β-D-glucopyranosyl)-cleroda-3,13(16),14-trien-17,12-olid-18-oic acid methyl ester | =O | Me | β-(β-D-Glc) | αH | | |
| 954 | epi-tinophyllololide | OGlc | Me | H | αH | <i>Tinospora capillipes</i> | <i>Chin. Chem. Lett.</i> , 1992, 3 , 185-188 |
| 955 | tinospinoside A | OGlc | Me | H | βH | <i>Tinospora sagittata</i> | <i>Planta Med.</i> , 2012, 78 , 82-85 |
| 956 | tinospinoside B | OGlc | Me | H | βOH | | |
| 957 | tinospinoside C | OGlc | Me | H | αOH | | |
| 958 | floribundic acid glucoside | H | Glc | H | αH | <i>Penianthus zenkeri</i> | <i>Phytochemistry</i> , 1991, 30 , 1957-1962 |
| 959 | zenkerin | H | -Xyl-(1→6)-Glc | H | αH | | |
| 960 | borapetoside C | H | Me | Glc | H | <i>Tinospora tuberculata</i> | <i>Liebigs Ann. Chem.</i> , 1993, 491-495 |
| 961 | borapetoside G | =O | Me | Glc | H | | |
| 962 | borapetoside F | H | — | — | — | | |
| 963 | tinoscorside C | αOH | — | — | — | <i>Tinospora cordifolia</i> | <i>Fitoterapia</i> , 2010, 81 , 485-489 |
| 964 | (2R,5R,6S,9S,10S,12S)-15,16-epoxy-2-hydroxy-6-O-(β-D-glucopyranosyl)-cleroda-3,7,13(16),14-tetraen-17,12-olid-18-oic acid methyl ester | αOH | — | — | — | <i>Tinospora crispa</i> | <i>J. Nat. Prod.</i> , 2010, 73 , 541-547 |
| 965 | (5R,6S,9S,10S,12S)-15,16-epoxy-2-oxo-6-O-(β-D-glucopyranosyl)-cleroda-3,7,13(16),14-tetraen-17,12-olid-18-oic acid methyl ester | =O | — | — | — | | |
| 966 | (3R,4R,5R,6S,8R,9S,10S,12S)-15,16-epoxy-3,4-epoxy-6-O-(β-D-glucopyranosyl)-cleroda-3,13(16),14-trien-17,12-olid-18-oic acid methyl ester | — | — | — | — | | |
| 967 | (1R,4S,5R,8S,9R,10S,12S)-15,16-epoxy-4-O-(β-D-glucopyranosyl)-cleroda-2,13(16),14-triene-17(12),18(1)-diolide | — | — | — | — | | |
| 968 | tinospinoside D | — | — | — | — | <i>Tinospora sagittata</i> | <i>Chem. Pharm. Bull.</i> , 2012, 60 , 1324-1328 |
| 969 | tinospinoside E | — | — | — | — | | |
| 970 | tinospinosinenside A | — | — | — | — | <i>Tinospora sinensis</i> | <i>J. Nat. Prod.</i> , 2007, 70 , 1971-1976 |
| 971 | tinospinosinenside B | OAc | — | — | — | | |
| 972 | tinospinosinenside C | OH | — | — | — | | |
| 973 | epi-12-palmatoside G | — | — | — | — | <i>Fibraurea tinctoria</i> | <i>J. Nat. Prod.</i> , 2007, 70 , 1930-1933 |
| 974 | chamaedryoside A | OH | Glc | — | — | <i>Teucrium chamaedrys</i> | <i>Magn. Reson. Chem.</i> , 2009, 47 , 1007-1012 |
| 975 | chamaedryoside B | O-6-α-Glc | H | — | — | | |

| | | | | | | | |
|------------|-------------------------------|-------------------------|---|---|---|---|---|
| 976 | chamaedryoside C | O-6-β-Glc | H | — | — | | |
| 977 | teulamioside | — | — | — | — | <i>Teucrium lamiiifolium</i> | <i>Phytochemistry</i> , 1993, 34 , 1095-1098 |
| 978 | syphonoside | H | — | — | — | <i>Syphonota geographica</i> , <i>Halophila stipulacea</i> | <i>J. Org. Chem.</i> , 2007, 72 , 5625-5630 |
| 979 | 6''-acetyl syphonoside | Ac | — | — | — | | <i>Tetrahedron</i> , 2008, 64 , 191-196 |
| 980 | syphonosideol | H | — | — | — | | |
| 981 | mixture of syphonoside esters | palmitic & stearic acid | — | — | — | | |

9. Clerodane Derivatives

9.1. N (or S or Cl)-Containing Derivatives (Table 29)





| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | R ₅ | Source | Ref. |
|------|---|----------------|---------------------------------|-----------------|----------------|----------------|--|---|
| 982 | cleroda-3,13(14) <i>E</i> -diene-15,16-diamide | — | — | — | — | — | <i>Polyalthia longifolia</i> | <i>Phytochem. Lett.</i> , 2015, 11 , 28-31 |
| 983 | cleroda-3-ene-15,16-diamide | — | — | — | — | — | | |
| 984 | agelasine K | — | — | — | — | — | <i>Agelas cf. mauritiana</i> | <i>J. Nat. Prod.</i> , 2008, 71 , 1451–1454 |
| 985 | agelasine L | — | — | — | — | — | | |
| 986 | axistatin 2 | — | — | — | — | — | <i>Agelas axifera</i> | <i>J. Nat. Prod.</i> , 2013, 76 , 420-424 |
| 987 | axistatin 1 | H | Me | αMe | βMe | Z ₈ | | |
| 988 | agelasine P | =O | CH ₂ OZ ₅ | αMe | βMe | Z ₆ | <i>Agelas sp.</i> | <i>Tetrahedron</i> , 2012, 68 , 9738-9744 |
| 989 | agelasine Q | H | CH ₂ OZ ₅ | αMe | αMe | Z ₆ | | |
| 990 | agelasine R | H | CH ₂ OZ ₅ | βMe | βMe | Z ₆ | | |
| 991 | agelasine U | Z ₆ | — | — | — | — | | |
| 992 | scutebarbatine Z | ONic | OH | H | H | H | <i>Scutellaria barbata</i> | <i>Chem. Pharm. Bull.</i> , 2010, 58 , 1267-1270 |
| 993 | scutebarbatine X | ONic | ONic | OH | OAc | OH | | |
| 994 | scutebata B | ONic | OBz | OH | OAc | OH | | |
| 995 | scutebata C | ONic | OH | OH | OAc | OH | | |
| 996 | barbatellarine B | ONic | OBz | OH | OAc | H | | |
| 997 | scutebarbatine Y | H | αOBz | βONic | βOH | — | | |
| 998 | scutehenanine A | H | αOH | βONic | βOH | — | | |
| 999 | 6- <i>O</i> -acetylscutehenanine A | H | αOAc | βONic | βOH | — | | |
| 1000 | 6- <i>O</i> -(2-carbonyl-3-methylbutanoyl)scutehenanine A | H | αOM ₁ | βONic | βOH | — | | |
| 1001 | scutebarbatine B | H | αONic | βOBz | βOH | — | | |
| 1002 | scutelinquanine C | H | αONic | βY ₈ | βOH | — | <i>Phytochem. Lett.</i> , 2010, 3 , 190-193 | |


| | | | | | | | | | |
|-------------------|--------------------------------------|-------------------|----------------|------------------------|--------------|---|---------------------------------|---|--|
| 1003 | scutebarbatine A | H | α ONic | β ONic | β OH | — | | <i>Chin. Chem. Lett.</i> , 1996, 7 , 333-334 | |
| 1004 | scutebarbatine K | H | α ONic | β OAc | β OH | — | | <i>Chem. Pharm. Bull.</i> , 2008, 56 , 207-209 | |
| 1005 | scutebarbatine L | H | α ONic | β Y ₄ | β OH | — | | <i>Planta Med.</i> , 2007, 73 , 1217-1220 | |
| 1006 | 2-carbonylscutebarbatine A | =O | α ONic | β ONic | β OH | — | | | |
| 1007 | 6-O-nicotinoylbarbatin A | H | α ONic | β OH | β OH | — | | | |
| 1008 | 8-O-nicotinoylbarbatin A | H | α OH | β OH | β ONic | — | | <i>Chem. Pharm. Bull.</i> , 2006, 54 , 869-872 | |
| 1009 | scutebarbatine C | — | — | — | — | — | | | |
| 1010 | scutebarbatine D | β OH | α ONic | β OBz | — | — | | | |
| 1011 | scutebarbatine E | =O | α ONic | β OBz | — | — | | <i>J. Asian Nat. Prod. Res.</i> , 2009, 11 , 451-456 | |
| 1012 | scutebarbatine O | α OH | β ONic | α ONic | — | — | | | |
| 1013 | cleroda-3-ene yrrolidine-15,16-dione | — | — | — | — | — | | <i>Polyalthia longifolia</i> | <i>Phytochem. Lett.</i> , 2015, 11 , 28-31 |
| 1014 | cleroda-3-ene yrrrole-15,16-dione | Me | =O | =O | | | | | |
| 1015 | echinophyllin C | CO ₂ H | H ₂ | O | — | — | <i>Echinodorus macrophyllus</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 1576-1579 | |
| 1016 | echinophyllin F | CO ₂ H | O | H ₂ | — | — | | | |
| 1017 | echinophyllin D | O | H ₂ | — | — | — | | | |
| 1018 | 15-oxo-echinophyllin A | O | O | — | — | — | <i>Casearia sylvestris</i> | <i>Fitoterapia</i> , 2009, 80 , 404-407 | |
| 1019 | echinophyllin A | H ₂ | O | — | — | — | <i>Echinodorus macrophyllus</i> | <i>Tetrahedron Lett.</i> , 2000, 41 , 2939-2943 | |
| 1020 | echinophyllin B | H ₂ | O | — | — | — | | | |
| 1021 | echinophyllin E | O | H ₂ | — | — | — | | | <i>J. Nat. Prod.</i> , 2000, 63 , 1576-1579 |
| 1022 | cleidbrevoid A | β T | — | — | — | — | <i>Cleidion brevipetiolatum</i> | <i>Fitoterapia</i> , 2012, 83 , 1100-1104 | |
| 1023 | cleidbrevoid B | α T | — | — | — | — | | | |
| 1024 | cleidbrevoid C | β T | — | — | — | — | | | |
| 1025 ^a | teupernin D | — | — | — | — | — | <i>Teucrium pernyi</i> | <i>Chem. Pharm. Bull.</i> , 1992, 40 , 2193-2195 | |
| 1026 | teuracemin | — | — | — | — | — | <i>Teucrium racemosum</i> | <i>Phytochemistry</i> , 1995, 40 , 505-507 | |
| 1027 | scutehenanine D | OBz | ONic | OH | — | — | <i>Scutellaria barbata</i> | <i>J. Nat. Prod.</i> , 2009, 72 , 1793-1797 | |
| 1028 | scutebarbatine H | ONic | OH | OH | — | — | | <i>Chem. Pharm. Bull.</i> , 2007, 55 , 1218-1221 | |
| 1029 | 7-O-nicotinoylscutebarbatine H | ONic | ONic | OH | — | — | | <i>Chem. Pharm. Bull.</i> , 2008, 56 , 207-209 | |
| 1030 | scutebarbatine I | β OEt | — | — | — | — | | | |
| 1031 | scutebarbatine J | α OEt | — | — | — | — | | <i>Phytochem. Lett.</i> , 2010, 3 , 190-193 | |
| 1032 | scutelinquinane B | — | — | — | — | — | | | |
| 1033 | scutelinquinane A | OAc | H | — | — | — | | | |
| 1034 | scutehenanine H | OBz | OH | — | — | — | | <i>Fitoterapia</i> , 2010, 81 , 737-741 | |
| 1035 | scutelinquinane D | OH | OH | — | — | — | | <i>J. Asian Nat. Prod. Res.</i> , 2010, 12 , 859-864 | |


| | | | | | | | | |
|-------------|--|------|------|------|------|---|--|---|
| 1036 | scutehenanine B* → scutebarbatine W (1047) | H | OBz | OH | ONic | | | <i>J. Nat. Prod.</i> , 2009, 72 , 1793-1797 |
| 1037 | scutebarbatine F* → scutebata F (1045) = barbatine C (1048) | H | ONic | OAc | OAc | | | <i>Chem. Pharm. Bull.</i> , 2006, 54 , 869-872 |
| 1038 | scutebarbatine G* → 1039 | H | OH | OH | ONic | | | |
| 1039 | — | ONic | OH | OH | H | | | |
| 1040 | 6,7-di- <i>O</i> -nicotinoylscutebarbatine G* → 1041 | H | ONic | ONic | ONic | | | <i>Chem. Pharm. Bull.</i> , 2007, 55 , 1218-1221 |
| 1041 | | ONic | ONic | ONic | H | | | |
| 1042 | 6- <i>O</i> -nicotinoyl-7- <i>O</i> -acetylscutebarbatine G* → barbatine D (1049) | H | ONic | OAc | ONic | | | |
| 1043 | 6- <i>O</i> -nicotinoylscutebarbatine G* → 1044 | H | ONic | OH | ONic | | | <i>J. Asian Nat. Prod. Res.</i> , 2009, 11 , 451-456 |
| 1044 | | ONic | ONic | OH | H | — | | |
| 1045 | scutebata F = barbatine C (1048) | ONic | OAc | OAc | H | — | | <i>J. Nat. Prod.</i> , 2010, 73 , 233-236 |
| 1046 | scutebata G | OBz | ONic | OBz | H | — | | |
| 1047 | scutebarbatine W | OBz | ONic | OH | H | — | | <i>Chem. Pharm. Bull.</i> , 2010, 58 , 1267-1270 |
| 1048 | barbatine C= scutebata F (1045) | ONic | OAc | OAc | H | — | | |
| 1049 | barbatine D | ONic | ONic | OAc | H | — | | <i>Eur. J. Org. Chem.</i> , 2009, 5810-5815 |
| 1050 | barbatine A | H | ONic | OAc | ONic | — | | |
| 1051 | barbatine B | H | ONic | ONic | ONic | — | | |
| 1052 | scutehenanine C | | | | | | | <i>J. Nat. Prod.</i> , 2009, 72 , 1793-1797 |
| 1053 | barbatellarine C | | | | | | | |
| 1054 | barbatellarine D | | | | | | | <i>Helv. Chim. Acta.</i> , 2011, 94 , 643-649 |

^a Compound **776** in Table 24 was also given the name ‘teupernin D’.

*Original references for isolations listed in table; however, structure revisions (denoted by “→ revised compound name or number”) were described in *Chem. Pharm. Bull.*, 2010, **58**, 1267-1270.

9.2. Degraded Derivatives (Table 30)







| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | Source | Ref. |
|------|--|---------------------|----------------|----------------|----------------|--|---|
| 1055 | pentandranic acid B | — | — | — | — | <i>Callicarpa pentandra</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 1062-1065 |
| 1056 | sagitone | — | — | — | — | <i>Tinosporasagittata</i> var. <i>yunnanensi</i> | <i>Molecules</i> , 2010, 15 , 8360-8365 |
| 1057 | fiaruretin D | H | — | — | — | <i>Fibraurea tinctoria</i> | <i>Bioorg. Med. Chem.</i> , 2008, 16 , 9603-9609 |
| 1058 | fiaruretin E | αOH | — | — | — | | |
| 1059 | fiaruretin F | =O | — | — | — | | |
| 1060 | peniankerine | — | — | — | — | <i>Penianthus zenkeri</i> | <i>Phytochemistry</i> , 1997, 46 , 165-167 |
| 1061 | — | — | — | — | — | <i>Nardophyllum lanatum</i> | <i>Phytochemistry</i> , 1990, 29 , 1227-1230 |
| 1062 | cajucarin B | — | — | — | — | <i>Croton cajucara</i> | <i>Chem. Pharm. Bull.</i> , 1990, 38 , 701-705 |
| 1063 | 3α,4β-dihydroxy-15,16-epoxy-19-nor-12-oxo-cleroda-5(10),13(16),14-triene | — | — | — | — | <i>Croton hovarum</i> | <i>Phytochemistry</i> , 1997, 45 , 379-381 |
| 1064 | crotoeurin C | CO ₂ Me | — | — | — | <i>Croton euryphyllus</i> | <i>Bioorg. Med. Chem. Lett.</i> , 2015, 25 , 1329-1332 |
| 1065 | 15,16-epoxy-6a-O-acetyl-8β-hydroxy-19-nor-clero-13(16),14-diene-17,12;18,2-diolide | — | — | — | — | <i>Dioscorea bulbifera</i> | <i>Nat. Prod. Commun.</i> , 2011, 6 , 1069-1072 |
| 1066 | jatrophoidin | αCO ₂ Me | — | — | — | <i>Croton jatrophoides</i> | <i>Planta Med.</i> , 2009, 75 , 262-267 |
| 1067 | isoteucvin | βH | — | — | — | | |
| 1068 | crassifolin G | αOH | — | — | — | <i>Croton crassifolius</i> | <i>J. Nat. Prod.</i> , 2012, 75 , 2188-2192 |
| 1069 | teuponin | — | — | — | — | <i>Teucrium japonicum</i> | <i>Phytochemistry</i> , 1991, 30 , 4175-4177 |
| 1070 | teucvisin C | βOH | βH | — | — | <i>Teucrium viscidum</i> | <i>Chem. Pharm. Bull.</i> , 2014, 62 , 472-476 |
| 1071 | teucvisin D | αH | αOH | — | — | | |
| 1072 | teucvisin E | OH | — | — | — | | |

| | | | | | | | |
|------|--|--------------|--------------------|-------------|---|---|--|
| 1073 | crassifolin H | H | — | — | — | <i>Croton crassifolius</i> | <i>Heterocycles</i> , 2014, 89 , 1585-1593 |
| 1074 | cajucarinolide | OH | =O | — | — | <i>Croton cajucara</i> | <i>Planta Med.</i> , 1992, 58 , 549-551 |
| 1075 | isocajucarinolide | =O | OH | — | — | | |
| 1076 | <i>trans</i> -dehydrocrotonin | — | — | — | — | <i>Croton cajucara</i> | <i>J. Braz. Chem. Soc.</i> , 2014, 25 , 629-638 |
| 1077 | teucorymbin | H | OAc | H | — | <i>Teucrium corymbosum</i> | <i>Phytochemistry</i> , 1995, 40 , 1481-1483 |
| 1078 | syspirensin A | α OH | OH | α OH | — | <i>Teucrium chamaedrys</i> ssp. <i>syspirensis</i> | <i>J. Nat. Prod.</i> , 1996, 59 , 457-460 |
| 1079 | teupolin IX | H | OH | β OMe | — | <i>Teucrium polium</i> | <i>Phytochemistry</i> , 2011, 72 , 2037-2044 |
| 1080 | 12- <i>epi</i> -montanin B | H | OH | β OH | — | <i>Teucrium maghrebinum</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 1029-1031 |
| 1081 | teupolin VIII | OH | — | — | — | <i>Teucrium polium</i> | <i>Phytochemistry</i> , 2011, 72 , 2037-2044 |
| 1082 | teupolin VII | OMe | — | — | — | | |
| 1083 | crotoeurin B | — | — | — | — | <i>Croton euryphyllus</i> | <i>Bioorg. Med. Chem. Lett.</i> , 2015, 25 , 1329-1332 |
| 1084 | crotoeurin A | — | — | — | — | | |
| 1085 | (+)-7 β -acetoxy-14,15,16-trinor-3-clerodene-13,18-dioate | — | — | — | — | <i>Sindora sumatrana</i> | <i>Chem. Pharm. Bull.</i> , 1994, 42 , 1202-1207 |
| 1086 | 2-oxo-14,15-bisnor-3,11 <i>E</i> -kolavadien-13-one | =O | Me | — | — | <i>Polyalthia simiarum</i> | <i>Nat. Prod. Commun.</i> , 2010, 5 , 1543-1546 |
| 1087 | 14,15-bisnor-3,11 <i>E</i> -kolavadien-13-one | H | Me | — | — | <i>Polyalthia viridis</i> | <i>Phytochemistry</i> , 1990, 29 , 653-655 |
| 1088 | 14,15-dinorclerod-3-ene-2,13-didione | =O | Me | — | — | <i>Clausena dunniana</i> | <i>Helv. Chim. Acta.</i> , 2003, 86 , 3187-3193 |
| 1089 | bis-norinfuscaic acid | H | CO ₂ H | — | — | <i>Jungermannia infusca</i> | <i>Phytochemistry</i> , 1998, 49 , 601-608 |
| 1090 | 2-oxo-5 α ,8 β -13,14,15,16-tetranorclerod-3-en-12-oic acid | — | — | — | — | <i>Vellozia bicolor</i> | <i>Phytochemistry</i> , 1994, 37 , 1115-1117 |
| 1091 | 18-hydroxyaylthonic acid | — | CH ₂ OH | — | — | <i>Dicranopteris dichotoma</i> | <i>J. Nat. Prod.</i> , 2007, 70 , 265-268 |
| 1092 | 18-oxo-aylthonic acid | — | CHO | — | — | | |
| 1093 | crotinsulactone | — | — | — | — | <i>Croton insularis</i> | <i>Helv. Chim. Acta.</i> , 2005, 88 , 2654-2660 |
| 1094 | ciliatolide A | — | — | — | — | <i>Scapania ciliata</i> | <i>Chem. Biodivers.</i> , 2013, 10 , 1606-1612 |
| 1095 | teucrolin D | α OAc | — | — | — | <i>Teucrium oliverianum</i> | <i>J. Nat. Prod.</i> , 1993, 56 , 830-842 |
| 1096 | teucrolivin F | =O | — | — | — | | <i>Phytochemistry</i> , 1991, 30 , 1603-1606 |
| 1097 | — | — | — | — | — | <i>Jamesoniella colorata</i> | <i>Nat. Prod. Commun.</i> , 2010, 5 , 999-1003 |

9.3 Ring-*seco* Derivatives (Table 31)







| No. | Compound Name | R ₁ | R ₂ | R ₃ | Source | Ref. |
|------|--|----------------|----------------|----------------|--------------------------------|---|
| 1098 | jamesoniellide F | — | — | — | <i>Jamesoniella autumnalis</i> | <i>Phytochemistry</i> , 1995, 39 , 859-868, |
| 1099 | jamesoniellide I | — | — | — | | <i>Phytochemistry</i> , 1999, 51 , 743-750 |
| 1100 | jamesoniellide K | OH | =O | — | <i>Jamesoniella colorata</i> | <i>Phytochemistry</i> , 2003, 63 , 227-233 |
| 1101 | jamesoniellide L | =O | OH | — | | |
| 1102 | salvianduline A | αOAc | — | — | <i>Salvia lavanduloides</i> | <i>Phytochemistry</i> , 1991, 30 , 3357-3360 |
| 1103 | salvianduline B | βOH | — | — | | |
| 1104 | pyrrhopappolide | H | — | — | <i>Microglossa pyrrhopappa</i> | <i>Phytochemistry</i> , 1990, 29 , 3233-3241 |
| 1105 | 6β-hydroxypyrrhopappolide | OH | — | — | | |
| 1106 | 6β-angeloyloxypyrrhopappolide | OAng | — | — | | |
| 1107 | 6β-seneciolyloxypyrrhopappolide | OSen | — | — | | |
| 1108 | 6β-tigloyloxypyrrhopappolide | OTig | — | — | | |
| 1109 | 6β-[2-methylbutyryloxy]pyrrhopappolide | OMeBu | — | — | | |
| 1110 | 15,16-epoxy-8-hydroxy-1,3,13(16),14-clerodatetraene-17,12:18,6-diolide | — | — | — | <i>Jamesoniella autumnalis</i> | <i>Phytochemistry</i> , 1995, 39 , 859-868 |
| 1111 | — | — | — | — | <i>Salvia miniata</i> | <i>Phytochemistry</i> , 2011, 72 , 265-275 |
| 1112 | jamesoniellide D | H | H | — | <i>Jamesoniella autumnalis</i> | <i>Phytochemistry</i> , 1995, 39 , 859-868, <i>Phytochemistry</i> , 1999, 51 , 743-750 |
| 1113 | jamesoniellide E | =O | — | — | | |
| 1114 | jamesoniellide H | αOH | — | — | | |
| 1115 | cephaloziellin H | αH | — | — | <i>Cephaloziella kiaeri</i> | <i>J. Nat. Prod.</i> , 2013, 76 , 1700-1708 |
| 1116 | cephaloziellin I | βH | — | — | | |
| 1117 | cephaloziellin J | — | — | — | | |
| 1118 | cephaloziellin K | — | — | — | | |
| 1119 | seco-eeoniolide | — | — | — | <i>Pteronia incana</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |
| 1120 | jamesoniellide A | — | — | — | <i>Jamesoniella autumnalis</i> | <i>J. Nat. Prod.</i> , 1992, 55 , 111-121 |


| | | | | | | |
|------|--|--------------|-------------------|--------------------|--------------------------------|---|
| 1121 | jamesoniellide B | — | — | — | | |
| 1122 | salvimicrophyllin A | — | — | — | <i>Salvia microphylla</i> | <i>J. Nat. Prod.</i> , 2014, 77 , 1088-1092 |
| 1123 | salvimicrophyllin B | — | — | — | | |
| 1124 | <i>seco</i> -hypoleucolide | — | — | — | <i>Conyza hypoleuca</i> | <i>Phytochemistry</i> 1991, 30 , 575-581 |
| 1125 | 17 α -hydroxy-12 β ,17-epoxystrictic acid | α OH | β H | CO ₂ H | <i>Conyza welwitschii</i> | <i>Phytochemistry</i> , 1990, 29 , 2247-2252 |
| 1126 | 17 β -hydroxy-12 β ,17-epoxystrictic acid | β OH | β H | CO ₂ H | | |
| 1127 | 17 α - <i>O</i> -methyl ether of 1125 | α OMe | β H | CO ₂ H | | |
| 1128 | 17 β - <i>O</i> -methyl ether of 1126 | β OMe | β H | CO ₂ H | | |
| 1129 | strictic acid 12 β ,17-olide | =O | β H | CO ₂ H | | |
| 1130 | — | =O | H | CO ₂ Me | <i>Croton jimenezii</i> | <i>Ingenieria Y Ciencia Quimica</i> , 2000, 19 , 68-73 |
| 1131 | 7 α -hydroxystrictic acid | OH | Me | H | <i>Conyza hypoleuca</i> | <i>Phytochemistry</i> 1991, 30 , 575-581 |
| 1132 | 11,12 <i>E</i> -dehydrostrictic acid | H | Me | H | | |
| 1133 | 12-hydroxystrictic acid | H | Me | β OH | <i>Conyza welwitschii</i> | <i>Phytochemistry</i> , 1990, 29 , 2247-2252 |
| 1134 | 12-oxo-17- <i>oic</i> -strictic diacid | H | CO ₂ H | =O | | |
| 1135 | dodonolide | — | — | — | <i>Dodonaea viscosa</i> . | <i>Tetrahedron</i> , 2001, 57 , 2981-2989 |
| 1136 | tonalensin | — | — | — | <i>Salvia tonalensis</i> | <i>J. Chem. Crystallogr.</i> , 1996, 26 , 239-242 |
| 1137 | tinosporafuranol | H | — | — | <i>Tinospora cordifolias</i> | <i>Nat. Prod. Res.</i> , 2010, 24 , 926-9354 |
| 1138 | tinosporafurandiol | OH | — | — | | |
| 1139 | fruticolide | — | — | — | <i>Teucrium fruticans</i> | <i>Phytochemistry</i> , 1992, 31 , 3531-3534 |
| 1140 | rhyacophiline | — | — | — | <i>Salvia rhyacophila</i> | <i>Tetrahedron</i> , 1991, 47 , 7199-7208 |
| 1141 | salvireptanolide | — | — | — | <i>Salvia reptans</i> | <i>Phytochemistry</i> , 1991, 30 , 2335-2338 |
| 1142 | jamesoniellide J | — | — | — | <i>Jamesoniella autumnalis</i> | <i>Phytochemistry</i> , 1999, 51 , 743-750 |
| 1143 | cephaloziellin E | — | — | — | <i>Cephaloziella kiaeri</i> | <i>J. Nat. Prod.</i> , 2013, 76 , 1700-1708 |
| 1144 | cephaloziellin F | — | — | — | | |
| 1145 | cephaloziellin G | — | — | — | | |

9.4. Rearranged Derivatives (Table 32)







1172




1173




1174,1175




1176,1177




1178,1179




1180




1181




1182




1183-1190




1191,1192




1193




1194,1195




1196-1198




1199




1200




1201,1202




1203,1204




1205




1206



1207-1210



1211



| No. | Compound Name | R ₁ | R ₂ | R ₃ | R ₄ | Source | Ref. |
|------|---|-------------------|----------------|--------------------|----------------|--|---|
| 1146 | — | — | — | — | — | <i>Solidago altissima</i> | <i>Phytochemistry</i> , 1999, 52 , 487-493 |
| 1147 | — | CHO | α OAc | CO ₂ Me | — | | |
| 1148 | (4→2)- <i>abeo</i> -cleroda-2,13 <i>E</i> -dien-2,14-dioic acid | CO ₂ H | H | CO ₂ H | — | <i>Polyalthia longifolia</i> var. <i>pendula</i> | <i>Molecules</i> , 2014, 19 , 2049-2060 |
| 1149 | (4→2)- <i>abeo</i> -2,13-diformyl-cleroda-2,13 <i>E</i> -dien-14-oic acid | — | — | — | — | | |
| 1150 | dunniana acid A | CO ₂ H | — | — | — | <i>Clausena dunniana</i> | <i>J. Nat. Prod.</i> , 2002, 65 , 392-394 |
| 1151 | dunniana acid B | CHO | — | — | — | | |
| 1152 | (+)-(4→2)- <i>abeo</i> -kolavelool-3-oic acid | — | — | — | — | <i>Aristolochia chamissonis</i> | <i>Phytochemistry</i> , 1999, 50 , 455-461 |
| 1153 | pentandranoic acid A | — | — | — | — | <i>Callicarpa pentandra</i> | <i>J. Nat. Prod.</i> , 2000, 63 , 1062-1065 |
| 1154 | pentandralactone | CHO | α OH | H | — | | |
| 1155 | (4→2)- <i>abeo</i> -16(R&S)-2,13 <i>Z</i> -kolavadien-16,15-olide-3-al | CHO | H | β OH | — | <i>Polyalthia viridis</i> | <i>Phytochemistry</i> , 1990, 29 , 653-655 |
| 1156 | polylongifoliaic A | CO ₂ H | H | OH | — | <i>Polyalthia longifolia</i> var. <i>pendula</i> | <i>RSC Advances</i> , 2014, 4 , 23707-23712 |
| 1157 | pilosanone C | — | — | — | — | <i>Portulaca pilosa</i> | <i>Phytochemistry</i> , 1995, 40 , 205-207 |
| 1158 | methyl dodovisate A | — | — | — | — | <i>Dodonaea viscosa</i> | <i>J. Asian. Nat. Prod. Res.</i> , 2010, 12 , 7-14 |
| 1159 | methyl dodovisate B | — | — | — | — | | |
| 1160 | scapanialide B | — | — | — | — | <i>Scapania parva</i> | <i>Phytochem. Lett.</i> , 2012, 5 , 535-540 |
| 1161 | isosalvipuberulin | — | — | — | — | <i>Salvia leucantha</i> | <i>J. Nat. Med.</i> , 2006, 60 , 206-209 |
| 1162 | salvileucantholide | — | — | — | — | | <i>Tetrahedron</i> , 1994, 50 , 11593-11600 |
| 1163 | dugesin B | — | — | — | — | <i>Salvia dugesii</i> | <i>Helv. Chim. Acta</i> , 2004, 87 , 949-955 |
| 1164 | spiroleucantholide | — | — | — | — | <i>Salvia leucantha</i> | <i>J. Nat. Med.</i> , 2006, 60 , 206-209 |
| 1165 | salvioccidentalinalin | — | — | — | — | <i>Salvia occidentalis</i> | <i>Molecules</i> , 2011, 16 , 9109-9115 |
| 1166 | dugesin C | — | — | — | — | <i>Salvia dugesii</i> | <i>Nat. Prod. Bioprospect.</i> , 2011, 1 , 81-86 |
| 1167 | blepharolide B | — | — | — | — | <i>Salvia blepharophylla</i> | <i>Phytochemistry</i> , 1999, 52 , 1535-1540 |
| 1168 | 2 β -hydroxysalvigenolide | — | — | — | — | <i>Salvia xalapensis</i> | <i>J. Nat. Prod.</i> , 2005, 68 , 787-790 |

| | | | | | | | |
|------|---|-------------|--------------|--------------|-------------|--------------------------------|---|
| 1169 | salviandulin E | — | — | — | — | <i>Salvia leucantha</i> | <i>Tetrahedron</i> , 1994, 50 , 11593-11600 |
| 1170 | dugesin A | — | — | — | — | <i>Salvia dugesii</i> | <i>Helv. Chim. Acta</i> , 2004, 87 , 949-955 |
| 1171 | dugesin D | — | — | — | — | | <i>Nat. Prod. Bioprospect.</i> , 2011, 1 , 81-86 |
| 1172 | blepharolide A | — | — | — | — | <i>Salvia blepharophylla</i> | <i>Phytochemistry</i> , 1999, 52 , 1535-1540 |
| 1173 | tilifodiolide | — | — | — | — | <i>Salvia dugesii</i> | <i>J. Org. Chem.</i> , 1990, 55 , 3522-3525 |
| 1174 | baenzigeride A | H | — | — | — | <i>Tinospora baenzigeri</i> | <i>Phytochemistry</i> , 1999, 52 , 1335-1340 |
| 1175 | baenzigeroside A | Glc | — | — | — | | |
| 1176 | baenzigeride B | H | — | — | — | | <i>Chem. Pharm. Bull.</i> , 2001, 49 , 854-857 |
| 1177 | baenzigeroside B | Glc | — | — | — | | |
| 1178 | — | α H | — | — | — | the smoke of salvinorin A | <i>Tetrahedron Lett.</i> , 2010, 51 , 5207-5209 |
| 1179 | — | β H | | | | | |
| 1180 | microphyllandiolide | — | — | — | — | <i>Salvia microphylla</i> | <i>Org. Lett.</i> , 2013, 15 , 3210-3213 |
| 1181 | teubetonin | | | | | <i>Teucrium betonicum</i> | <i>Tetrahedron</i> , 1995, 51 , 2363-2368 |
| 1182 | 7 β ,8 β -epoxyisochiliolide lactone | — | — | — | — | <i>Microglossa pyrrhopappa</i> | <i>Phytochemistry</i> , 1990, 29 , 3233-3241 |
| 1183 | 6 β -angeloxyisochiliolide lactone | OAng | H | H | =O | | |
| 1184 | 6 β -[2-methylbutyryloxy]-isochiliolide lactone | OMeBu | H | H | =O | | |
| 1185 | 8 β -hydroxyisochiliolide lactone | H | H | OH | =O | | |
| 1186 | isochiliolide lactone | H | H | H | =O | | |
| 1187 | 7 β -angeloxyisochiliolide lactone | H | β OAng | β H | =O | <i>Pteronia divaricata</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |
| 1188 | 3 α -hydroxypteronia lactone | OAc | β H | β OAng | α OH | | |
| 1189 | 3 β -hydroxypteronia lactone | OAc | β H | β OAng | β OH | | |
| 1190 | 3-oxo-pteronia lactone | OAc | β H | β OAng | =O | <i>Microglossa pyrrhopappa</i> | <i>Phytochemistry</i> , 1990, 29 , 3233-3241 |
| 1191 | 8(17)-dehydroisochiliolide lactone | H | — | — | — | | |
| 1192 | 7 β -acetoxy-8(17)-dehydro- isochiliolide lactone | OAc | — | — | — | | |
| 1193 | 6 β -hydroxyisochiliolide | — | — | — | — | | |
| 1194 | 6 β -hydroxy-incana-pteroniolide | β OH | — | — | — | <i>Pteronia incana</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |
| 1195 | incanapteroniolide | H | — | — | — | | |
| 1196 | 19 α -hydroxypteronia-dilactone | α OH | — | — | — | <i>Pteronia eenii</i> | <i>Phytochemistry</i> , 1990, 29 , 1231-1245 |
| 1197 | 19 β -hydroxypteronia-dilactone | β OH | — | — | — | | |
| 1198 | pteroniatriolactone | =O | — | — | — | | |
| 1199 | eeniolide | — | — | — | — | | |
| 1200 | jamesoniellide C | — | — | — | — | <i>Jamesoniella autumnalis</i> | <i>Phytochemistry</i> , 1994, 37 , 491-494 |

| | | | | | | | |
|------|---------------------|------------|------------|------------|---|------------------------------|---|
| 1201 | alysine A | OH | OAc | — | — | <i>Teucrium alyssifolium</i> | <i>Tetrahedron</i> , 1995, 51 , 11793-11800. |
| 1202 | alysine C | H | OH | — | — | | |
| 1203 | alysine B | OAc | — | — | — | | |
| 1204 | 3-deacetylalysine B | OH | — | — | — | | |
| 1205 | alysine D | — | — | — | — | | |
| 1206 | alysine E | — | — | — | — | | |
| 1207 | salvilanguiduline A | H | H | H | — | <i>Salvia languidula</i> | <i>Tetrahedron Lett.</i> , 1992, 33 , 581-584 |
| 1208 | salvilanguiduline B | OH | H | H | — | | |
| 1209 | salvilanguiduline C | H | OH | H | — | | |
| 1210 | salvilanguiduline D | H | H | OH | — | | |
| 1211 | cephaloziellin Q | — | — | — | — | <i>Jamesoniella colorata</i> | <i>Nat. Prod. Commun.</i> , 2010, 5 , 999-1003 |
| 1212 | cephaloziellin L | β H | α H | Δ^3 | | <i>Cephaloziella kiaeri</i> | <i>J. Nat. Prod.</i> , 2013, 76 , 1700-1708 |
| 1213 | cephaloziellin M | β H | β H | Δ^3 | | | |
| 1214 | cephaloziellin N | α H | α H | | | | |
| 1215 | cephaloziellin O | β H | α H | | | | |
| 1216 | cephaloziellin P | β H | β H | | | | |
| 1217 | teubrevin E | — | — | — | — | <i>Teucrium brevifolium</i> | <i>Tetrahedron</i> , 1995, 51 , 837-848 |
| 1218 | teubrevin F | — | — | — | — | | |
| 1219 | teubrevin G | — | — | — | — | | |
| 1220 | teubrevin H | — | — | — | — | | |
| 1221 | teubrevin I | — | — | — | — | | |
| 1222 | salvixalapoxide | — | — | — | — | <i>Salvia xalapensis</i> | <i>J. Nat. Prod.</i> , 2005, 68 , 787-790 |
| 1223 | salvixalapadiene | — | — | — | — | | |
| 1224 | isosalvixalapadiene | — | — | — | — | | |

Abbreviation of Functional Groups

