A novel fluorescent retrograde neural tracer: cholera toxin B conjugated carbon dots

Nan Zhou,^a Zeyu Hao,^a Xiaohuan Zhao,^b Suraj Maharjan,^a Shoujun Zhu,^b Yubin Song,^b Bai Yang^{*b} and Laijin Lu^{*a}

Table of the Electronic Supplementary Information

- 1. Fig. S1 Photoluminescence (PL) spectra of CTB.
- 2. Fig. S2 UV-vis absorption spectra of the dialyzate.
- 3. Fig. S3 Fluorescence signal of CTB-CDs in L4 DRG.
- 4. Fig. S4 Immunohistochemistry of CTB-CDs in L4 DRG.
- 5. Fig. S5 Low magnification image of immunohistochemistry of L5 spinal cord.
- 6. Fig. S6 Stability of CTB-CDs. The PL intensity of CTB-CDs was stable after 4 h of continuous UV excitation (a) and it remained constant in a solution of pH 4-9 (b).
- 7. Fig. S7 CTB-CDs labelled neural cells in the longitudinal section of spinal cord.
- **8.** Fig. S8 Histological analysis of brain and spinal cord in the control group and CDs-injected rats (100 mg/kg) at 1 day, 7 days and 28 days. The scale bar is 50 μm.



Fig. S1 Photoluminescence (PL) spectra of CTB.



Fig. S2 UV-vis absorption spectra of the dialyzate.



Fig. S3 Fluorescence signal of CTB-CDs in L4 DRG.



Fig. S4 Immunohistochemistry of CTB-CDs in L4 DRG.



Fig. S5 Low magnification image of immunohistochemistry of L5 spinal cord.



Fig. S6 Stability of CTB-CDs. The PL intensity of CTB-CDs was stable after 4 h of continuous UV excitation (a) and it remained constant in a solution of pH 4-9 (b).



Fig. S7 CTB-CDs labelled neural cells in the longitudinal section of spinal cord.



Fig. S8 Histological analysis of brain and spinal cord in the control group and CDs-injected rats (100 mg/kg) at 1 day, 7 days and 28 days. The scale bar is 50 μ m.