

Supporting Information for

Manganese(III)-mediated alkenyl C_{sp2}-P bond formation from the reaction of β -nitrostyrenes with dialkyl phosphites

Jian-Fei Xue,^a Shao-Fang Zhou,^a Ye-Ye Liu,^a Xiangqiang Pan,^{*a} Jian-Ping Zou^{*a,b} and

Olayinka Taiwo Asekun^c

^a *Key Laboratory of Organic Synthesis of Jiangsu Province, College of Chemistry and Chemical Engineering, Soochow University, 199 Renai Street, Suzhou, Jiangsu 215123, China; E-mail: jpzou@suda.edu.cn, panxq@suda.edu.cn*

^b *Key Laboratory of Synthetic Chemistry of Natural Substances, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, China.*



^c *Department of Chemistry, University of Lagos, Akoka, Yaba, Nigeria.*


Table of contents

S-2 ¹H, ¹³C NMR spectra of 3a-3u, and ³¹P NMR spectra of compounds 3a, 3f-3o, 3q-3u


NMR Spectra


3a







3b







3c







3d







3e







3f







3g


S8








3h















3j








3k








31








3m









3n







3o













3q







3r










3t

S27





3u

