

Electronic Supplementary Information

“Corrole and nucleophilic aromatic substitution are not incompatible: a novel route to 2,3-difunctionalized copper corroles”

M. Stefanelli, ^{*a} F. Mandoj,^a S. Nardis,^a M. Raggio,^a F.R. Fronczek,^b G.T. McCandless,^c K. M. Smith,^b and R. Paolesse^a

manuela.stefanelli@uniroma2.it; kmsmith@lsu.edu

^a Dipartimento di Scienze e Tecnologie Chimiche, Università di Roma Tor Vergata, via della Ricerca Scientifica 1, 00133 Rome, Italy

^b Department of Chemistry, Louisiana State University, Baton Rouge, LA 70803, USA

^c Department of Chemistry, The University of Texas at Dallas, Richardson, TX 75080, USA.

Copies of ¹H NMR and FAB mass spectra




Figure S1: ¹H NMR spectrum of 2-DEM-3NO₂-TtBuPCorrCu(**1**)




Figure S2: ${}^1\text{H}$ NMR spectrum of 2-DEM(OH)-3NO₂-TtBuPCorrCu(2)




Figure S3: ^1H NMR spectrum of 2-EtO-3-(NO₂)TtBuCorrCu(**3**)




Figure S4: FAB mass spectrum of 2-DEM-3NO₂-TtBuPCorrCu(**1**)




Figure S5: FAB mass spectrum of 2-DEM(OH)-3NO₂-TtBuPCorrCu(**2**)




Figure S6: FAB mass spectrum of 2-EtO-3-(NO₂)TtBuCorrCu(**3**)