## *meso*-Tetraphenylporphyrin with pi-system extended by fusion with anthraquinone

Mikhail A. Filatov, \*a,b Ernesta Heinrich, a Katharina Landfester a and Stanislav Baluschev a,c,d

<sup>a</sup> Max Planck Institute for Polymer Research, Ackermannweg 10, D-55128 Mainz, Germany, Fax: +496131379100 e-mail: filatov@polymer.bas.bg

<sup>b</sup> Institute of Polymers, Bulgarian Academy of Sciences, Acad. G. Bonchev Str., block 103-A, BG - 1113 Sofia, Bulgaria

<sup>c</sup> Optics and Spectroscopy Department, Faculty of Physics, Sofia University 'St. Kliment Ochridski'', 5 James Bourchier, 1164 Sofia, Bulgaria

<sup>d</sup> Freiburg Institute for Advanced Studies (FRIAS), Albert-Ludwigs-Universität Freiburg, Albertstraße 19, D-79104 Freiburg, Germany



Fig. S2. <sup>13</sup>C NMR spectrum of compound **2**.















Fig. S14. <sup>13</sup>C NMR spectrum of compound 8.







Fig. S18. <sup>13</sup>C NMR spectrum of compound **10**.



Fig. S20. <sup>13</sup>C NMR spectrum of compound **11**.





Fig. S23. MALDI TOF spectrum of 12 (positive mode).



Fig. S24. <sup>1</sup>H NMR spectrum of compound **Ph<sub>4</sub>TAQP**.



Fig. S25. <sup>13</sup>C NMR spectrum of compound Ph<sub>4</sub>TAQP.



Fig. S26. MALDI TOF spectrum of **Ph<sub>4</sub>TAQP** (positive mode).



Fig. S27. MALDI TOF spectrum of Ph<sub>4</sub>TAQP -Zn (positive mode).



Fig. S28. MALDI TOF spectrum of Ph<sub>4</sub>TAQP -Pd (positive mode).