


Supporting Information

Epigenetic stimulation of polyketide production in *Chaetomium cancroideum* by an NAD⁺-dependent HDAC inhibitor


Teigo Asai^{*a}, Shuntaro Morita^a, Tohru Taniguchi^b, Kenji Monde^b and Yoshiteru Oshima^{*a}


^a Graduate School of Pharmaceutical Sciences, Tohoku University, Aoba-yama, Aoba-ku, Sendai 980-8578, Japan. ^b Faculty of Advanced Life Science, Frontier Research Center for Post-Genome Science and Technology, Hokkaido University, Kita 21 Nishi 11, Sapporo 001-0021, Japan.

¹ H NMR spectrum of 1	S2
¹³ C NMR spectrum of 1	S3
¹ H NMR spectrum of 2	S4
¹³ C NMR spectrum of 2	S5
¹ H NMR spectrum of 3	S6
¹³ C NMR spectrum of 3	S7





^1H NMR spectrum of **1** (600 MHz, CDCl_3)







^{13}C NMR spectrum of **1** (150 MHz, CDCl_3)










^{13}C NMR spectrum of **2** (150 MHz, CDCl_3)





^1H NMR spectrum of **3** (600 MHz, CDCl_3)





^{13}C NMR spectrum of **3** (150 MHz, CDCl_3)

