

Electronic Supplementary Information for

Semisynthetic photoprotein reporters for tracking fast fast Ca²⁺ transients

Natalia P. Malikova,^a Aren J. Borgdorff^b, and Eugene S. Vysotski^{a,†}

^a*Photobiology Laboratory, Institute of Biophysics, Russian Academy of Sciences, Siberian Branch, Krasnoyarsk, 660036, Russia;*

^b*Institut des Neurosciences Alfred Fessard, UPR 3294, Centre National de la Recherche Scientifique, Avenue de la Terrasse, 91198 Gif-sur-Yvette, France*

†Corresponding author:

Fax: +7 (391) 243-3400; Tel: +7 (391) 249-4430;

E-mail: <eugene.vysotski@gmail.com> or <eugene_vysotski@ibp.ru>




Figure S1. Stopped-flow records of the rising phase (left panel) and full-time course (right panel) of luminescence signal for aequorin reconstituted with native CTZ (A), CTZ-*f* (B), CTZ-*i* (C), and CTZ-*hcp* (D) without Mg^{2+} (black) and with 1 mM Mg^{2+} (red).




Figure S2. Stopped-flow records of the rising phase (left panel) and full-time course (right panel) of luminescence signal for obelin reconstituted with native CTZ (A), CTZ-*f* (B), CTZ-*i* (C), and CTZ-*hcp* (D) without Mg^{2+} (black) and with 1 mM Mg^{2+} (red). Inset: Stopped-flow records of the full-time course of luminescence signal for obelin-*hcp* without Mg^{2+} (black) and with 1 mM Mg^{2+} (red).




Figure S3. Stopped-flow records of the rising phase (left panel) and full-time course (right panel) of luminescence signal for clytin reconstituted with native CTZ (A), CTZ-*f* (B), CTZ-*i* (C), and CTZ-*hcp* (D) without Mg^{2+} (black) and with 1 mM Mg^{2+} (red). Inset: Stopped-flow records of the full-time course of luminescence signal for clytin-*hcp* without Mg^{2+} (black) and with 1 mM Mg^{2+} (red).