

The Electrophilic Effect of Thiol Group on Thiol-yne Thermal Click Polymerization for Hyperbranched Polythioether

Bo Zhao,^a Yaochen Zheng,^{a,b} Zhulin Weng,^{a,c} Shengying Cai,^a and Chao Gao^{a*}

^a MOE Key Laboratory of Macromolecular Synthesis and Functionalization, Department of Polymer Science and Engineering, Zhejiang University, Hangzhou 310027, P. R. China

^b Department of Polymer Science and Engineering, College of Chemistry and Chemical Engineering, Yantai University, 30 Qingquan Road, Yantai 264005, P. R. China

^c School of Mathematical and Physical Sciences, Hubei University for Nationalities, Enshi, 445000, P. R. China

Figure and table of contents:




Fig. S1 ¹H NMR spectra of the starting materials 1-5.




Fig. S2 ^{13}C NMR spectra of HP1-8.




Fig. S3 ^1H NMR spectra of the HP 2 recorded at given time intervals.




Fig. S4 ^{13}C NMR spectra of HP 2-8.




Fig. S5 FTIR spectra of the HP 2 recorded at given time intervals.




Fig. S6 ^1H NMR spectra of the HP 4 recorded at given time intervals.




Fig. S7 ^{13}C NMR spectra of HP 4-8.




Fig. S8 FTIR spectra of the HP 4 recorded at given time intervals.