

Supporting Information

Palladium-Schiff-base-Silica Framework as a Robust and Recyclable Catalyst for the Suzuki-Miyaura Cross-Coupling in Aqueous Media

Tahshina Begum^a, Manoj Mondal^a, Pradip K. Gogoi*^a and Utpal Bora*^{ab}

^aDepartment of Chemistry, Dibrugarh University, Dibrugarh 786004, Assam, India

^bDepartment of Chemical Sciences, Tezpur University, Napaam, Tezpur 784028, Assam, India

E-mails: dr.pradip54@gmail.com (P. K. Gogoi), utbora@yahoo.co.in, ubora@tezu.ernet.in (U. Bora)

Content:

- | | |
|---|--------|
| 1. General information about the NMR instrument..... | S2 |
| 2. ¹ H NMR spectra of the products of the Suzuki-Miyaura Reaction..... | S3-S17 |

General information about the ^1H NMR spectra:

^1H spectra were recorded in CDCl_3 using tetramethylsilane (TMS) as an internal standard on a Advance DPX 300 MHz FT-NMR spectrometer operating at 300 MHz (Table 3, entries 2, 3, 6, 7, 8, 9, 11 and 17), on JEOL, JNM ECS NMR spectrometer operating at 400 MHz (Table 3, entries 1, 4, 5, 12, 14, 15, 18, 19 and 20), and Bruker Avance 500 MHz spectrometer operating at 500 MHz (Table 3, entries 10 and 13).





























