

Supporting Information for

Disulfide cross-linked cholic-acid modified PEG-poly(amino acid) block copolymer micelles for controlled drug delivery of doxorubicin

5 Yuling Li,^{*a} Yuling Shen,^a Sai Wang,^a Dandan Zhu,^a Baixiang Du^a and Jihong Jiang^b

^a School of Chemistry and Chemical Engineering, Jiangsu Normal University, Xuzhou 221116, P. R. China.

^b Key Laboratory of Biotechnology for Medicinal Plant of Jiangsu Province, Jiangsu Normal University, 10 Xuzhou 221116, P. R. China.

*Corresponding Author. Tel: +86-516-83403165; E-mail: ylli19722@163.com

15

20

25




Figure S1. The calibration curve of DOX in PB (10 mM, pH 7.4) based on fluorescence measurements (at 560 nm).

5




Figure S2. Stability of cross-linked PEG-pGlu (EDA-LA)₁₅-CA micelles against a) 100-fold dilution, b) 0.15 M or 2 M NaCl. The parent non-cross-linked micelles were used as a control.

15




Figure S3. DOX release from core-cross-linked PEG-pGlu (EDA-LA)₂₆-CA micelles in PB buffer (10 mM) at 37 °C. Non-cross-linked micelles were used as a control. Data are presented as mean ± SD (n = 3).

