Supplementary information: Carbon supported catalysts in low temperature steam reforming of ethanol: Study of catalyst performance

Anne-Riikka Rautio,1* Prem Kumar Seelam,2 Päivi Mäki-Arvela,3 Olli Pitkänen,1 Mika Huuhtanen,2

Riitta L. Keiski,² Krisztian Kordas¹

¹Microelectronics and Materials Physics Laboratories, Department of Electrical Engineering, Faculty of Information Technology and Electrical Engineering, University of Oulu, P.O. Box 4500, FI-90014 Oulu, Finland

² Environmental and Chemical Engineering, Faculty of Technology, P.O. Box 4500, FI-90014 Oulu, Finland

³Johan Gadolin Process Chemistry Centre, Åbo Akademi University, FI-20500 Åbo, Finland




Figure S1 The concentrations (vol.%) of minor products formed over Ni supported catalysts (Ni/CNT and Ni/GC) in SRE reaction: (a) CH_3CHO , (b) CH_4 , (c) CO and (d) CO_2 .

^{*} Corresponding author. Tel+358 50 364 5199.E-mail: <u>anneriik@ee.oulu.fi</u> (A.-R. Rautio)




Figure S2 The concentrations (vol.%) of minor products formed over Pt supported catalysts (Pt/CNT and Pt/GC) in SRE reaction: (a) CH₃CHO, (b) CH₄, (c) CO and (d) CO₂.




Figure S3 The concentrations (vol.%) of minor products formed over Pd supported catalysts (Pd/CNT and Pd/GC) in SRE reaction: (a) CH_3CHO and (b) CH_4 .




Figure S4 TPR-H₂ profiles of Ni doped carbon-based catalysts.




Figure S5 XRD patterns recorded to catalyst materials and their supports. (XRD patterns: Pt: 00-004-0802; Pd: 00-046-1043.)