

**Effective catalytic hydrodechlorination of chlorophenoxyacetic acids over  
Pd/graphitic carbon nitride**

Fang Jiang<sup>a</sup>, Wenhui Tan<sup>a</sup>, Huan Chen<sup>a\*</sup>, Ling Tan<sup>a</sup>, Jingliang Liu<sup>b</sup>

*a Key Laboratory of Jiangsu Province for Chemical Pollution Control and Resources Reuse,  
School of Environmental and Biological Engineering, Nanjing University of Science and  
Technology, Nanjing 210094, China.*

*b School of Biochemical Environmental and Engineering, Nanjing XiaoZhuang University,  
Nanjing 211171, China.*

\*Corresponding author. Tel: +86-25-84303209; Fax: +86-25-84315352.

*E-mail: hchen404@njjust.edu.cn (H. Chen)*

Fig. 1S Liquid phase catalytic HDC of 2,4-D over catalyst 3 with various catalyst dosages. ( $\square$ )  $0.05 \text{ g L}^{-1}$ , ( $\blacksquare$ )  $0.1 \text{ g L}^{-1}$ , ( $\triangle$ )  $0.2 \text{ g L}^{-1}$ , ( $\blacktriangle$ )  $0.3 \text{ g L}^{-1}$ . The insert figure is the dependence of initial activity on catalyst dosage.




Fig. 2S XRD patterns of g-C<sub>3</sub>N<sub>4</sub> and Pd/g-C<sub>3</sub>N<sub>4</sub> catalysts.




Fig. 3S XPS profiles of Pd 3d spectra on used catalyst 3. Black line refers to the experimental data; Red line refers to the fitting curve; Blue line refers to Pd<sup>0</sup>; Green line refers to Pd<sup>2+</sup>.




Fig. 4S XPS profiles of Pd 3d spectra on catalyst 7, 8 and 9. Black line refers to the experimental data; Red line refers to the fitting curve; Blue line refers to Pd<sup>0</sup>; Green line refers to Pd<sup>2+</sup>.

