

Supporting Information

Biocatalytic green approach for epoxidation of fatty compounds derived from soyadeodistillate under acid free conditions

Sunil Kumar Suman,^{1a} Manisha Dhawaria,^{1a} Jyoti Porwal,^{1a} Mounika Aila,^{1a} Neha Karanwal,^{1a} Babita Behra,^{1b} Savita Kaul,^{1a} Sanjay Ghosh² and Suman L. Jain^{1c*}

^{1a}Biofuel Division, CSIR-Indian Institute of Petroleum, Dehradun-248005 (India)

^{1b}Analytical Sciences Division, CSIR-Indian Institute of Petroleum, Dehradun-248005 (India)

^{1c}Chemical Sciences Division, CSIR-Indian Institute of Petroleum, Dehradun-248005 (India); Tel. 91-135-2525788; *Email: suman@iip.res.in

²Department of Biotechnology, Indian Institute of Technology Roorkee-247667, India

Fig. S 1. FTIR spectra of: A) Linoleic acid; B) Epoxide of Linoleic acid




Fig. S 2. ^1H NMR of (A) linoleic acid, (B) epoxide of linoleic acid




Fig. S 3. ¹H NMR of (A) soyadeodistillate, (B) epoxide of soyadeodistillate, (C) Ester of soyadeodistillate [(9Z,12Z)-2-ethylhexyl octadeca-9,12-dienoate], (D) Epoxide of ester of soyadeodistillate [(9Z,12Z)-2-ethylhexyl octadeca-9,12-dienoate]

