Relationship between measurement conditions and energy levels in the organic dyes used in dye-sensitized solar cells

Valentas Gaidelis^a, Egidijus Kamarauskas^a, Tadas Malinauskas^b, Vytautas Getautis^b, Robert Send^c, Henrike Wonneberger^c, and Ingmar Bruder^c

^aDepartment of Solid State Electronics, Vilnius University, Saulėtekio 9, LT-10222 Vilnius, Lithuania

^bDepartment of Organic Chemistry, Kaunas University of Technology, Radvilėnų pl. 19, Kaunas, 50254, Lithuania

^ctrinamiX GmbH, Carl-Bosch-Str.38, Ludwigshafen, 67056, Germany

July 9, 2015

Supplementrary Materials



Figure S1: IPCE and IPCE of dye D1



Figure S2: Absorbance spectra of dye D1



Figure S3: IPCE and IPCE of dye D2



Figure S4: Absorbance and IPCE of dye D2



Figure S5: Absorbance and IPCE of dye D3



Figure S6: IPCE and IPCE of dye D3



Figure S7: Absorbance and IPCE of dye D4



Figure S8: IPCE and IPCE of dye D4



Figure S9: Absorbance and IPCE of dye D5



Figure S10: IPCE and IPCE of dye D5



Figure S11: Absorbance spectra of dye D6



Figure S12: IPCE and IPCE of dye D7



Figure S13: Absorbance and IPCE of dye D8



Figure S14: IPCE and IPCE of dye D8



Figure S15: IPCE and IPCE of dye D9



Figure S16: Absorbance spectra of dye D9