

Supporting Information

Synthesis and characterization of cyclic 9,9-bis(4-hydroxyphenyl)fluorine (phenylene phosphonate) oligomer and its flame retardancy application

Nana Li ^{a,b}, Guowei Jiang ^a, Guangyuan Zhou ^{*,a}

^a Key Laboratory of Polymer Ecomaterials, Changchun Institute of Applied Chemistry, Chinese Academy of Sciences, Changchun, Jilin, 130022, China.

^b University of Chinese Academy of Sciences, Beijing, 100049, China.

Author: Nana Li

E-mail: lnnyy1314@163.com

Author: Guowei Jiang

E-mail: gwjiang@ciac.ac.cn

*Corresponding author: Guangyuan Zhou

E-mail: gyzhou@ciac.ac.cn

Phone number: +86-0431-85262204

Fax number: +86-0431-85685653

Key words: Cyclic 9,9-bis(4-hydroxyphenyl)fluorine (phenylene phosphonate), polybutylene terephthalate, flame retardancy, TGA, FTIR, pyrolysis/GC/MS




Fig. S1 ^1H NMR spectra of CPFP oligomer.

^1H NMR (DMSO- d_6 , ppm): δ 6.62 (d, 4H), 6.88(m, 4H), 7.28(m, 2H), 7.36(m, 4H), 7.58(m, 2H), 7.69(m, 1H), 7.88(m, 2H), 7.97(d, 2H) ppm.